

Document downloaded from:

<http://hdl.handle.net/10251/83602>

This paper must be cited as:

Conejero Casares, JA.; Lizama, C.; Murillo Arcila, M. (2016). On the existence of chaos for the viscous van Wijngaarden-Eringen equation. *Chaos, Solitons and Fractals*. 89:100-104. doi:10.1016/j.chaos.2015.10.009.



The final publication is available at

<http://dx.doi.org/10.1016/j.chaos.2015.10.009>

Copyright Elsevier

Additional Information

This is the author's version of a work that was accepted for publication in *Chaos, Solitons and Fractals*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Chaos, Solitons and Fractals* 89 (2016) 100–104. DOI 10.1016/j.chaos.2015.10.009.

# On the existence of chaos for the viscous van Wijngaarden–Eringen equation<sup>☆</sup>

J. Alberto Conejero

*Instituto Universitario de Matemática Pura y Aplicada.  
Universitat Politècnica de València, València, Spain*

Carlos Lizama

*Departamento de Matemática y Ciencia de la Computación.  
Universidad de Santiago de Chile,  
Las Sophoras 173, Estación Central, Santiago, Chile.*

Marina Murillo-Arcila

*Instituto Universitario de Matemática Pura y Aplicada.  
Universitat Politècnica de València, València, Spain*

---

## Abstract

We study the viscous van Wijngaarden–Eringen equation:

$$\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = (\text{Re}_d)^{-1} \frac{\partial^3 u}{\partial t \partial x^2} + a_0^2 \frac{\partial^4 u}{\partial t^2 \partial x^2} \quad (1)$$

which corresponds to the linearized version of the equation that models the acoustic planar propagation in bubbly liquids. We show the existence of an explicit range, solely in terms of the constants  $a_0$  and  $\text{Re}_d$ , in which we can ensure that this equation admits a uniformly continuous, Devaney chaotic and topologically mixing semigroup on Herzog's type Banach spaces.

*Keywords:* Acoustics,  $C_0$ -semigroups, bubble liquids, Devaney chaos, hypercyclicity, van Wijngaarden–Eringen equation, wave propagation, pressure waves.

---

## 1. Introduction

In the 1940's and 50's, the interest in studying the propagation of pressure waves of small amplitude in bubbly liquids appeared. The reason was to determine whether it was possible to take advantage of these acoustical properties to control the sound produced by propellers, both of surface ships and submerged ship. A vast literature on the subject deals with theoretical and experimental studies of the various aspects of propagation of pressure waves of small amplitude in bubbly liquids, see for instance [32].

---

<sup>☆</sup>The first and third authors are supported in part by MEC Project MTM2013-47093-P. The second author is partially supported by CONICYT, under Fondecyt Grant number 1140258.

*Email addresses:* [aconejero@upv.es](mailto:aconejero@upv.es) (J. Alberto Conejero), [carlos.lizama@usach.cl](mailto:carlos.lizama@usach.cl) (Carlos Lizama), [mamuar1@posgrado.upv.es](mailto:mamuar1@posgrado.upv.es) (Marina Murillo-Arcila)

The acoustic planar propagation perpendicular to and along the x-axis (i.e., 1D flow) in bubbly liquids is given by the following equation [19, Eq. 4.14]

$$\frac{\partial^2 u}{\partial t^2} - (1 - 2\varepsilon(\beta - 1)) \frac{\partial^2 u}{\partial x^2} + \varepsilon \frac{\partial^2}{\partial t^2} \left( \frac{\partial^2 u}{\partial x^2} \right)^2 = (\text{Re}_d)^{-1} \frac{\partial^3 u}{\partial t \partial x^2} + a_0^2 \frac{\partial^4 u}{\partial t^2 \partial x^2}, \quad t \geq 0, \quad x \in \mathbb{R}, \quad (2)$$

being  $\text{Re}_d = c_e L / \delta$  a *Reynolds number*, where  $c_e (> 0)$  is the adiabatic sound speed,  $\delta$  is the diffusivity of sound [31], and  $L$  is a characteristic (macroscopic length). The constant  $a_0$  is a *Knudsen number* that corresponds to the dimensionless bubble radius. In addition,  $\gamma$  stands for the adiabatic index of the liquid and  $\beta (> 1)$  is known as the *coefficient of nonlinearity* [5]. This coefficient is given by  $\beta = (\gamma + 1)/2$  in the case of a perfect gas.

More details on the formulation of equation (2) can be found in [19] and [18]. The linearized version,  $\varepsilon = 0$ , of equation (2) is known as the viscous (or dissipative) van Wijngaarden–Eringen equation, see [32, 14].

$$\frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} = (\text{Re}_d)^{-1} \frac{\partial^3 u}{\partial t \partial x^2} + a_0^2 \frac{\partial^4 u}{\partial t^2 \partial x^2}, \quad t \geq 0, \quad x \in \mathbb{R}, \quad (3)$$

Some other classical models in nonlinear acoustics are the Kuznetsov equation, the Westervelt equation, and the Kokhlov-Zabolotskaya-Kuznetsov equation. Several initial boundary problems for these second order in time partial differential equations have been already solved (see for instance [20], [21], [27] and [28]).

Many problems in acoustics for the sound propagation are described in terms of the linear wave equation. The transport equation governing the propagation of second-sound in both metals and dielectric solids is the well known damped wave equation [6, 24] (see also [17] for recent advances in the area). For these two cases, the 1D version of this PDE coincides with the hyperbolic heat transfer equation. The chaotic behaviour of the solutions of the abstract Cauchy problem stated with this equation was investigated in [9], see also [16]. For high wave amplitudes and intensities, phenomena such as wave distortion and formation of shocks appear, and then it is natural to study chaos for this type of equations. In [8] the chaotic behaviour of the one-dimensional version of the Moore-Gibson-Thompson equation is studied. See also the seminal formulation of this equation in [30, Eq. 7]. Moreover, the strong connection between acoustics and bioengineering and industry of high intensity sound waves has contributed to the improvement of the research in this area (see [10], [22], [7]).

To the best of our knowledge, no study on chaotic behavior for equation (1) has been carried on. Hence, our aim here is to examine the van Wijngaarden–Eringen equation in the context of a dynamic, yet still analytically tractable, setting.

In this paper, we succeed in proving the existence of a chaotic dynamics for the van Wijngaarden–Eringen equation. More precisely, we are able to show that whenever  $a_0 < 1$  and

$$\frac{\sqrt{5}}{6} < a_0 \text{Re}_d < \frac{1}{2}, \quad (4)$$

then equation (1) admits a uniformly continuous semigroup which is Devaney chaotic on an isomorphic copy of the sequence space  $c_0^2(\mathbb{N}_0)$ .

The paper is organized as follows. In section 2 we introduce some basic concepts related to the study of  $C_0$ -semigroups and chaos. In Section 3, chaos is also studied for the viscous van Wijngaarden–Eringen equation. Moreover, we provide a range for the bubble radius that ensures the existence of chaotic behaviour for the solutions of this equation. Finally, in Section 4, we give some physical interpretation of the results proved previously.

## 2. Preliminaries

Let  $X$  be a separable infinite-dimensional Banach space. We recall that  $\{T_t\}_{t \geq 0}$ , with  $T_t : X \rightarrow X$  a continuous and linear map on  $X$  for each  $t \geq 0$ , is a  $C_0$ -semigroup if  $T_0 = I$ ,  $T_{t+s} = T_t \circ T_s$  and  $\lim_{s \rightarrow t} T_s x = T_t x$  for all  $x \in X$  and  $t \geq 0$ .

Let  $\{T_t\}_{t \geq 0}$  an arbitrary  $C_0$ -semigroup on  $X$ . It can be shown that

$$Ax := \lim_{t \rightarrow 0} \frac{1}{t}(T_t x - x), \quad (5)$$

exists on a dense subspace of  $X$ ; the set of these  $x$ , the domain of  $A$ , is denoted by  $D(A)$ . Then  $A$ , or rather  $(A, D(A))$ , is called the *infinitesimal generator of the semigroup*.

Conversely, the Hille-Yosida theorem states that, under suitable assumption on a densely defined operator  $(A, D(A))$ , the solutions of the following abstract Cauchy problem on  $X$ :

$$\begin{cases} u_t(t, x) = Au(t, x), \\ u(0, x) = \varphi(x) \quad \varphi \in X, \end{cases} \quad (6)$$

can be represented by a  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  on  $X$  whose infinitesimal generator is  $A$ . If  $A \in L(X)$ , then the operators in the  $C_0$ -semigroup can be represented as  $T_t = e^{tA} = \sum_{k=0}^{\infty} (tA)^k / k!$  for all  $t \geq 0$  (see for instance [13, Ch. I, Prop. 3.5]).

A  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  on  $X$  is said to be *hypercyclic* if there exists  $x \in X$  such that the set  $\{T_t x : t \geq 0\}$  is dense in  $X$ . An element  $x \in X$  is called a *periodic point* for the semigroup  $\{T_t\}_{t \geq 0}$  if there exists some  $t > 0$  such that  $T_t x = x$ . A  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  is *topologically mixing* if for any pair  $U, V$  of nonempty open sets of  $X$ , there exists some  $t_0 \geq 0$  such that  $T_t(U) \cap V \neq \emptyset$  for all  $t \geq t_0$ . A  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  is called *Devaney chaotic* if it is hypercyclic and the set of periodic points is dense in  $X$ . We point out that these two requirements also yield the sensitive dependence on the initial conditions, as it was seen by Banks et al [1, 16]. Further information on the dynamics of  $C_0$ -semigroups can be found in [16, Ch. 7]. See also [12] for information regarding the stability properties of  $C_0$ -semigroups.

Another variation of the definition of chaos is the notion of *distributional chaos* introduced by Schweizer and Smítal [29], see also [23, 26] for its presentation in the infinite-dimensional linear setting. A  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  on  $X$  is said to be *distributionally chaotic* if there exists an uncountable subset  $S \subset X$  and  $\delta > 0$  such that, for each pair of distinct points  $x, y \in S$  and for every  $\varepsilon > 0$ , we have  $\overline{\text{Dens}}(\{s \geq 0; \|T_s x - T_s y\| > \delta\}) = 1$  and  $\overline{\text{Dens}}(\{s \geq 0; \|T_s x - T_s y\| < \varepsilon\}) = 1$ , where  $\overline{\text{Dens}}$  stands for the upper density of a set of real positive numbers. The set  $S$  is called the scrambled set and the  $C_0$ -semigroup is said to be *densely distributionally chaotic* if  $S$  is dense on  $X$ .

Now, we present a criterion that ensures Devaney chaos for  $C_0$ -semigroups. It is a variation of the (DSW) criterion [11] which depends on verifying that the point spectrum of the infinitesimal generator of the  $C_0$ -semigroup contains “enough” eigenvalues. A first criterion stated in these terms was given for operators by Godefroy and Shapiro in [15]. We will use the following version of the (DSW) Criterion, see [16, Th. 7.30]. It is also well known that distributional chaos holds whenever the DSW criterion can be applied [2, 4].

**Theorem 2.1.** *Let  $X$  be a complex separable Banach space, and  $\{T_t\}_{t \geq 0}$  a  $C_0$ -semigroup on  $X$  with infinitesimal generator  $(A, D(A))$ . Assume that there exists an open connected subset  $U$  and a weakly holomorphic function  $f : U \rightarrow X$ , such that*

- (1)  $U \cap i\mathbb{R} \neq \emptyset$ ,
- (2)  $f(\lambda) \in \ker(\lambda I - A)$  for every  $\lambda \in U$ ,
- (3) for any  $x^* \in X^*$ , if  $\langle f(\lambda), x^* \rangle = 0$  for all  $\lambda \in U$ , then  $x^* = 0$ .

*Then the semigroup  $\{T_t\}_{t \geq 0}$  is Devaney chaotic and topologically mixing.*

A Borel probability measure  $(\mu, \mathfrak{B})$  is said to have *full support* if for all non-empty open set  $U \subset X$  we have  $\mu(U) > 0$ , and  $\mu$  is said to be  *$T_t$ -invariant* if for all  $A \in \mathfrak{B}$  we have that  $\mu(A) = \mu(T_t^{-1}(A))$  for all  $t \geq 0$ . A  $C_0$ -semigroup  $\{T_t\}_{t \geq 0}$  is  *$\mu$ -strongly mixing* if  $\lim_{t \rightarrow \infty} \mu(A \cap T_t^{-1}(B)) = \mu(A)\mu(B)$ , for any  $A, B$  borelian sets of  $X$ . The existence of these measures are ensured if hypothesis of the DSW criterion are satisfied [25].

Finally, we recall the definition of the space of analytic functions of Herzog type. Given  $\rho > 0$ , let:

$$X_\rho = \left\{ f : \mathbb{R} \rightarrow \mathbb{C}; f(x) = \sum_{n=0}^{\infty} \frac{a_n \rho^n}{n!} x^n, (a_n)_{n \geq 0} \in c_0(\mathbb{N}_0) \right\}$$

endowed with the norm  $\|f\|_\rho = \sup_{n \geq 0} |a_n|$ . This space is isometrically isomorphic to  $c_0(\mathbb{N}_0)$ . For examples and references on Herzog spaces, we refer the reader to [8].

### 3. Existence of chaotic behavior

In this section, we will study the chaotic behavior of the viscous van Wijngaarden–Eringen equation

$$\frac{\partial^2 u}{\partial t^2}(t, x) - \frac{\partial^2 u}{\partial x^2}(t, x) = (\text{Re}_b)^{-1} \frac{\partial^3 u}{\partial t \partial x^2} u(t, x) + a_0^2 \frac{\partial^4 u}{\partial t^2 \partial x^2}(t, x), \quad t \geq 0, \quad x \in \mathbb{R}, \quad (7)$$

in the space of analytic functions of Herzog type. We recall that  $a_0 > 0$  denotes the dimensionless bubble radius and  $\text{Re}_b$  is a Reynolds number. Since the second order differential operator  $\partial_{xx}$  turns out to be a bounded operator on  $X_\rho$ , assuming the condition

$$a_0 < 1, \quad (8)$$

we obtain  $\|a_0^2 \partial_{xx}\|_\rho < 1$  and, consequently, the inverse operator  $(1 - a_0^2 \partial_{xx})^{-1}$  exists on  $X_\rho$ . Then we can express (7) as a first order equation on the product space  $X := X_\rho \oplus X_\rho$ . Setting  $u_1 = u$  and  $u_2 = \frac{\partial u}{\partial t}$  we can pose the following abstract Cauchy problem:

$$\begin{cases} \frac{\partial}{\partial t} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 & I \\ (1 - a_0^2 \partial_{xx})^{-1} \partial_{xx} & (\text{Re}_b)^{-1} (1 - a_0^2 \partial_{xx})^{-1} \partial_{xx} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}; \\ \begin{pmatrix} u_1(0, x) \\ u_2(0, x) \end{pmatrix} = \begin{pmatrix} \varphi_1(x) \\ \varphi_2(x) \end{pmatrix}, \quad x \in \mathbb{R}. \end{cases} \quad (9)$$

Then, the operator-valued matrix

$$A := \begin{pmatrix} 0 & I \\ -(1 - a_0^2 \partial_{xx})^{-1} \partial_{xx} & (\text{Re}_b)^{-1} (1 - a_0^2 \partial_{xx})^{-1} \partial_{xx} \end{pmatrix} \quad (10)$$

defines a bounded operator on  $X$  and, consequently, we have that  $\{e^{tA}\}_{t \geq 0}$  is the solution  $C_0$ -semigroup of (9). See also [9, Sec. 2.2] for a similar representation of the solution in the case of the wave equation, and [8, Sec. 3] for the case of the Moore-Gibson-Thompson equation.

The following theorem is the main result in this paper.

**Theorem 3.1.** *Suppose that  $a_0 < 1$  and*

$$0.3726 \approx \frac{\sqrt{5}}{6} < a_0 \text{Re}_b < \frac{1}{2}, \quad (11)$$

*then for each  $\rho$  satisfying*

$$\rho > \frac{r_0}{\left(\frac{1}{2a_0^2 \text{Re}_b} - 3r_0\right)a_0}, \quad (12)$$

*where  $r_0 := \frac{1}{2} \sqrt{\frac{1 - 4a_0^2 \text{Re}_b^2}{2a_0^2 \text{Re}_b}}$ , the operator  $A$  generates a uniformly continuous semigroup which is Devaney and distributionally chaotic, topologically mixing and admits a strongly mixing measure with full support on  $X_\rho \oplus X_\rho$ .*

*Proof.* Fix an arbitrary  $\rho > 0$  satisfying (12). Our purpose is to apply Theorem 2.1. Firstly, we define

$$U := \{z \in \mathbb{C} : |z| < r_0\}. \quad (13)$$

Note that  $1 - 4a_0^2 \operatorname{Re}_b^2 > 0$  due to the second inequality in (11). Hence  $r_0 > 0$  and  $U \cap i\mathbb{R} \neq \emptyset$ . This proves condition (1) in Theorem 2.1.

Secondly, for each  $\lambda \in U$  we define

$$R_\lambda := \frac{\lambda^2}{1 + (\operatorname{Re}_b)^{-1}\lambda + a_0^2\lambda^2}, \quad (14)$$

and weakly analytic functions  $f_{z_0, z_1} : U \rightarrow X_\rho \oplus X_\rho$  by

$$f_{z_0, z_1}(\lambda) := \begin{pmatrix} \varphi_\lambda \\ \lambda \varphi_\lambda \end{pmatrix}, \quad (15)$$

where  $\varphi_\lambda(x) := z_0 \cosh(\sqrt{R_\lambda}x) + z_1 \sinh(\sqrt{R_\lambda}x)$ , with  $z_0, z_1 \in \mathbb{C}$ .

It is easy to verify that

$$\varphi_\lambda''(x) + (\operatorname{Re}_b)^{-1}\lambda\varphi_\lambda''(x) + a_0^2\lambda^2\varphi_\lambda''(x) = \lambda^2\varphi_\lambda(x), \quad x \in \mathbb{R}, \quad (16)$$

and therefore  $Af_{z_0, z_1}(\lambda) = \lambda f_{z_0, z_1}(\lambda)$ . We will show that  $f_{z_0, z_1}(\lambda) \in X_\rho \oplus X_\rho$  for all  $\lambda \in U$ . Indeed, first note that we can rewrite  $\varphi_\lambda$  as follows:

$$\varphi_\lambda(x) = \cosh\left(\rho x \sqrt{\frac{R_\lambda}{\rho^2}}\right) z_0 + \sinh\left(\rho x \sqrt{\frac{R_\lambda}{\rho^2}}\right) z_1 = \sum_{n=0}^{\infty} a_n(\lambda) \frac{(\rho x)^n}{n!}, \quad x \in \mathbb{R}, \quad (17)$$

where  $a_n(\lambda) = z_0 \frac{R_\lambda^{n/2}}{\rho^n}$ ,  $n = 0, 2, 4, \dots$  and  $a_n(\lambda) = z_1 \sqrt{R_\lambda} \frac{R_\lambda^{(n-1)/2}}{\rho^n}$ ,  $n = 1, 3, 5, \dots$

Therefore, by definition, it is enough to prove that  $|\frac{R_\lambda}{\rho^2}| < 1$ . Indeed, observe that

$$a_0^2\lambda^2 + (\operatorname{Re}_b)^{-1}\lambda + 1 = a_0^2 \left[ \left( \lambda + \frac{1}{2a_0^2 \operatorname{Re}_b} \right) + 2r_0 \right] \left[ \left( \lambda + \frac{1}{2a_0^2 \operatorname{Re}_b} \right) - 2r_0 \right]. \quad (18)$$

Define  $\alpha := \frac{1}{2a_0^2 \operatorname{Re}_b} + 2r_0$  and  $\beta := \frac{1}{2a_0^2 \operatorname{Re}_b} - 2r_0$ . Then  $0 < \beta < \alpha$  and for all  $\lambda \in U$  we have

$$|\lambda + \alpha| \geq \alpha - |\lambda| \geq \beta - r_0 \quad \text{and} \quad |\lambda + \beta| \geq \beta - |\lambda| \geq \beta - r_0. \quad (19)$$

The hypothesis  $\sqrt{5}/6 < a_0 \operatorname{Re}_b$  implies that  $\frac{1}{2a_0^2 \operatorname{Re}_b} - 3r_0 > 0$ . Therefore  $\rho > 0$  and a calculation gives

$$\beta - r_0 > \frac{r_0}{\rho a_0}. \quad (20)$$

The above considerations imply,

$$\left| \frac{R_\lambda}{\rho^2} \right| = \frac{|\lambda|^2}{\rho^2 a_0^2 |\lambda + \alpha| |\lambda + \beta|} < r_0^2 \frac{1}{\rho^2 a_0^2} \frac{\rho^2 a_0^2}{r_0^2} = 1,$$

for all  $\lambda \in U$ , proving the claim. It proves condition (2) in Theorem 2.1.

It only remains to show that for any  $x^* \in X_\rho^* \oplus X_\rho^*$  the functions  $\lambda \rightarrow \langle f_{z_0, z_1}(\lambda), x^* \rangle$ ,  $z_0, z_1 \in \mathbb{C}$ , are holomorphic on  $U$ , and if they all vanish on  $U$ , then  $x^* = 0$ . Since  $X_\rho$  is isometrically isomorphic to  $c_0$ , in what follows, we identify the dual space  $X_\rho^*$  with  $\ell^1$ .

Let  $x^* \in X_\rho^* \oplus X_\rho^*$ . It can be represented in a canonical way by  $x^* = (x_1^*, x_2^*) = ((x_{1,n}^*)_{n \geq 0}, (x_{2,n}^*)_{n \geq 0}) \in \ell^1 \oplus \ell^1$ . Then, we have

$$0 = \langle f_{z_0, z_1}(\lambda), x^* \rangle = \langle \varphi_\lambda, x_1^* \rangle + \langle \lambda \varphi_\lambda, x_2^* \rangle, \quad (21)$$

for all  $\lambda \in U, z_0, z_1 \in \mathbb{C}$ . This last equation can be reformulated in the following way:

$$\begin{aligned} 0 &= \sum_{n=0}^{\infty} a_n(\lambda) x_{1,n}^* + \lambda \sum_{n=0}^{\infty} a_n(\lambda) x_{2,n}^* \\ &= z_0 x_{1,0}^* + \lambda z_0 x_{2,0}^* + \frac{z_1}{\rho} \sqrt{R_\lambda} x_{1,1}^* + \frac{z_1}{\rho} \lambda \sqrt{R_\lambda} x_{2,1}^* + \frac{z_0}{\rho^2} R_\lambda x_{1,2}^* + \frac{z_0}{\rho^2} R_\lambda \lambda x_{2,2}^* + \dots \end{aligned} \quad (22)$$

Let  $\lambda_0 = 0$ . It is clear that  $\lambda_0 \in U$  and  $R_{\lambda_0} = 0$ . Evaluating (22) in  $\lambda_0$ , we have the following equation:

$$z_0 x_{1,0}^* = 0 \quad (23)$$

for all  $z_0 \in \mathbb{C}$ . Therefore,  $x_{1,0}^* = 0$ .

Now, we divide (22) by  $\lambda$  and we get:

$$\begin{aligned} 0 &= \frac{1}{\lambda} \left( \sum_{n=0}^{\infty} a_n(\lambda) x_{1,n}^* + \lambda \sum_{n=0}^{\infty} a_n(\lambda) x_{2,n}^* \right) \\ &= z_0 x_{2,0}^* + \frac{z_1}{\rho} \frac{\sqrt{R_\lambda}}{\lambda} x_{1,1}^* + \frac{z_1}{\rho} \sqrt{R_\lambda} x_{2,1}^* + \frac{z_0}{\rho^2} \frac{R_\lambda}{\lambda} x_{1,2}^* + \frac{z_0}{\rho^2} R_\lambda x_{2,2}^* + \dots \\ &= z_0 x_{2,0}^* + \frac{z_1}{\rho} \frac{1}{\sqrt{1 + (\operatorname{Re} b)^{-1} \lambda + a_0^2 \lambda^2}} x_{1,1}^* + \frac{z_1}{\rho} \sqrt{R_\lambda} x_{2,1}^* \\ &\quad + \frac{z_0}{\rho^2} \frac{\lambda}{(1 + (\operatorname{Re} b)^{-1} \lambda + a_0^2 \lambda^2)} x_{1,2}^* + \frac{z_0}{\rho^2} R_\lambda x_{2,2}^* + \dots \end{aligned} \quad (24)$$

As  $R_{\lambda_0} = 0$ , evaluating (24) in  $\lambda_0$  we obtain:

$$z_0 x_{2,0}^* + \frac{z_1}{\rho} x_{1,1}^* = 0. \quad (25)$$

for all  $z_0, z_1 \in \mathbb{C}$ . Then,  $x_{2,0}^* = 0$  and  $x_{1,1}^* = 0$ .

Now, we divide (22) by  $\lambda \sqrt{R_\lambda}$  and we get:

$$0 = \frac{1}{\lambda \sqrt{R_\lambda}} \left( \sum_{n=0}^{\infty} a_n(\lambda) x_{1,n}^* + \lambda \sum_{n=0}^{\infty} a_n(\lambda) x_{2,n}^* \right). \quad (26)$$

So that equation (26) can be reduced to:

$$\begin{aligned} 0 &= \frac{z_1}{\rho} x_{2,1}^* + \frac{z_0}{\rho^2} \frac{\sqrt{R_\lambda}}{\lambda} x_{1,2}^* + \frac{z_0}{\rho^2} \sqrt{R_\lambda} x_{2,2}^* + \dots \\ &= \frac{z_1}{\rho} x_{2,1}^* + \frac{z_0}{\rho^2} \frac{1}{\sqrt{1 + (\operatorname{Re} b)^{-1} \lambda + a_0^2 \lambda^2}} x_{1,2}^* + \frac{z_0}{\rho^2} \frac{\lambda}{\sqrt{1 + (\operatorname{Re} b)^{-1} \lambda + a_0^2 \lambda^2}} x_{2,2}^* + \dots \end{aligned} \quad (27)$$

Evaluating (27) in  $\lambda_0$ , we have:

$$\frac{z_1}{\rho} x_{2,1}^* + \frac{z_0}{\rho^2} x_{1,2}^* = 0 \quad (28)$$

for all  $z_0, z_1 \in \mathbb{C}$ . Therefore,  $x_{2,1}^* = 0$  and  $x_{1,2}^* = 0$ .

Proceeding inductively, we will get that  $x_{i,n}^* = 0$  for  $i = 1, 2$  and  $n \in \mathbb{N}$ . We finally have  $x^* = 0$  and we conclude the result by applying Theorem 2.1.  $\square$

*Remark 3.2.* Recalling that  $a_0 > 0$  denotes the bubble radius and  $\text{Re}_b = c_e L / \delta$  is a Reynolds number, where  $L$  denote a characteristic length,  $c_e$  the adiabatic sound speed and the positive constant  $\delta > 0$  is known as the diffusivity of sound, we observe that the condition (11), namely

$$0.3726 \approx \frac{\sqrt{5}}{6} < a_0 \text{Re}_b < \frac{1}{2}, \quad (29)$$

together with the general condition  $a_0 < 1$ , give an explicit range for the bubble radius in order to obtain chaotic behavior in the given model.

This estimation can be compared with the physical one of

$$0.4899 \approx \frac{\sqrt{6}}{5} < a_0 \text{Re}_b, \quad (30)$$

given in [19, Eq. 4.20] that has to be satisfied by the Reynolds number.

#### 4. Conclusions

These results can be understood in the following sense. For the hypercyclicity, given an arbitrary acoustic planar wave close to the origin one can determine a wave far away enough in order that, as time goes by, we can have that its propagation through a bubbly liquid gives us a wave at the origin as close as we want to the initial one. The Devaney chaos also yields the existence of periodic waves similar as much as we want to a prescribed one close to the origin.

These theoretical results present some limitations: The amplitudes required to resemble a prescribed wave at the origin can be so wide that cannot even be easily generated.

Furthermore, we recall that the existence of Devaney chaos yields the sensitive dependence of the solutions to the problem respect to the initial conditions. This affirms that given some planar wave, one can find a small perturbation, such that after a long enough period of time the behavior of both waves is completely different.

One last comment, the existence of distributional chaos asserts, roughly speaking, that there is an uncountable set of initial conditions such that we can pick a pair of initial conditions from this set and, as time goes by, there will be long time intervals in which the behaviour of the waves close to the origin are very similar for both initial waves. On the other hand, there will be also intervals as long as the previous ones in which the waves are quite different depending on which one of these two initial conditions we have chosen, see [3].

#### 5. Acknowledgements

We want to thank Pedro M. Jordan for proposing us to conduct this research.

#### References

- [1] J. Banks, J. Brooks, G. Cairns, G. Davis, and P. Stacey. On Devaney's definition of chaos. *Amer. Math. Monthly*, 99(4):332–334, 1992.
- [2] X. Barrachina and J. A. Conejero. Devaney chaos and distributional chaos in the solution of certain partial differential equations. *Abstr. Appl. Anal.*, pages Art. ID 457019, 11, 2012.
- [3] X. Barrachina, J. A. Conejero, M. Murillo-Arcila, and J. B. Seoane-Sepúlveda. Distributional chaos for the forward and backward control traffic model. *Linear Algebra Appl.*, 479:202–215, 2015.
- [4] T. Bermúdez, A. Bonilla, F. Martínez-Giménez, and A. Peris. Li-Yorke and distributionally chaotic operators. *J. Math. Anal. Appl.*, 373(1):83–93, 2011.
- [5] R. T. Beyer. *The parameter B=A, in Nonlinear Acoustics*. 25-39. Academic Press, San Diego, CA, 1997.
- [6] M. Chester. Second sound in solids. *Phys. Rev.*, 131:2013–2015, 1963.
- [7] F. Colouvrat. On the equations of nonlinear acoustics. *Journal de Acoustique*, 5:321–359, 1992.



- [8] J. A. Conejero, C. Lizama, and F. Rodenas. Chaotic behaviour of the solutions of the Moore-Gibson-Thompson equation. *Applied Math. & Inf. Sciences*, 9(5):2233–2238, 2015.
- [9] J. A. Conejero, A. Peris, and M. Trujillo. Chaotic asymptotic behavior of the hyperbolic heat transfer equation solutions. *Internat. J. Bifur. Chaos Appl. Sci. Engrg.*, 20(9):2943–2947, 2010.
- [10] D. Crighton. Model equations of nonlinear acoustics. *Ann. Rev. Fluid Mech.*, 11:11–33, 1979.
- [11] W. Desch, W. Schappacher, and G. F. Webb. Hypercyclic and chaotic semigroups of linear operators. *Ergodic Theory Dynam. Systems*, 17(4):793–819, 1997.
- [12] T. Eisner. *Stability of operators and operator semigroups*, volume 209 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2010.
- [13] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafuno, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.
- [14] A. C. Eringen. Theory of thermo-microstretch fluids and bubbly liquids. *International Journal of Engineering Science*, 28(2):133 – 143, 1990.
- [15] G. Godefroy and J. H. Shapiro. Operators with dense, invariant, cyclic vector manifolds. *J. Funct. Anal.*, 98(2):229–269, 1991.
- [16] K.-G. Grosse-Erdmann and A. Peris. *Linear chaos*. Universitext. Springer, London, 2011.
- [17] P. M. Jordan. Second-sound propagation in rigid, nonlinear conductors. *Mec. Res. Com.*, 68:52–59, 2015.
- [18] P. M. Jordan and C. Feuillede. On the propagation of transient of acoustic waves in isothermal bubbly liquids. *Physics Letters A*, 350(12):56–62, 2006.
- [19] P. M. Jordan, R. S. Keiffer, and G. Saccomandi. Anomalous propagation of acoustic traveling waves in thermoviscous fluids under the Rubin-Rosenau-Gottlieb theory of dispersive media. *Wave Motion*, 51(2):382–388, 2014.
- [20] B. Kaltenbacher and I. Lasiecka. Well-posedness of the Westervelt and the Kuznetsov equation with nonhomogeneous Neumann boundary conditions. *Discrete Contin. Dyn. Syst.*, (Dynamical systems, differential equations and applications. 8th AIMS Conference. Suppl. Vol. II):763–773, 2011.
- [21] B. Kaltenbacher, I. Lasiecka, and R. Marchand. Wellposedness and exponential decay rates for the Moore-Gibson-Thompson equation arising in high intensity ultrasound. *Control Cybernet.*, 40(4):971–988, 2011.
- [22] V. P. Kuznetso. Equations of nonlinear acoustics. *Soviet Physics Acoustics-USSR*, 16(4):467–470, 1971.
- [23] F. Martínez-Giménez, P. Oprocha, and A. Peris. Distributional chaos for backward shifts. *J. Math. Anal. Appl.*, 351(2):607–615, 2009.
- [24] M. Maurer. Relaxation model for heat conduction in metals. *J. Appl. Phys.*, 40:5123–5130, 1969.
- [25] M. Murillo-Arcila and A. Peris. Strong mixing measures for  $C_0$ -semigroups. *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM*, 109(1):101–115, 2015.
- [26] P. Oprocha. A quantum harmonic oscillator and strong chaos. *J. Phys. A*, 39(47):14559–14565, 2006.
- [27] A. Rozanova. The Khokhlov-Zabolotskaya-Kuznetsov equation. *C. R. Math. Acad. Sci. Paris*, 344(5):337–342, 2007.
- [28] A. Rozanova-Pierrat. Qualitative analysis of the Khokhlov-Zabolotskaya-Kuznetsov (KZK) equation. *Math. Models Methods Appl. Sci.*, 18(5):781–812, 2008.
- [29] B. Schweizer and J. Smítal. Measures of chaos and a spectral decomposition of dynamical systems on the interval. *Trans. Amer. Math. Soc.*, 344(2):737–754, 1994.
- [30] G. Stokes. XXXVIII. an examination of the possible effect of the radiation of heat on the propagation of sound. *Philosophical Magazine. Series 4*, 1(4):3605–317, 1851.
- [31] P. A. Thompson. *Compressible-Fluid Dynamics*. McGraw Hill, New York, NY, 1972.
- [32] L. V. Wijngaar. One-dimensional flow of liquids containing small gas bubbles. *Annual review of fluid Mechanics*, 4:369–396, 1972.