

Document downloaded from:

<http://hdl.handle.net/10251/115608>

This paper must be cited as:

Rodríguez Burruezo, A.; Gurrea-Ysasi, G.; Prohens Tomás, J.; Fita, A.; Cortés Olmos, C. (2018). In vitro germination and growth protocols of the ornamental *Lophophora williamsii* (Lem.) Coult. as a tool for protecting endangered wild populations. *Scientia Horticulturae*. 237:120-127. doi:10.1016/j.scienta.2018.03.064



The final publication is available at

<http://doi.org/10.1016/j.scienta.2018.03.064>

Copyright Elsevier

Additional Information

1 ***In vitro* germination and growth protocols of the ornamental**
2 ***Lophophora williamsii* (Lem.) Coult. as a tool for protecting endangered**
3 **wild populations**

4
5 Cortés-Olmos, C.; Gurrea-Ysasi, G.; Prohens, J.; Rodríguez-Burruezo, A.; Fita, A.*

6 *Instituto de Conservación y Mejora de la Agrodiversidad Valenciana (COMAV),*
7 *Universitat Politècnica de València. Camino Vera s/n, CP 46022, Valencia, Spain.*

8 * To whom correspondence should be addressed (anfifer@btc.upv.es)

9
10 **ABSTRACT**

11 *Lophophora williamsii* is an ornamental slow growth cactus highly appreciated by cacti
12 growers and hobbyists. Its demand is often satisfied through illegal collection of wild
13 plants and many populations are threatened with extinction. Thus, an efficient *in vitro*
14 protocol without plant growth regulators will be of great interest for conservation
15 purposes of this cactus. Eight different germination media, combining Murashige and
16 Skoog medium (MS, full and half-strength), sucrose (20 and 30 g L⁻¹) and agar (8 and 10
17 g L⁻¹), were used to study germination rate, number of seedlings with areoles and initial
18 seedling development. Germination rates among culture media only differed significantly
19 in the first 14 days after sowing (DAS), reaching 67-75% at the end of the assay (49
20 DAS). Remarkable interactions among media components were detected, and 20 g L⁻¹
21 sucrose and 8 g L⁻¹ agar combination gave the highest performance for both size and
22 number of areoles. Following germination assay, a growth assay was conducted during
23 105 days using three growth media (GrM) at different sucrose concentration (15, 30 and

24 45 g L⁻¹) to evaluate the increase in seedling size and number of areoles. Regardless of
25 their initial size, 15 g L⁻¹ sucrose provided the best results for both traits. Size increase
26 was higher in the 4-5 mm seedling group, while increase in areoles was greater in 2-3 mm
27 seedlings. It was possible to develop an *in vitro* protocol, in absence of plant growth
28 regulators, which allows maximizing *L. williamsii* germination and growth during its first
29 stages of development, which may increase the availability of plants in the market and
30 avoid exhaustion of wild populations. Furthermore, plants grown *ex situ* could be
31 reintroduced in endangered natural populations.

32

33 Keywords: peyote; Cactaceae; ex-situ conservation; threatened species; plant growth
34 regulator

35

36

37 **1. Introduction**

38 *Lophophora williamsii* (Lem.) Coult., commonly known as ‘peyote’, is a small (5-12 cm
39 in diameter) blue-green, button-like, spineless and slow growth cactus with napiform root
40 (Figure 1) whose wild populations are distributed in Mexican highlands and in the arid
41 regions of South-western United States (Anderson, 1996). This plant has been used during
42 centuries in several rituals and ceremonies by Indian Tribes (McLaughlin, 1973; Borchers
43 et al., 2000; Halpern et al., 2005) due to its content in alkaloids (of which the major one
44 is mescaline) with psychotropic activity (Casado et al., 2008). In addition to its ancestral
45 ethnobotanical use, *L. williamsii* has always aroused a great interest among cacti lovers
46 and collectors. In fact, its growing demand in the market (as in other Cactaceae) has been

47 often satisfied through illegal collection of wild individuals (Anderson et al., 1994; Sajeve
48 et al., 2013), in part due to the fact that its growth from seeds is very slow.

49 <Figure 1>

50 The plundering of wild plants, added to other problems related to human activity, such as
51 agricultural and urban expansions, introduction of exotic grasses, use of herbicides and
52 pesticides among others, has led to many cacti being threatened with extinction (Taylor
53 1997; Sanchez-Martinez et al., 2009), especially in those places close to urban areas
54 where access to natural populations is easier. This high pressure on ecosystems
55 compromises the viability of certain populations and could result in an irremediable loss
56 of unique genetic pools. In this way, genus *Lophophora* is protected by Mexican laws and
57 is included in the Convention on International Trade in Endangered Species of Fauna and
58 Flora (CITES) (Sajeve et al., 2013).

59 *In vitro* culture techniques may play a key role to accelerate the growth of *L. williamsii*
60 plants, especially after sowing from seeds and during early stages of the development,
61 when seedlings are more vulnerable. Furthermore, areoles are characteristic of Cactaceae,
62 equivalent to the buds of other higher plants, and include two points of growth: one that
63 leads to the thorns and another that originates the flowers and new buds (Ballester-
64 Olmos, 1997). In this sense, as active growth organs, the number of areoles must
65 be considered for *in vitro* protocols as they represent the capacity of each plant for a higher
66 multiplication. *In vitro* culture could help to obtain larger and flowering plants faster than
67 by conventional seed reproduction, satisfying the demand of the market and thus reducing
68 consequently the need to plunder wild plants. Also, *in vitro* culture could contribute to
69 the *ex situ* conservation of plants and populations with the aim to reintroduce them in
70 their habitat for restoring extinct or critically endangered natural populations.

71 *In vitro* plants propagation and micropropagation for conservation purposes requires
72 efficient methods. In some cases, it has been reported that medium composition and its
73 supplementation with plant growth regulators (PGRs) may alter morphological and
74 physiological characters, even genetic stability, in the obtained plants (Lema-Ruminska
75 and Kulus, 2014). Therefore, media without PGRs would be desirable as they are less
76 prone to induct of somaclonal variations on material collected from the wild. In addition,
77 media containing no PGRs are cheaper and easier to prepare than those formulations
78 including PGR. Thus, in this study we compared different *in vitro* culture media in
79 absence of PGRs, in order to detect those that maximize the development and growth of
80 *L. williamsii* seedlings during the early stages after sowing. This would also be the first
81 step to establish a *Lophophora* micropropagation protocol in absence of PGRs.

82

83 **2. Material and methods**

84 *2.1 Seed disinfection*

85 Seeds of a population of *L. williamsii* were kindly donated by Cactusloft (Valencia,
86 Spain). Three hundred and twenty seeds were disinfected for 1 min in 70% etanol (v/v),
87 followed by 25 min in 15% domestic bleach solution (v/v; 4% sodium hypochlorite),
88 supplemented with 0.08% Tween-20 (v/v) and rinsed 3 times in distilled sterilized water
89 under aseptic conditions under laminar flow cabinet conditions (model AH-100, Telstar,
90 Terrassa, Spain).

91

92 *2.2 Germination assay*

93 Disinfected seeds were sown on different germination media (Table 1) in Petri dishes (10
94 seeds per dish, 4 dishes per medium, n=4). A total of eight *in vitro* formulations were
95 evaluated in this assay, which included all combinations of Murashige and Skoog medium
96 (MS, at full-strength or half-strength, 1×MS or 1/2×MS, i.e. at original or half
97 concentration of commercial MS salts concentration, respectively), sucrose concentration
98 (at 20 or 30 g L⁻¹), and agar concentration (at 8 or 10 g L⁻¹). The pH of all media were
99 adjusted to 5.7 before autoclaving. Four replicates per medium were cultured for 49 days
100 in a growth chamber at 25 ± 2 °C, 16 h day length and photosynthetic photon flux of 50
101 μmol m⁻²s⁻¹.

102 <Table 1>

103 Germination ratio, number of seedlings with areoles, number of areoles per seedling, and
104 seedling size were recorded weekly. Germination was evaluated during 49 days after
105 sowing (DAS) of *in vitro* culture. For the rest of the traits (i.e. seedlings with areoles,
106 areoles per seedling, and seedling size), only seedlings germinated within the first 28
107 DAS were evaluated, as data from seedlings germinated later, due to their low size, could
108 bias the analysis. The germination viability and the number of seedlings with areoles were
109 measured and expressed in percentage. Data in these traits were transformed by arcsine
110 square root (Little and Hills, 1978), but differences between the results obtained by
111 transformed and non-transformed data were negligible. Thus results presented in this
112 work were performed on the original non-transformed data.

113 2.3 Plant growth assay

114 After 49 DAS (germination assay), 197 healthy plants from the germination assay were
115 subcultivated to analyze growth parameters. Seedlings were distributed in three groups
116 depending on their size and similar root development: i) seedlings with sizes between 2

117 and 3 mm (64 plants); ii) seedlings with sizes between 3 and 4 mm (100 plants); iii)
118 seedlings with sizes between 4 and 5 mm (33 plants). Each size group was divided in
119 three subgroups and each subgroup was cultivated in a different growth medium (GrM)
120 (Table 1) for 15 weeks, maintaining the same pH and culture conditions described in the
121 germination assay. Subculturing was performed monthly.

122 The number of areoles per seedling and seedling size were quantified by using a binocular
123 microscope and recorded weekly.

124

125 *2.4 Statistical Analysis*

126 Unifactorial and multifactorial (using MS concentration, sucrose, and agar as factors)
127 ANOVA analyses were performed using Statgraphics X64 (Statpoint Technologies, The
128 Plains, VA, USA). Culture media and media component effects were analysed in both
129 assays.

130

131 **3. Results and discussion**

132 *3.1 Germination Assay*

133 *3.1.1 Germination viability*

134 Significant differences were found among media in the first two weeks of the experiment
135 in terms of germination rate (Table 2). From 21 DAS on, germination values among
136 media did not differ statistically. Therefore, only the two first weeks were of interest to
137 study the effect of media formulation. As a whole, although differences were not
138 significant in all cases, our results suggest that most half-strength MS formulations may
139 enhance early germination, especially at 7 DAS. Thus, with the only exception of M8,

140 media M5, M7 and, particularly, M6 provided \geq germination rates at 7 DAS in
141 comparison to full-strength MS media (Table 2). Thus, M6 followed by M4, M5 and M7,
142 had the highest percentage of germination ($>10\%$) at 7 DAS, while M6, with 67% of
143 germination efficiency, was also found the best formulation at 14 DAS, followed to a
144 lesser extent by the rest of media (42-62%) (Table 2).. In this regard, full-strength MS is
145 capable to cause a one-week delay in germination compared to half-strength MS in some
146 species like *Pilosocereus robinii* and *Astrophytum asterias* (Quiala et al., 2009; Lema-
147 Ruminska and Kulus, 2012). Even in the case of immature embryos from other species
148 grown *in vitro*, like *Capsella* and *Capsicum*, it has been reported that full-strength MS
149 level could be slightly toxic for them and decrease their efficiency rates (Monnier, 1995;
150 Manzur et al., 2013).

151 <Table 2>

152 After 49 DAS, the percentage of germination observed ranged between 67.5% (M1 and
153 M7) to 77.5% (M8), similar to many other *in vitro* germination rates in cacti, as detected
154 in *Ariocarpus kotschoubeyanus* and *Cereus hildmannianus* (Moebius-Goldammer et al.,
155 2003; Langer and Mergener, 2013). Furthermore, our results revealed that germination
156 rates in the different media reached similar values (nonsignificant differences) at the end
157 of the experiment (Table 2). This fact indicates that the seed population used as a starting
158 material was homogeneous for this trait and differences in germination observed at the
159 earlier stages have been solely caused by the different media formulation.

160

161 3.1.2 Number of seedlings with areoles

162 There were no significant differences between MS levels (full-strength MS vs. half-
163 strength MS), sucrose levels (20 g L⁻¹ vs. 30 g L⁻¹) or agar levels (8 g L⁻¹ vs. 10 g L⁻¹)

164 along the experiment for the percentage of seedlings with areoles, which suggests that
165 there is no clear effect of each of these factors for this trait (Table 2). By contrast,
166 significant differences were found among the different media formulations, indicating a
167 considerable interaction among the components of the media (Table 2). In this regard,
168 media which enabled an earlier germination (i.e. M4, M5, M6 and M7) also favored an
169 earlier emergence of areoles until 14 DAS (Table 2).

170 Regarding the values corresponding to the most advanced stages (from 21 DAS on), we
171 found that M3 and particularly M5, both with agar concentration of 8 g L⁻¹, displayed the
172 highest percentages of plants with areoles (Table 2). On the contrary, media such as M1
173 and M7, also containing 8 g L⁻¹ of agar, were among the worst in the experiment. These
174 findings suggest the occurrence of strong interaction between MS and sucrose levels at 8
175 g L⁻¹ agar to explain such differences (Table 2). Thus, the simultaneous combination of
176 high levels of both MS and sucrose (M3) or alternatively low levels of both (M5) are
177 required for a good response on this trait, while the contrary is true for combinations of
178 high levels of MS and low levels of sucrose *or vice versa* in M1 and M7, respectively.

179 In addition, it should be noted that in general M4 and M8 displayed a lower efficiency
180 than the other media from 7 to 42 DAS, with significant differences in many cases,
181 particularly M4 (Table 2). Probably their high osmotic gradient, due to high sucrose and
182 agar levels, is the main reason for the poor performance observed in these formulations
183 (Pérez-Molphe-Balch *et al.*, 1998). In this respect, many authors have reported in a broad
184 range of species that isolated mature embryos cultivated *in vitro* had better response and
185 good growth in media at low sugar contents (Fisher and Neuhaus, 1995; Monnier, 1995;
186 Bhojwani and Razdan, 1996; Manzur *et al.*, 2013).

187 Finally, we found that from 35-42 DAS on, the percentage of plants with areoles did not
188 increase significantly in most of the media, reaching their highest values at this stage
189 (Table 2). This is correlated with the corresponding germination rates, which also reached
190 their highest values at 21-28 DAS, indicating that all germinated seedlings develop
191 areoles in all media about 2-3 weeks after germination.

192

193 *3.1.3 Initial seedling development*

194 The average size of seedlings germinated in M5 was greater than those observed in the
195 other media until 21 DAS. However, M1 offered the highest performance, even higher
196 than M5, in the later stages (Table 3). Also, for all the media studied most growth was
197 detected during the first 14 DAS. Therefore, M5 provided the fastest initial growth, while
198 M1 enabled a more sustained trend. In terms of number of areoles per seedling, a similar
199 behaviour was found. Thus, as a whole, M5 followed by M2, M6 and M7, resulted in a
200 higher number of areoles per seedling until 21 DAS (Table 3). However, from 28 DAS
201 to the end of the germination assay, M1 had the highest values, followed by M5. Both
202 media had the same levels of sucrose and agar, suggesting that the combination of sucrose
203 20 g L⁻¹ and agar 8 g L⁻¹ favors both traits.

204

<Table 3>

205 Considering the global effect of each factor, it seems that both sucrose and agar
206 concentration affected to a greater extent seedling growth and number of areoles per
207 seedling. In the case of the former, levels of 20 g L⁻¹ increased on average both traits,
208 particularly at the later stages, i.e. from 35 DAS on (Table 3). Considering the latter,
209 seedlings grown in media with agar concentration at 8 g L⁻¹ showed on average a faster
210 growth and number of areoles since the beginning of the assay, in comparison to those

211 observed in media with agar at 10 g L⁻¹. In fact, it was observed that media formulations
212 including agar at 10 g L⁻¹ hindered radicle penetration just after germination (Figure 2),
213 causing developmental delay with respect to those grown with 8 g L⁻¹. Furthermore, high
214 concentration of gelling agents may decrease the amount of available water (Pérez-
215 Molphe-Balch et al., 2002; Garcia-Osuna et al., 2011), affecting adversely seedling
216 development. These findings may also explain why media M4 and M8, with the highest
217 sucrose and agar levels, showed the poorest values in general, particularly in the most
218 advanced stages (i.e. after 28 DAS) (Table 3, Figure 3).

219 <Figure 2>

220 <Figure 3>

221 Finally, half-strength MS media appeared to be more adequate on average during the first
222 21 days, while no significant differences were found at more advanced stages (Table 3).
223 These results could explain the differences between M1 and M5, which only differed in
224 MS concentration. Thus, half-strength MS in M5 favoured early growth of seedlings,
225 while full-strength MS provided a better performance in the most advanced stages as was
226 observed in M1 (Figure 4). Therefore, similarly to germination efficiency, it looks like
227 that full-strength MS may have a negative interaction with plantlets at the earliest stages
228 of development, while it has no effect or even positive effect on later stages. Many studies
229 have reported that 1/2×MS, or even 1/4×MS, provides a better effect not only in
230 germination but also in subsequent early development than full MS (Gland-Zwenger,
231 1995; Xu et al., 2007; Manzur et al., 2013; Koné et al., 2015). This is probably due to a
232 deleterious effect of some of the salts present in the MS formulation (Monnier, 1995). By
233 contrast, full strength MS is advised for more advanced seedling stages as, once seedlings

234 increase their size and become photosynthetically active, their requirements in micro and
235 macronutrients are higher (Stewart and Kane, 2006; Paul et al., 2012).

236 <Figure 4>

237 *3.2 Growth assay*

238 As observed in the germination assay, seedlings from M1 showed a more prominent
239 development, largely due to the positive interaction between its factors. As the
240 concentration of agar was determinant and the presence of full-strength MS displayed
241 positive interactions, we decided to evaluate the influence of sucrose on the further
242 development of germinated seedlings. Thus, media based on full-strength MS and 8 g L⁻¹
243 of agar were supplemented with three different sucrose concentrations (Table 1), which
244 encompassed a range larger than initially studied in the germination assay, keeping 30 g
245 L⁻¹ as an intermediate concentration (i.e. 15, 30 and 45 g L⁻¹). Also, seedlings were
246 evaluated separately on the basis of their initial size/diameter: 2-3 mm, 3-4 mm and 4-5
247 mm.

248

249 *3.2.1 Seedling size*

250 There were no significant differences between the initial sizes of seedlings within each
251 size group. In fact, differences were ≤ 0.1 mm (Table 4). This confirms a precise selection
252 of seedlings for the growth assay as they were similar in size for the three treatments
253 within each group.

254 <Table 4>

255 In all size groups it was observed that seedlings grown in GrM-1 showed both a larger
256 final size and increase in size at the end of the experiment than those grown in GrM-2 or

257 GrM-3 after 105 days of cultivation (Table 4, Figure 5). These findings suggest that,
258 regardless the initial size of seedlings, relatively low levels of sucrose provided by GrM-
259 1 (15 g L⁻¹) are better for seedlings in terms of growth. By contrast, higher sucrose
260 concentrations (≥ 30 g L⁻¹) had comparatively a deleterious effect on growth (Table 4).
261 These results are in agreement with those from the germination assay, on which the early
262 growth of seedlings after germination was considerably higher under low sucrose levels
263 (20 g L⁻¹ vs. 30 g L⁻¹) (Table 3). Therefore, from germination on, *L. williamsii* seedlings
264 evolve better under low sucrose levels (15-20 g L⁻¹). Probably, excessive levels of sucrose
265 (>30 g L⁻¹) in some culture media caused unfavourable osmotic gradient that limits the
266 development of seedlings in this species. In this regard, several authors reported that only
267 early immature embryos require high levels of sucrose as source of energy and for
268 osmotic reasons, while both immature advanced and mature embryos (equivalent to our
269 *L. williamsii* seeds and seedlings) have a good response to moderate levels of sucrose in
270 terms of germination efficiency and further growth (Monnier, 1995; Bhojwani and
271 Razdan, 1996; Manzur et al., 2013).

272 <Figure 5>

273 Moreover, as a whole, we found that those seedlings with initial size of 4-5 mm had a
274 higher size increase than 2-3 mm and 3-4 mm seedlings (Table 4). These results could be
275 related to the greater initial vigour and the presence of more developed root systems in
276 larger seedlings compared to smaller seedlings. Consequently, these results suggest that
277 subcultivating *L. williamsii* seedlings of 4-5 mm are more appropriate for growing
278 purposes in order to boost their development in multiplication protocols.

279 Furthermore, no morphological variations, hyperhydration, or *calli* occurrence were
280 detected during the growth (neither in the germination assay nor in the growth assay). In

281 this way the plants obtained at the end of the assay displayed the usual appearance
282 observed in seedlings grown *ex vitro*. These results suggest that media used during
283 seedlings growth do not alter their genetic integrity, with seedlings having a normal
284 morphology. Therefore, plants from seeds harvested in nature and grown with our *in vitro*
285 protocols could be subsequently used to repopulate threatened populations without
286 evidence of modification of their genetic structure.

287

288 3.2.2 Number of areoles

289 Firstly, as can be observed on Table 4, the initial number of areoles was highly correlated
290 to the initial size of seedlings. Thus, 2-3 mm seedlings had 1-2 areoles, while seedlings
291 with 3-4 mm and 4-5 mm seedlings had 3-4 and 6-7 areoles, respectively. Nevertheless,
292 little differences were found among seedlings of the same size within 2-3 mm and 3-4
293 mm groups. This was probably due to: a) seedlings for this assay were merely selected
294 from germination assay based on size, without considering the number of areoles (random
295 effect), and/or b) plants of the same size came from different germination media (initial
296 media effect). Therefore, some differences in this trait could appeared randomly at the
297 beginning within the same size group.

298 Nevertheless, despite some initial little differences within each group, remarkable effects
299 of growth media were found for this trait. Thus, in the same way than for size, medium
300 GrM-1 enabled a greater increase in the number of areoles than any of the other two
301 media, regardless the initial size of the seedlings (Table 4). However, it was observed that
302 within the 4-5 mm group, there were no significant differences between the increase in
303 the number of areoles displayed by seedlings grown in GrM-1 and GrM-2, even
304 considering the five best seedlings of each group (data not shown).

305 In addition, seedlings from the smallest initial size, i.e. the 2-3 mm group displayed in
306 general a similar or higher increase in the number of areoles than those from higher initial
307 sizes. It was observed that the appearance of the areoles was especially intense after
308 germination (until seedlings reached 2-3 mm of diameter) and the distance between
309 areoles changed during growth (data not shown), being closer in the early stages of
310 seedling development (2-3 mm group) and separating as seedlings increased their volume
311 (3-4 mm and 4-5 mm groups). These observations may explain the lower increase in the
312 number of areoles observed in the 3-4 mm and 4-5 mm groups compared to the 2-3 mm
313 group, even more considering that the distance between areoles in adult plants can also
314 vary from 13 to 25 mm depending on the culture or environmental factors and it is closely
315 related with hydration of the plant (Calhoun, 2012; Lodé, 2015).

316 Probably, this greater increase in the number of areoles observed in the 2-3 group can be
317 expected in seedlings whose initial size and volume is smaller as an ancestral mechanism
318 of protection against adverse environmental conditions and/or presence of herbivores in
319 more susceptible seedlings (Ballester-Olmos, 1997; Hewitt, 2014). In fact, very often *L.*
320 *williamsii* genotypes develop a delicate transient spinescence in the first areoles, while
321 they lack pilosity (which appears later at more advanced stages) (Kunte and Subik, 2003;
322 Hunt, 2006). This is most clearly observed in *L. jourdaniana* (Haberm.) plantlets, on
323 which these small spines are kept even for years.

324

325 **4. Conclusions**

326 On the basis of our findings, we consider that the best protocol for *L. williamsii*
327 propagation would involve a seed germination and initial seedling growth in an *in vitro*
328 medium including half-strength MS, 8 g L⁻¹ agar and 20 g L⁻¹ sucrose (M5). Then, after

329 14-21 DAS, seedlings should be subcultivated in a full-strength MS, keeping initial agar
330 and sucrose concentrations (M1) to accelerate seedling growth. Once seedlings reach 4
331 mm in diameter, they should be subcultivated to a growth medium formulated with full-
332 strength MS, 15 g L⁻¹ of sucrose and 8 g L⁻¹ of agar and keep them there until reaching
333 the optimum size for rustication.

334 In the present work it has been possible to establish an *in vitro* germination and growth
335 protocol for *L. williamsii* seedlings that allows maximizing their initial development,
336 without the use of plant growth regulators. No morphological variations or *calli*
337 appearance were found in the seedlings during our culture assays. In this sense, apparently
338 these media without PGRs do not compromise the genetic or morphological integrity of
339 the cultivated seedlings. In consequence, these seedlings could be later reintroduced into
340 the habitat of threatened populations for repopulation purposes without altering the
341 genetic structure of the population. Furthermore, this *in vitro* germination and growth
342 protocol PGRs free is also the first step to establish massive propagation techniques
343 focused on maintaining certain clones with ornamental interest.

344

345 **References**

- 346 Anderson, E.F., Arias S., Taylor N.P., Cattabriga A. 1994. Threatened cactus of Mexico.
347 London: Royal Botanical Gardens, Kew.
- 348 Anderson, E.F. 1996. Peyote: The Divine Cactus, 2nd ed. University of Arizona Press,
349 Tucson.
- 350 Ballester-Olmos, J.F. 1997. Cactus y plantas suculentas. Ed.R.Guillén. Floraprint España.,
- 351 Bhojwani, S.S., Razdan, M.K. 1996. Plant tissue culture: Theory and practice. Elsevier,
352 Amsterdam.

353 Borchers, A.T., Keen, C.L., Stern, J.S., Gershwin, M.E. 2000. Inflammation and Native
354 American medicine: the role of botanicals. *Am J Clin Nutr.* 72: 339-347.

355 Calhoun, S. 2012. *The Gardener's Guide to Cactus: The 100 Best Paddles, Barrels,*
356 *Columns, and Globes.* Timber Press. Portland. USA.

357 Casado, R., Uriarte, I., Cavero, R.Y., Calvo, M.I. 2008. LC-PAD Determination of
358 Mescaline in Cactus "Peyote" (*Lophophora williamsii*). *Chromatographia.* 67: 665-
359 667.

360 Fischer, C., Neuhaus, G. 1995. *In vitro* development of globular zygotic wheat embryos.
361 *Plant Cell. Rep.* 15: 186-191.

362 Garcia-Osuna, H.T., Benavides-Mendoza, A., Escobedo-Bocardo, L., Villareal-
363 Quintanilla, J.A., Cornejo-Oviedo, E. 2011. Hyperhydricity control of *in vitro*
364 shoots of *Turbinicarpus valdezianus* (Moller) Gl. & F. *Phyton.* 80: 175-179.

365 Gland-Zwerger, A. 1995. Culture conditions affecting induction and regeneration in
366 isolated microscope cultures of different *Brassica* species. In: GCIRC Proceedings
367 of the ninth international Rapeseed Congress. GCIRC, Cambridge, UK. Pp 799-
368 801.

369 Halpern, J.H., Sherwood, A.R., Hudson, J.I., Yurgelun-Todd, D., Pope, H.G. 2005.
370 Psychological and cognitive effects of long-term peyote use among Native
371 Americans. *Biol Psychiatry.* 58: 624-631.

372 Hewitt, T. 2014. *Cacti.* Anness Publishing. London, U.K. ISBN – 9781780192840.

373 Hunt, D. 2006. *The new cactus lexicon.* Ed. DH Books. England. ISBN: 0-9538134-4-4.

374 Koné, M., Koné, T., Silué, N., Soumahoro, A.B., Kouakou, T.H. 2015. *In vitro* seeds
375 germination and seedling growth of Bambara groundnut (*Vigna subterranean* (L.)
376 Verdc. (Fabaceae)). *The Scientific World J.* 2015:595073.

377 Kunte, L., Subik, R. 2003. The complete Encyclopedia of Cacti. Rebo Productions. The
378 Netherlands.

379 Langer, D.F., Mergener, R.A. 2013. Cultivo *in vitro* de *Cereus Hildmannianus* K. Shum.
380 Unoesc & Ciência – ACBS 4: 7-14. (in Portuguese)

381 Lema-Ruminska, J., Kulus, D. 2012. Induction of somatic embryogenesis in *Astrophytum*
382 *asterias* (Zucc.) Lem. In the aspect of light conditions and auxin 2,4-D
383 concentrations. Acta Scientiarum Polonorum. Hortorum Cultus. 11: 77-87.

384 Lema-Ruminska, J., Kulus, D. 2014. Micropropagation of Cacti – A Review. Haseltonia.
385 19: 46-63.

386 Little, T.M., Hills, F.J. 1978. Agricultural experimentation: Design and analysis. Wiley
387 & Sons, New York.

388 Lodé, J. 2015. Taxonomy of the Cactaceae. Cactus Adventures Ed. Almeria. Spain.

389 Manzur, J.P., Penella, C., Rodríguez-Burruezo, A. 2013. Effect of the genotype,
390 developmental stage and médium composition on the *in vitro* culture efficiency of
391 immature zygotic embryos from genus *Capsicum*. Sci. Hort. 161:181-187.

392 McLaughlin, J.L., 1973. Peyote: an introduction. Lloydia. 36: 1-8.

393 Moebius-Goldammer, K.G., Mata-Rosas, M., Chávez-Ávila, V.M. 2003. Organogenesis
394 and somatic embryogenesis in *Ariocarpus kotschoubeyanus* (Lem.) K. Schum.
395 (Cactaceae), an endemic and endangered Mexican species. In Vitro Cell. Biol.-
396 Plant 39: 388-393.

397 Monnier, M., 1995. Culture of zygotic embryos. In: Thorpe, T.A. (Ed.), *In vitro*
398 embryogenesis in plants. Kluwer Academic Publishers, Dordrecht, pp. 117–153.

399 Paul, S., Kumaria, S., Tandon, P. 2012. An effective nutrient medium for asymbiotic seed
400 germination and large-scale *in vitro* regeneration of *Dendrobium hookerianum*, a
401 threatened orchid of northeast India. AoB Plants 2012: plr032.

402 Pérez-Molphe-Balch, E., Pérez-Reyes, M.E., Villalobos-Amador, E., Meza-Rangel, E.,
403 Del Rocío-Morones-Ruiz, L., Lizalde-Viramontes, H.J. 1998. Micropropagation of
404 21 species of Mexican cacti by axillary proliferation. *In vitro* celular and
405 developmental biology. *Plant* 34: 131-135.

406 Pérez-Molphe-Balch, E., Pérez-Reyes, M.E., Dávila-Figueroa, C.A., Villalobos-Amador,
407 E. 2002. *In vitro* propagation of three species of columnar cacti from the Sonoran
408 Desert. *Horticultural Sci.* 37: 693-696.

409 Quiala, E., Matos J., Montalvo, G., Feria, M., Chavez, M., Capote, A., Perez, N., Barbon,
410 R., Kowalski, B. 2009. *In vitro* propagation of *Pilosocereus robinii* (Lemaire) Byles
411 et Rowley, endemic and endangered cactus. *Journal of the Professional Association*
412 *for Cactus Development.* 11: 18-25.

413 Sajeva, M., Augugliaro, C., Smith, M., Oddo, E. 2013. Regulating internet trade in CITES
414 species. *Conserv Biol.* 27: 429-430.

415 Sanchez-Martinez, E., Hernandez-Martinez, M.M., Hernandez-Oria, J.G., Campos, G.R.,
416 Vara, P.M. Torres-Galeana, L.E. 2009. *Mammillaria herrerae* Werderm., a
417 Mexican species on the brink of extinction. *Mitteilungsblatt des Arbeitskreises für*
418 *Mammillarienfreunde.* 33: 26-37.

419 Stewart, S.L., Kane, M.E. 2006. Asymbiotic seed germination and *in vitro* seedling
420 development of *Habenaria macroceratitis* (Orchidaceae), a rare Florida terrestrial
421 orchid. *Plant Cell, Tissue and Organ Culture.* 86:147–158

422 Taylor, N. 1997. Cactaceae. In: Oldfield S. (ed.) *Cactus and succulent plants – status*
423 *survey and conservation action.* Plan IUC/SSCN Cactus and Succulent Specialist
424 Group, Gland, Switzerland and Cambridge, UK, pp. 1-2.

425 Xu, L., Najeeb, U., Tang, G.X., Gu, H.H., Zhang, G.Q., He, Y., Zhou, W.J. 2007. Haploid
426 and double haploid technology. In: Advances in botanical research: Rapeseed
427 Breeding (Ed: Gupta, S.K.).

428

429

430

431

432

433

434

435

436

437

438 Table 1. Composition of *in vitro* media studied in both the germination and plant growth
439 assays of *L. williamsii*.

Media	Murashige & Skoog	Sucrose (g L⁻¹)	Agar (g L⁻¹)
<i>Germination Assay</i>			
M1	full-strength	20	8
M2	full-strength	20	10
M3	full-strength	30	8

M4	full-strength	30	10
M5	half-strength	20	8
M6	half-strength	20	10
M7	half-strength	30	8
M8	half-strength	30	10
<i>Growth Assay</i>			
GrM-1	full-strength	15	8
GrM-2	full-strength	30	8
GrM-3	full-strength	45	8

440

441

442

443

444

445

446

447 Table 2. Germination assay: germination efficiency and percentage of *L. williamsii* seedlings
 448 with areoles for the eight *in vitro* media evaluated along 49 days after sowing (DAS).

449

Media	7 DAS	14 DAS	21 DAS	28 DAS	35 DAS	42 DAS	49 DAS
-------	-------	--------	--------	--------	--------	--------	--------

Germination efficiency (%)

M1	0.0 a ¹	42.5 a	60.0 a	60.0 a	62.5 a	67.5 a	67.5 a
M2	7.5 abc	55.0 ab	67.5 a	70.0 a	70.0 a	70.0 a	70.0 a
M3	5.0 ab	60.0 b	65.0 a	70.0 a	70.0 a	70.0 a	75.0 a
M4	12.5 bc	57.5 ab	70.0 a	70.0 a	70.0 a	72.5 a	75.0 a
M5	12.5 bc	55.0 ab	62.5 a	67.5 a	70.0 a	72.5 a	72.5 a
M6	17.5 c	67.5 b	67.5 a	67.5 a	67.5 a	70.0 a	70.0 a
M7	12.5 bc	52.5 ab	55.0 a	65.0 a	65.0 a	67.5 a	67.5 a
M8	0.0 a	62.5 b	67.5 a	67.5 a	70.0 a	72.5 a	77.5 a
1/2MS	10.6 A	59.4 A ²	63.1 A	66.9 A	68.1 A	70.6 A	71.9 A
MS	6.3 A ²	53.8 A	65.6 A	67.5 A	68.1 A	70.0 A	71.9 A
Sucrose 20 g L ⁻¹	9.4 A	55.0 A	64.4 A	66.3 A	67.5 A	70.0 A	70.0 A
Sucrose 30 g L ⁻¹	7.5 A	58.1 A	64.4 A	68.1 A	68.8 A	70.6 A	73.8 A
Agar 8 g L ⁻¹	7.5 A	52.5 A	60.6 A	65.6 A	66.9 A	69.4 A	70.6 A
Agar 10 g L ⁻¹	9.4 A	60.6 A	68.1 A	68.8 A	69.4 A	71.3 A	73.1 A

Seedlings with areoles (%)

M1	0.0	0.0 a	32.5 ab	42.5 ab	42.5 ab	42.5 ab	42.5 a
M2	0.0	0.0 a	35.0 ab	50.0 b	52.5 ab	52.5 ab	52.5 a
M3	0.0	5.0 ab	42.5 ab	52.5 b	55.0 ab	60.0 b	62.5 a
M4	0.0	7.5 ab	22.5 a	25.0 a	35.0 a	37.5 a	45.0 a
M5	0.0	10.0 b	45.0 b	52.5 b	62.5 b	62.5 b	65.0 a
M6	0.0	12.5 b	40.0 ab	45.0 ab	52.5 ab	52.5 ab	55.0 a
M7	0.0	10.0 b	30.0 ab	40.0 ab	45.0 ab	47.5 ab	50.0 a
M8	0.0	0.0 a	22.5 a	32.5 ab	42.5 ab	47.5 ab	50.0 a
1/2MS	0.0	8.1 A ²	34.4 A	42.5 A	50.6 A	52.5 A	55.0 A

MS	0.0	3.1 A	33.1 A	42.5 A	46.3 A	48.1 A	50.6 A
Sucrose 20 g L ⁻¹	0.0	5.6 A	38.1 A	47.5 A	52.5 A	52.5 A	53.8 A
Sucrose 30 g L ⁻¹	0.0	5.6 A	29.4 A	37.5 A	44.4 A	48.1 A	51.9 A
Agar 8 g L ⁻¹	0.0	6.3 A	37.5 A	46.9 A	51.3 A	53.1 A	55.0 A
Agar 10 g L ⁻¹	0.0	5.0 A	30.0 A	38.1 A	45.6 A	47.5 A	50.6 A

450 ¹ Values with different letters among *in vitro* media indicate significant differences according to
451 the least significant difference test (LSD) at $P < 0.05$.

452 ² Values with different capital letters indicate significant differences between levels of each
453 component of *in vitro* media according to the least significant difference test (LSD) at $P < 0.05$.

454

455

456

457

458

459

460 Table 3. Germination assay: seedling size and number of areoles per *L. williamsii* seedling for
461 the eight *in vitro* media evaluated along 49 days after sowing (DAS).

Media	7 DAS	14 DAS	21 DAS	28 DAS	35 DAS	42 DAS	49 DAS
<i>Seedling size (mm)</i>							
M1	0.0	2.7 ab ¹	3.4 c	3.6 e	3.9 e	4.0 e	4.1 e

M2	0.0	2.6 ab	2.5 a	2.5 ab	2.8 ab	2.8 ab	2.9 ab
M3	0.0	2.6 ab	2.6 a	2.9 bcd	3.1 bc	3.2 bc	3.3 bc
M4	0.0	2.4 a	2.4 a	2.4 a	2.5 a	2.6 a	2.6 a
M5	0.0	3.4 c	3.4 c	3.3 de	3.7 de	3.8 de	3.9 de
M6	0.0	2.6 ab	2.7 ab	2.8 abc	2.9 ab	2.9 ab	2.9 abc
M7	0.0	3.0 b	3.3 bc	3.3 cde	3.4 cd	3.5 cd	3.5 cd
M8	0.0	2.4 a	2.4 a	2.5 ab	2.7 ab	2.7 ab	2.8 ab
1/2MS	0.0	2.9 B ²	3.0 B	3.0 A	3.2 A	3.2 A	3.3 A
MS	0.0	2.6 A	2.7 A	2.8 A	3.1 A	3.1 A	3.2 A
Sucrose 20 g L ⁻¹	0.0	2.8 A	3.0 A	3.1 A	3.3 B	3.4 B	3.4 B
Sucrose 30 g L ⁻¹	0.0	2.6 A	2.7 A	2.8 A	2.9 A	3.0 A	3.0 A
Agar 8 g L ⁻¹	0.0	2.9 B	3.2 B	3.3 B	3.5 B	3.6 B	3.7 B
Agar 10 g L ⁻¹	0.0	2.5 A	2.5 A	2.6 A	2.7 A	2.8 A	2.8 A

Areoles per seedling

M1	0.0	0.00 a	1.13 a	2.81 d	3.91 d	4.53 e	5.19 e
M2	0.0	0.00 a	1.23 ab	1.49 a	1.85 ab	2.26 ab	2.80 ab
M3	0.0	0.50 ab	1.04 a	1.91 abc	2.39 b	3.33 cd	3.83 cd
M4	0.0	0.50 ab	1.08 a	1.56 ab	1.46 a	1.88 a	1.94 a
M5	0.0	0.75 b	1.49 b	2.67 d	3.08 c	4.09 de	4.64 de
M6	0.0	0.75 b	1.25 ab	2.09 bc	2.29 b	2.88 bc	3.34 bc
M7	0.0	1.00 b	1.23 ab	2.33 cd	3.06 c	3.56 cd	3.77 cd
M8	0.0	0.00 a	1.13 a	1.81 abc	1.80 ab	2.62 abc	2.84 b
1/2MS	0.0	0.63 B	1.27 B	2.23 A	2.56 A	3.29 A	3.65 A
MS	0.0	0.25 A	1.05 A	1.94 A	2.40 A	3.00 A	3.44 A

Sucrose 20 g L ⁻¹	0.0	0.38	A	1.27	A	2.27	A	2.78	B	3.44	B	3.99	B
Sucrose 30 g L ⁻¹	0.0	0.50	A	1.12	A	1.90	A	2.18	A	2.85	A	3.10	A
Agar 8 g L ⁻¹	0.0	0.56	A	1.22	A	2.43	B	3.11	B	3.88	B	4.36	B
Agar 10 g L ⁻¹	0.0	0.31	A	1.10	A	1.74	A	1.85	A	2.41	A	2.73	A

462 ¹ Values with different letters among *in vitro* media indicate significant differences according to
463 the least significant difference test (LSD) at $P < 0.05$.

464 ² Values with different capital letters indicate significant differences between levels of each
465 component of *in vitro* media according to the least significant difference test (LSD) at $P < 0.05$.

466

467

468

469

470

471

472

473 Table 4. Growth assay: evolution of seedling size and mean number of areoles per *L. williamsii*
474 seedling in germinated seedlings, grouped by initial sizes, after 105 days of *in vitro* growth
475 culture in three different growth media (GrM at 15, 30 and 45 g sucrose·L⁻¹).

	Size (mm)			Areoles per seedling		
	Initial	Final	Increase	Initial	Final	Increase
2-3 mm seedlings						
GrM-1	2,5 a	4,4 b	1,9 b	1,32 a	9,11 b	7,79 b

GrM-2	2,5	a	3,6	a	1,1	a	1,42	a	6,84	a	5,42	a
GrM-3	2,5	a	3,5	a	1,0	a	1,92	b	6,62	a	4,69	a
3-4 mm seedlings												
GrM-1	3,1	a	4,4	b	1,3	b	3,09	a	9,03	a	5,94	b
GrM-2	3,1	a	3,9	a	0,8	a	3,68	b	8,26	a	4,58	a
GrM-3	3,2	a	4,0	a	0,8	a	3,90	b	8,30	a	4,40	a
4-5 mm seedlings												
GrM-1	4,3	a	6,5	b	2,3	b	6,33	a	12,83	a	6,50	b
GrM-2	4,3	a	5,8	a	1,5	ab	6,80	a	12,80	a	6,00	ab
GrM-3	4,4	a	5,6	a	1,1	a	6,73	a	11,36	a	4,64	a

476 ¹ Values with different letters among *in vitro* media within each size group indicate significant
477 differences according to the least significant difference test (LSD) at $P < 0.05$.

478

479

480

481

482

483

484

485 FIGURES

486

487 Figure 1. *Lophophora williamsii* morphology: a) adult plant, b) napiform root, c)
488 flowering and d) immature fruit and mature dry fruit containing viable seeds. Black bar
489 indicates 10 mm.

490 Figure 2. Comparative radicle development at 21 days after sowing in media with
491 different agar concentration: M1 (8 g L⁻¹) (left) vs. M8 (10 g L⁻¹) (right). Each small
492 square in the grid indicates a size of 1x1 mm.

493 Figure 3. Comparative initial seedling development at 49 days after sowing: medium 1
494 (left) vs. medium 4 (right). Each small square in the grid indicates a size of 1x1 mm.

495 Figure 4. Comparative seedlings sizes per week in each media. Each color represents the
496 increase of size (in mm) during each time interval (days after sowing, DAS).

497 Figure 5. Comparison of final size at the end of the growth assay (105 days after
498 subcultivation) for each combination of initial size x growing media. Each small square
499 in the grid indicates a size of 1x1 mm.

500

501

502

503