

Metal-Free Diastereo- and Enantioselective Dearomative Formal [3 + 2] Cycloaddition of 2-Nitrobenzofurans and Isocyanoacetate Esters

Adrian Laviós, Amparo Sanz-Marco, Carlos Vila, M. Carmen Muñoz, José R. Pedro,* and Gonzalo Blay*



Cite This: *Org. Lett.* 2022, 24, 2149–2154




Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: The diastereo- and enantioselective dearomative formal $[3 + 2]$ cycloaddition of 2-nitrobenzofurans and α -aryl- α -isocyanoacetate esters provides tricyclic compounds bearing the $3a,8b$ -dihydro- $1H$ -benzofuro[2,3-*c*]pyrrole framework with three consecutive stereogenic centers. The reaction was enabled by a cuprene-ether organocatalyst. The reaction products were obtained with almost full diastereoselectivity and with excellent enantiomeric excesses for a number of substituted 2-nitrobenzofurans and isocyanoacetates.

The asymmetric dearomatization of arenes and heteroarenes has become a powerful tool for accessing chiral structures from simple, easily available planar scaffolds.¹ Many of these reactions involve electron-poor aromatic *aza*-heterocycles such as pyridines, pyrimidines, and their benzo-fused analogues as an approach toward the synthesis of chiral six-membered nonaromatic *aza*-heterocycles.² Dearomatization of these compounds usually involves the participation of species with a positive nitrogen that are obtained through either protonation or *N*-alkylation. In contrast, the asymmetric dearomatization of five-membered heterocycles normally has exploited the nucleophilic character of these electron-rich heterocycles. Nevertheless, installing proper electron-withdrawing substituents on the heteroarenes can reverse them into electron-deficient compounds, thus serving as electrophiles for Umpolung-like reactions.³

Following this strategy, electron-deficient nitroheteroarenes have been successfully used as dienophiles and dipolarophiles in cycloaddition reactions, which are normally initiated by Michael addition followed by intramolecular trapping of the anion by an electrophilic group. Pioneering work by Arai⁴ and Trost⁵ showed the potential of 3-nitroindoles as dipolarophiles in catalytic asymmetric dearomative formal $[3 + 2]$ cycloaddition reactions. Since then, 2- and 3-nitroindoles⁶ and related nitroarenes⁷ have been disclosed as dipolarophiles in cycloaddition reactions with a variety of formal dipoles to provide a great variety of polycyclic structures containing multiple stereogenic centers. Among these, dearomatization of 2-nitrobenzofurans lead to compounds containing a chiral 2,3-dihydrobenzofuran scaffold, which is an important pharmacophore present in biologically active natural products and pharmaceuticals.⁸ Despite these precedents, dearomative cycloaddition reactions with 2-nitrobenzofurans remain rare, and only a reduced number of them have been developed.^{6b,e,9}

On the other hand, α -isocyano esters stand as useful and versatile formal 1,3-dipoles. The high α - acidity together with the electrophilic ability of the isocyano group allows isocyano esters to undergo tandem/cascade reactions with electrophilic unsaturated systems, leading to a variety of five-membered *aza*-heterocycles.¹⁰ These formal $[3 + 2]$ cycloaddition reactions normally involve nucleophilic addition of the α -enolate followed by intramolecular attack of the resulting anion to the empty orbital of the isocyano group. Despite some success in the application of this strategy, there is still a considerable interest in developing new cycloaddition reactions involving isocyanoacetate esters as formal dipoles to access a diversity of novel structures.

Gribble and coworkers studied the reaction of N-protected 3-nitroindoles with ethyl isocyanoacetate, which provided nonchiral pyrrolo[2,3-*b*]indoles or pyrrolo[3,4-*b*]indoles depending on the N1 protecting group via a Barton-Zard reaction involving the elimination of HNO_2 (Scheme 1).¹¹ However, the Barton-Zard reaction can be interrupted by

Received: February 7, 2022

Published: March 16, 2022




Scheme 1. (Interrupted) Barton–Zard Reactions with Nitroarenes


Gribble et al.¹¹




Yuan and You¹³



Yuan and Zhao¹⁴



This work




using α -substituted α -isocyano esters.¹² Using this modification, Yuan and You developed a catalytic asymmetric dearomatization reaction of 3-nitroindoles with α -substituted α -isocyano esters through an interrupted Barton–Zard reaction enabled by a silver/cinchona-derived aminophosphine catalyst (Scheme 1).¹³ The reaction provided the expected products with fair to good diastereoselectivity and excellent enantioselectivity for α -alkyl isocyanoacetates. A similar catalyst has been recently reported by Yuan and Zhao to catalyze the reaction of 2-nitroindoles with α -aryl α -isocyano esters. The authors also reported the reaction with 3-

nitroindoles by using an Ag/BINAP complex (Scheme 1).¹⁴ To the best of our knowledge, a similar reaction has not been reported under metal-free conditions or with other nitroarenes. Herein we report our results on the organocatalytic enantioselective dearomative [3 + 2]-cycloaddition of 2-nitrobenzofurans and isocyanoacetate esters (Scheme 1).


To carry out the optimization process, we used the reaction between 2-nitrobenzofuran (**1a**) and methyl 2-phenyl-2-isocyanoacetate (**2a**). Following our previous research on isocyanoacetate cycloadditions,¹⁵ we first tested the reaction in the presence of bifunctional squaramide **I** and Ag₂O in dichloromethane. Under these conditions, compound **3a** was obtained with fair diastereoselectivity, although with low enantiomeric excess for the major diastereomer. We also observed that silver oxide alone was able to catalyze the reaction. Accordingly, further optimization was pursued in the absence of silver salts (Table 1). In this way, squaramide **I** provided compound **3a** with null diastereoselectivity, although the ee increased for both diastereomers. A number of bifunctional squaramides were tested, but none of them allowed us to obtain a good enantiomeric excess for the major diastereomer. (See the SI.) In a similar way, thiourea **II** provided better diastereoselectivity but again with low ee for the major diastereomer. Fortunately, cupreine derivative **III** allowed us to obtain compound **3a** as just one diastereomer with a promising 73% ee (Table 1, entry 4). Increasing the catalyst load to 15 mol % had little effect on the stereoselectivity, whereas lowering the reaction temperature to 0 °C improved both the diastereo- and enantioselectivity, although at the expense of reducing the yield (Table 1, entry 6). Increasing the concentration permitted us to improve the yield while keeping the high stereoselectivity. Finally, several solvents were tested, with the best result being obtained in

Table 1. Optimization of the Reaction Conditions^a



entry	OC (mol %)	solvent	[1a] (M)	T (°C)	yield (%) ^b	dr ^c	ee (%) ^d
1 ^e	I (10)	CH ₂ Cl ₂	0.1	rt	95	61:39	10/30
2	I (10)	CH ₂ Cl ₂	0.1	rt	95	50:50	35/37
3	II (10)	CH ₂ Cl ₂	0.1	rt	62	17:83	80/6
4	III (10)	CH ₂ Cl ₂	0.1	rt	73	3:97	9/73
5	III (15)	CH ₂ Cl ₂	0.1	rt	65	1:99	nd/76
6	III (15)	CH ₂ Cl ₂	0.1	0	48	1:99	nd/93
7	III (15)	CH ₂ Cl ₂	0.2	0	75	1:99	nd/90
8	III (10)	CH ₂ Cl ₂	0.2	0	74	1:99	nd/90
9	III (10)	MTBE	0.2	0	46	1:99	nd/74
10	III (10)	toluene	0.2	0	40	1:99	nd/93
11	III (10)	CHCl ₃	0.2	0	79	1:99	nd/93
12 ^f	III' (10)	CHCl ₃	0.2	0	19	1:99	nd/10

^aConditions: **1a** (0.1 mmol), **2a** (0.13 mmol), OC, solvent, 48 h. ^bIsolated yield after column chromatography. ^cDetermined by ¹H NMR. ^dDetermined by HPLC over chiral stationary phases. ^eAg₂O (5 mol %) was used. ^fControl experiment; see the mechanistic discussion.

Scheme 2. Scope of the Reaction of 2-Nitrobenzofurans 1 and Isocyanoacetates 2^a

^a Conditions: 1a (0.15 mmol), 2a (0.19 mmol), III (0.015 mmol), CHCl₃ (0.75 mL), 0 °C. Yields after column chromatography, *dr* determined by ¹H NMR, ee determined by HPLC. ^b Reaction carried out on a 1 mmol scale.

chloroform, which furnished compound 3a in 79% yield as only one diastereomer with 93% ee (Table 1, entry 11).


Under the optimized conditions, the scope of the reaction of methyl 2-phenylisocyanoacetate (2a) and several substituted 2-nitrobenzofurans 1 was studied (Scheme 2, 1a–1o). The reaction could be successfully achieved with 2-nitrobenzofurans bearing substituents of varied electronic natures at different positions of the homoaromatic ring. Good yields, full diastereoselectivities, and high enantioselectivities were obtained with nitrobenzofurans substituted at the five (3ba–3ga), six (3ha, 3ia), or seven (3ja–3la) position with either electron-donating or electron-withdrawing groups. In general, somehow better yields were obtained when this substituent was a halogen or an electron-withdrawing group. Unfortunately, substitution at position 4 of the benzofuran ring brought about a serious decrease in yield and enantioselectivity (3ma). We also tested some disubstituted nitrobenzofurans. 5,7-Dichloro-2-nitrobenzofuran (1o) reacted with 2a to give the expected cycloaddition product 3oa in excellent yield with excellent enantioselectivity. On the contrary, compound 1n bearing two bulky *tert*-butyl groups reacted to give 3na with good enantioselectivity, although in lower yield.

Next, we studied the reaction with different isocyanoacetates 2. Methyl (4-methoxyphenyl)isocyanoacetate (2b, R² = 4-

MeOC₆H₄) reacted with several 2-nitrobenzofurans to give the corresponding products 3ab, 3eb, and 3ib in good yields with excellent enantioselectivities, higher than those obtained with isocyanoacetate 2a. However, when an electron-withdrawing group was introduced on the aryl ring of the isocyanoacetate (2c, R² = 4-ClC₆H₄; 2d, R² = 4-NO₂C₆H₄) the reaction products 3ac and 3ad were obtained with lower enantioselectivities, although still in good yields with full diastereoselectivity. On the contrary, methyl 2-isocyanopropanoate (2e, R² = Me) reacted with 1a and provided a racemic product 3ae in low yield. We also tested the reaction of 2a with 2-nitrobenzotriphene, which was sluggish and gave compound 4 in low yield with fair enantioselectivity (72% ee). Finally, the reaction with the less synthetically accessible 3-nitrobenzofurans was attempted. The reaction of 2a with commercially available 5-hydroxy-3-nitrobenzofuran did not proceed under our reaction conditions, probably due to the presence of the free OH group in the substrate (Scheme 2, 5a). In fact, 5-acetoxy-3-nitrobenzofuran reacted with 2a to give compound 5b in ca. 30% yield, although unfortunately, the enantiomeric excess could not be determined.¹⁶

To highlight the robustness of the method, we carried out the reaction of 1a and 2a on a 1 mmol scale to obtain compound 3aa in 75% yield with 91% ee, which was

comparable to that obtained in the model reaction. Furthermore, some synthetic transformations were carried out with compound **3aa** (**Scheme 3**). Elimination of the nitro group upon treatment with DBU in dichloromethane gave compound **6** in 53% yield with 90% ee. Reduction of the C–N double bond was achieved with Et₃SiH in the presence of BF₃·Et₂O to give compound **7** in quantitative yield with a slight erosion of the ee. Finally, aminoacetal **8** could be obtained in 99% yield with full diastereoselectivity and 98% ee after refluxing in MeOH. In a similar way, starting from **3da**, we could obtain compound **9**, which could be crystallized and subjected to single-crystal X-ray diffraction analysis (**Figure 1**).



group upon treatment with DBU in dichloromethane gave compound **6** in 53% yield with 90% ee. Reduction of the C–N double bond was achieved with Et₃SiH in the presence of BF₃·Et₂O to give compound **7** in quantitative yield with a slight erosion of the ee. Finally, aminoacetal **8** could be obtained in 99% yield with full diastereoselectivity and 98% ee after refluxing in MeOH. In a similar way, starting from **3da**, we could obtain compound **9**, which could be crystallized and subjected to single-crystal X-ray diffraction analysis (**Figure 1**).




Figure 1. Proposed stereochemical model and ORTEP plot for the X-ray structure of compound **9** with thermal ellipsoids drawn at the 50% probability level. Flack parameter 0.005(15).

In this way, we could establish its absolute stereochemistry and hence that of compound **3da**. The stereochemistry (1*R*,3*aR*,8*bR*) of all compounds **3** was assigned by analogy, assuming a common stereochemical pathway.

To get some insight into the reaction mechanism, we performed a control experiment running the reaction of **1a** and **2a** in the presence of compound **III'**, the methyl ether of catalyst **III** lacking a free OH phenol group. Under these conditions, the reaction took place sluggishly to give compound **3aa** in 19% yield with 10% enantiomeric excess (**Table 1**, entry 12). These results indicate that compound **III** most probably works as a bifunctional catalyst with the free OH phenol group and the basic amine acting synergistically. According to literature precedents and the observed stereochemistry,¹⁷ we propose the stereochemical model outlined in **Figure 1**. Catalyst **III** activates the electrophile through H bonding of the phenol and the nitro group, whereas the tertiary base deprotonates the isocyano ester. The approach of the

nucleophile is guided by ion pairing with the ammonium salt facing the *Si* face of the ester enolate and the *Si* face of the furan double bond to account for the observed stereochemistry.

In summary, we have established a highly diastereo- and enantioselective organocatalytic procedure for the dearomatic formal [3 + 2] cycloaddition of 2-nitrobenzofurans and 2-aryl-2-isocyanoacetate esters. Although related reactions with 2- and 3-nitroindoles have been reported, this is the first example using 2-nitrobenzofurans. The reaction is catalyzed by a cupreine derivative in the absence of metals and provides chiral tricyclic compounds in good yields with full diastereoselectivity and excellent enantioselectivity for a number of 2-nitrobenzofuran derivatives substituted at the five, six, or seven position. The arylisocyanoacetate ester allows substitution at the aryl ring, with the best results being obtained with phenyl or 4-methoxyphenyl derivatives. The potential synthetic applicability of the reaction has been demonstrated with a 1 mmol scale reaction and versatile modifications of compound **3aa**.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.2c00427>.

Experimental procedures, characterization data, NMR spectra, and HPLC traces (**PDF**)

FAIR data, including the primary NMR FID files, for compounds **3–8** (**ZIP**)

Accession Codes

CCDC 2149627 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

Gonzalo Blay – Departament de Química Orgànica, Facultat de Química, Universitat de València, 46100 València, Spain;
✉ orcid.org/0000-0002-7379-6789; Email: gonzalo.blay@uv.es

José R. Pedro – Departament de Química Orgànica, Facultat de Química, Universitat de València, 46100 València, Spain;
✉ orcid.org/0000-0002-6137-866X; Email: jose.r.pedro@uv.es

Authors

Adrian Laviós – Departament de Química Orgànica, Facultat de Química, Universitat de València, 46100 València, Spain;
✉ orcid.org/0000-0002-0259-7336

Amparo Sanz-Marco – Departament de Química Orgànica, Facultat de Química, Universitat de València, 46100 València, Spain;
✉ orcid.org/0000-0002-1729-598X

Carlos Vila – Departament de Química Orgànica, Facultat de Química, Universitat de València, 46100 València, Spain;
✉ orcid.org/0000-0001-9306-1109

M. Carmen Muñoz – Departament de Física Aplicada, Universitat Politècnica de València, 46022 València, Spain;
✉ orcid.org/0000-0003-2630-3897

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.orglett.2c00427>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We acknowledge grant PID2020-116944GB-100 funded by MCIN/AEI/10.13039/501100011033 and by the “European Union Next Generation EU/PRTR”, grant RyC-2016-20187 funded by MCIN/AEI/10.13039/501100011033 and by “ESF Investing in your future” to C.V., and grant FPU18/03038 funded by MEC to A.L. We acknowledge access to the NMR, MS, and X-ray facilities from the Servei Central de Suport a la Investigació Experimental (SCSIE)-UV. Dedicated to Professor Joan Bosch, Universitat de Barcelona, on the occasion of his 75th birthday.

REFERENCES

- (1) (a) *Asymmetric Dearomatization Reactions*; Yhou, S. L., Ed.; Wiley-VCH: Weinheim, Germany, 2016. (b) Sheng, F.-T.; Wang, J.-Y.; Tan, W.; Zhang, Y.-C.; Shi, F. *Org. Chem. Front.* **2020**, *7*, 3967–3998. (c) Zheng, C.; You, S.-L. Advances in Catalytic Asymmetric Dearomatization. *ACS Central Sci.* **2021**, *7*, 432–444.
- (2) (a) Matador, E.; Iglesias-Sigüenza, J.; Monge, D.; Merino, P.; Fernandez, R.; Lassaletta, J. M. Enantio- and Diastereoselective Nucleophilic Addition of *N*-*tert*-butylhydrazones to Isoquinolinium Ions Through Anion-Binding Catalysis. *Angew. Chem., Int. Ed.* **2021**, *60*, 5096–5101. (b) Segovia, C.; Nocquet, P. A.; Levacher, V.; Briere, J. F.; Oudeyer, S. Organocatalysis: A Tool of Choice for the Enantioselective Nucleophilic Dearomatization of Electron-Deficient Six-Membered Ring Azaarenum Salts. *Catalysts* **2021**, *11*, 1249. (c) Gomez-Martinez, M.; del Carmen Perez-Aguilar, M.; Piekarski, D. G.; Daniliuc, C. G.; Garcia-Mancheño, O. *N,N*-Dialkylhydrazones as Versatile Umpolung Reagents in Enantioselective Anion-Binding Catalysis. *Angew. Chem., Int. Ed.* **2021**, *60*, 5102–5107.
- (3) Chataigner, I.; Piettre, S. R. Multicomponent Domino [4 + 2]/[3 + 2] Cycloadditions of Nitroheteroaromatics: An Efficient Synthesis of Fused Nitrogenated Polycycles. *Org. Lett.* **2007**, *9*, 4159–4162.
- (4) Awata, A.; Arai, T. PyBidine/Copper Catalyst: Asymmetric exo-Selective [3 + 2] Cycloaddition using Imino Ester and Electrophilic Indole. *Angew. Chem., Int. Ed.* **2014**, *53*, 10462–10465.
- (5) Trost, B. M.; Ehmke, V.; O’Keefe, B. M.; Bringley, D. A. Palladium-Catalyzed Dearomative Trimethylenemethane Cycloaddition Reactions. *J. Am. Chem. Soc.* **2014**, *136*, 8213–8216.
- (6) For a recent review, see: (a) Sheng, F.-T.; Wang, J.-Y.; Tan, W.; Zhang, Y.-C.; Shi, F. Progresses in organocatalytic asymmetric dearomatization reactions of indole derivatives. *Org. Chem. Front.* **2020**, *7*, 3967–3998. Selected examples: (b) Jiang, Z.; Liu, X.; Zhang, H.; Tan, J.-P.; Ren, X.; Gao, G.; Wang, T. Bifunctional Phosphonium Salt-Catalyzed Asymmetric Cyclodearomatization of 2-Nitroindoles and 2-Nitrobenzofurans for Constructing CF₃-Containing Spiro-Polycycles. *Adv. Synth. Catal.* **2021**, *363*, 3115–3120. (c) Ling, J.; Mara, D.; Roure, B.; Laugeois, M.; Vitale, M. R. Copper(I)-Catalyzed Dearomative (3 + 2) Cycloaddition of 3-Nitroindoles with Propargylic Nucleophiles: An Access to Cyclopenta[b]indolines. *J. Org. Chem.* **2020**, *85*, 3838–3848. (d) Mei, M.-S.; Wang, Y.-H.; Hu, Q.; Li, Q.-H.; Shi, D.-Y.; Gao, D.; Ge, G.; Lin, G.-Q.; Tian, P. Enantioselective [3 + 2] annulation of isatin-derived MBH-carbonates and 3-nitroindoles enabled by a bifunctional DMAP-thiourea. *Chem. Commun.* **2020**, *56*, 10718–10721. (e) Wang, H.; Zhang, J.; Tu, Y.; Zhang, J. Phosphine-Catalyzed Enantioselective Dearomative [3 + 2]-Cycloaddition of 3-Nitroindoles and 2-Nitrobenzofurans. *Angew. Chem., Int. Ed.* **2019**, *58*, 5422–5426. (f) Chen, X.-M.; Lei, C.-W.; Yue, D.-F.; Zhao, J.-Q.; Wang, Z.-H.; Zhang, X.-M.; Xu, X.-Y.; Yuan, W.-C. Organocatalytic Asymmetric Dearomatization of 3-Nitroindoles and 3-Nitrobenzothiophenes via Thiol-Triggered Diastereo- and Enantioselective Double Michael Addition Reaction. *Org. Lett.* **2019**, *21*, 5452–5456. (g) Rivinoja, D. J.; Gee, Y. S.; Gardiner, M.-G.; Ryan, J. H.; Hyland, C. J. T. The Diastereoselective Synthesis of Pyrroloindolines by Pd-Catalyzed Dearomative Cycloaddition of 1-Tosyl-2-vinyl-aziridine to 3-Nitroindoles. *ACS Catal.* **2017**, *7*, 1053–1056. (h) Cheng, Q.; Zhang, F.; Cai, Y.; Guo, Y.-L.; You, S.-L. Stereodivergent Synthesis of Tetrahydrofuroindoles through Pd-Catalyzed Asymmetric Dearomative Formal [3 + 2] Cycloaddition. *Angew. Chem., Int. Ed.* **2018**, *57*, 2134–2138. (i) Zhao, J.-Q.; Zhou, X.-J.; Chen, Y.-Z.; Xu, X.-Y.; Zhang, X.-M.; Yuan, W.-C. Zn-Catalyzed Diastereo- and Enantioselective Dearomative [3 + 2] Cycloaddition Reaction of 2-Nitroindoles and 2-Nitrobenzothiophenes. *Adv. Synth. Catal.* **2018**, *360*, 2482–2487. (j) Suo, J.-J.; Liu, W.; Du, J.; Ding, C.-H.; Hou, X.-L. Diastereo- and Enantioselective Palladium-Catalyzed Dearomative [3 + 2] Cycloaddition of 3-Nitroindoles. *Chem.—Asian J.* **2018**, *13*, 959–963. (k) Sun, M.; Zhu, Z.-Q.; Gu, L.; Wan, X.; Mei, G.-J.; Shi, F. Catalytic Asymmetric Dearomative [3 + 2] Cycloaddition of Electron-Deficient Indoles with All-Carbon 1,3-Dipoles. *J. Org. Chem.* **2018**, *83*, 2341–2348. (l) Zhang, J.-Q.; Tong, F.; Sun, B.-B.; Fan, W.-T.; Chen, J.; Hu, D.; Wang, X.-W. Pd-Catalyzed Asymmetric Dearomative Cycloaddition for Construction of Optically Active Pyrroloindoline and Cyclopentaindoline Derivatives: Access to 3a-Aminopyrroloindolines. *J. Org. Chem.* **2018**, *83*, 2882–2891. (m) Gerten, A. L.; Stanley, L. M. Enantioselective dearomative [3 + 2] cycloadditions of indoles with azomethine ylides derived from alanine imino esters. *Org. Chem. Front.* **2016**, *3*, 339–343. (n) Zhao, J.-Q.; Zhou, M.-Q.; Wu, Z.-J.; Wang, Z.-H.; Yue, D.-F.; Xu, X.-Y.; Zhang, X.-M.; Yuan, W.-C. Asymmetric Michael/Cyclization Cascade Reaction of 3-Isothiocyanato Oxindoles and 3-Nitroindoles with Amino-Thiocarbamate Catalysts: Enantioselective Synthesis of Polycyclic Spirooxindoles. *Org. Lett.* **2015**, *17*, 2238–2241. (o) Zhao, J.-Q.; Wu, Z.-J.; Zhou, M.-Q.; Xu, X.-Y.; Zhang, X.-M.; Yuan, W.-C. Zn-Catalyzed Diastereo- and Enantioselective Cascade Reaction of 3-Isothiocyanato Oxindoles and 3-Nitroindoles: Stereocontrolled Syntheses of Polycyclic Spirooxindoles. *Org. Lett.* **2015**, *17*, 5020–5023. (p) Yue, D.-F.; Zhao, J.-Q.; Chen, Y.-Z.; Zhang, X.-M.; Xu, X.-Y.; Yuan, W.-C. Zinc-Catalyzed Enantioselective Dearomative [3 + 2] Cycloaddition Reaction of 3-Nitrobenzothiophenes and 3-Nitrothieno[2,3-b]pyridine with 3-Isothiocyanato Oxindoles. *Adv. Synth. Catal.* **2018**, *360*, 1420–1425. (q) (a) Wang, T.; Tan, J.; Chen, Y. Chiral Polycyclic Tropane Compound and Its Preparing Method and Application. China Patent CN113292574 A, 2021. (B) Tan, J.-P.; Li, X.; Chen, Y.; Rong, X.; Zhu, L.; Jiang, C.; Xiao, K.; Wang, T. Highly Stereoselective Construction Of Polycyclic Benzofused Tropane Scaffolds and their Latent Bioactivities: Bifunctional Phosphonium Salt-Enabled Cyclo-dearomatization Process. *Sci. China Chem.* **2020**, *63*, 1091–1099. (c) Duan, J.; Dhar, T. G. M.; Marcoux, D.; Shi, Q.; Batt, D. G.; Liu, Q.; Cherney, R. J.; Cornelius, L. A. M.; Srivastava, A. S.; Beaudoin Bertrand, M.; et al. Tricyclic Sulfones as ROR γ Modulators and Their Preparations. World Intellectual Property Organization Patent WO2016179460 A1, 2016. (d) Dong, J.-J.; Bao, J.; Zhang, X.-Y.; Xu, X.-Y.; Nong, X.-H.; Qi, S.-H. Alkaloids and citrinins from marine-derived fungus *Nigrospora oryzae* SC5GAF 0111. *Tetrahedron Lett.* **2014**, *55*, 2749–2753. (e) Metal-catalyzed reactions: (a) Zhou, P.; Yi, Y.; Hua, Y.-Z.; Jia, S.-K.; Wang, M.-C. Dinuclear Zinc Catalyzed Enantioselective Dearomatization [3 + 2] Annulation of 2-Nitrobenzofurans and 2-Nitrobenzothiophenes. *Chem.—Eur. J.* **2022**, *28*, No. e202103688. (b) Wang, Z.; Wang, D.-C.; Xie, M.-S.; Qu, G.-R.; Guo, H.-M. Enantioselective Synthesis of Fused Polycyclic Tropanes via Dearomative [3 + 2] Cycloaddition Reactions of 2-Nitrobenzofurans. *Org. Lett.* **2020**, *22*, 164–167. (c) Liang, L.; Niu, H.-Y.; Wang, D.-C.; Yang, X.-H.; Qu, G.-R.; Guo, H.-M. Facile synthesis of chiral [2,3]-fused hydrobenzofuran via asymmetric Cu(I)-catalyzed dearomative 1,3-dipolar cycloaddition. *Chem. Commun.* **2019**, *55*, 553–556.

- (d) Zhao, J.-Q.; Zhou, X.-J.; Zhou, Y.; Xu, X.-Y.; Zhang, X.-M.; Yuan, W.-C. Diastereo- and Enantioselective Dearomative [3 + 2] Cycloaddition Reaction of 2-Nitrobenzofurans with 3-Isothiocyanato Oxindoles. *Org. Lett.* **2018**, *20*, 909–912. (e) Cheng, Q.; Zhang, H.-J.; Yue, W.-J.; You, S.-L. Palladium-Catalyzed Highly Stereoselective Dearomative [3 + 2] Cycloaddition of Nitrobenzofuran. *Chem.* **2017**, *3*, 428–436. Organocatalyzed reactions: (f) Zhang, H.; He, J.; Chen, Y.; Zhuang, C.; Jiang, C.; Xiao, K.; Su, Z.; Ren, X.; Wang, T. Regio- and Stereoselective Cascade of β,γ -Unsaturated Ketones by Dipeptided Phosphonium Salt Catalysis: Stereospecific Construction of Dihydrofuro-Fused [2,3-b] Skeletons. *Angew. Chem., Int. Ed.* **2021**, *60*, 19860–19870. (g) Dou, P.-H.; Chen, Y.; You, Y.; Wang, Z.-H.; Zhao, J.-Q.; Zhou, M.-Q.; Yuan, W.-C. Organocatalyzed Asymmetric Dearomative [3 + 2] Annulation of Electron-Deficient 2-Nitrobenzo Heteroarenes with 3-Isothiocyanato Oxindoles. *Adv. Synth. Catal.* **2021**, *363*, 4047–4053. (h) Zhao, J.-Q.; Zhou, S.; Wang, Z.-H.; You, Y.; Chen, S.; Liu, X.-L.; Zhou, M.-Q.; Yuan, W.-C. Catalytic asymmetric dearomative [4 + 2] annulation of 2-nitrobenzofurans and 5H-thiazol-4-ones: stereoselective construction of dihydrobenzofuran-bridged polycyclic skeletons. *Org. Chem. Front.* **2021**, *8*, 6330–6336. (i) Zhou, X.-J.; Zhao, J.-Q.; Chen, X.-M.; Zhuo, J.-R.; Zhang, Y.-P.; Chen, Y.-Z.; Zhang, X.-M.; Xu, X.-Y.; Yuan, W.-C. Organocatalyzed Asymmetric Dearomative Aza-Michael/Michael Addition Cascade of 2-Nitrobenzofurans and 2-Nitrobenzothiophenes with 2-Aminochalcones. *J. Org. Chem.* **2019**, *84*, 4381–4391. (j) Zhao, J.-Q.; Yang, L.; Zhou, X.-J.; You, Y.; Wang, Z.-H.; Zhou, M.-Q.; Zhang, X.-M.; Xu, X.-Y.; Yuan, W.-C. Organocatalyzed Dearomative Cycloaddition of 2-Nitrobenzofurans and Isatin-Derived Morita–Baylis–Hillman Carbonates: Highly Stereoselective Construction of Cyclopenta[b]benzofuran Scaffolds. *Org. Lett.* **2019**, *21*, 660–664. (k) Yang, X.-H.; Li, J.-P.; Wang, D.-C.; Xie, M.-S.; Qu, G.-R.; Guo, H. M. Enantioselective dearomative [3 + 2] cycloaddition of 2-nitrobenzofurans with aldehyde-derived Morita–Baylis–Hillman carbonates. *Chem. Commun.* **2019**, *55*, 9144–9147.
- (10) (a) Blay, G.; Vila, C.; Martínez-Pardo, P.; Pedro, J. R. Enantioselective synthesis of five membered heterocycles through [3 + 2] Cycloadditions with Isocyano Esters. *Targets Heterocycl. Syst.* **2018**, *22*, 165–193. (b) Gulevich, A. V.; Zhdanko, A. G.; Orru, R. V. A.; Nenajdenko, V. G. Isocyanoacetate Derivatives: Synthesis, Reactivity, and Application. *Chem. Rev.* **2010**, *110*, 5235–5331.
- (11) (a) Pelkey, E. T.; Chang, L.; Gribble, G. W. An abnormal Barton-Zard reaction leading to the pyrrolo[2,3-b]indole ring system. *Chem. Commun.* **1996**, 1909–1910. (b) Pelkey, E. T.; Gribble, G. W. Synthesis and Reactions of *N*-Protected 3-Nitroindoles. *Synthesis* **1999**, *1999*, 1117–1122.
- (12) Guo, C.; Xue, M.-X.; Zhu, M.-K.; Gong, L.-Z. Organocatalytic asymmetric formal [3 + 2] cycloaddition reaction of isocyanoesters to nitroolefins leading to highly optically active dihydropyrroles. *Angew. Chem., Int. Ed.* **2008**, *47*, 3414–3417.
- (13) Wan, Q.; Xie, J.-H.; Zheng, C.; Yuan, Y.-W.; You, S.-L. Silver-Catalyzed Asymmetric Dearomatization of Electron-Deficient Heteroarenes via Interrupted Barton–Zard Reaction. *Angew. Chem., Int. Ed.* **2021**, *60*, 19730–19734.
- (14) Yuan, W.-C.; Chen, X.-M.; Zhao, J.-Q.; Zhang, Y.-P.; Wang, Z.-H.; You, Y. Ag-Catalyzed Asymmetric Interrupted Barton–Zard Reaction Enabling the Enantioselective Dearomatization of 2- and 3-Nitroindoles. *Org. Lett.* **2022**, *24*, 826–831.
- (15) (a) Martínez-Pardo, P.; Blay, G.; Muñoz, M. C.; Pedro, J. R.; Sanz-Marco, A.; Vila, C. Enantioselective synthesis of chiral oxazolines from unactivated ketones and isocyanoacetate esters by synergistic silver/organocatalysis. *Chem. Commun.* **2018**, *54*, 2862–2865. (b) Martínez-Pardo, P.; Blay, G.; Muñoz, M. C.; Pedro, J. R.; Sanz-Marco, A.; Vila, C. Enantioselective Synthesis of 5-Trifluoromethyl-2-oxazolines under Dual Silver/Organocatalysis. *J. Org. Chem.* **2019**, *84*, 314–325. (c) Martínez-Pardo, P.; Blay, G.; Escrivá-Palomo, A.; Sanz-Marco, A.; Vila, C.; Pedro, J. R. Catalytic Diastereo- and Enantioselective Synthesis of 2-Imidazolinones. *Org. Lett.* **2019**, *21*, 4063–4066. (d) Martínez-Pardo, P.; Lavios, A.; Sanz-Marco, A.; Vila, C.; Pedro, J. R.; Blay, G. Enantioselective Synthesis of Functionalized Diazaspirocycles from 4-Benzylideneisoxazol-5(4H)-one Derivatives and Isocyanoacetate Esters. *Adv. Synth. Catal.* **2020**, *362*, 3564–3569.
- (16) Full conversion of 5-acetoxy-3-nitrobenzofuran was observed. The reaction gave a complex mixture that contained ca. 30% of **5b**. Attempts to purify compound **5b** by column chromatography with different eluent systems were unsuccessful.
- (17) (a) Montesinos-Magraner, M.; Canton, R.; Vila, C.; Blay, G.; Fernandez, I.; Muñoz, M. C.; Pedro, R. C. *RSC Adv.* **2015**, *5*, 60101–60105. (b) Li, H.; Wang, Y.; Tang, L.; Wu, F.; Liu, X.; Guo, C.; Foxman, B. M.; Deng, L. *Angew. Chem., Int. Ed.* **2005**, *44*, 105–108.