

Document downloaded from:

<http://hdl.handle.net/10251/103261>

This paper must be cited as:

Martínez-Sales, Ml.; García Ximenez, F.; Espinos Gutierrez, FJ. (2016). Zebrafish as a possible bioindicator of organic pollutants in drinking waters with effects on reproduction: are effects accumulative or reversible ?. *Zygote*. 24(5):692-699.  
doi:10.1017/S0967199415000702



The final publication is available at

<https://doi.org/10.1017/S0967199415000702>

Copyright Cambridge University Press

Additional Information

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

ZEBRAFISH AS A POSSIBLE BIOINDICATOR OF ORGANIC POLLUTANTS  
IN DRINKING WATERS WITH EFFECTS ON REPRODUCTION: ARE  
EFFECTS CUMULATIVE OR REVERSIBLE?

M. Martinez-Sales<sup>1</sup>, F. García-Ximénez, FJ. Espinós.

Aquaculture and Environmental Research Group (ACUMA), Universidad  
Politécnica de Valencia, Camino de Vera 14, 46022, Valencia, Spain.

<sup>1</sup> All correspondence to: M. Martinez-Sales. Aquaculture and Environmental  
Research Group (ACUMA), Universitat Politècnica de València, Camino de Vera  
14, 46022, Valencia, Spain. Tel: +34963879433. E-mail: [mimarsa@alumni.upv.es](mailto:mimarsa@alumni.upv.es)

**Short title:**

ARE THE POLLUTANTS CUMULATIVE OR REVERSIBLE IN ZEBRAFISH?

26 ABSTRACT

27

28 Due to inefficient detection and removal treatments, organic pollutants are present in  
29 drinking waters. For this reason, zebrafish is proposed as a complementary control  
30 measure in conventional potabilization treatments.

31

32 According to the most sensitive parameters (hatching rate, fertility rate and  
33 underdeveloped specimens) detected in our previous work, in the current work we  
34 attempt to study, in these parameters, the possible cumulative effect of  
35 environmental pollutants likely present in drinking waters, between generations,  
36 when specimens are cultured in the same water in both generations and/or the  
37 possible reversibility of these effects when cultured in control water.

38

39 To this end, batches of 20 embryos with the chorion intact were cultured in 3  
40 drinking waters from different sources (A, B and C) and in one control water up to 5  
41 months, in 20 l tanks. Four replicates were performed in all water groups, with a  
42 total of 28 aquariums.

43

44 Results in water C revealed a non-reversible effect on fertility rate, and also in water  
45 C an alteration of sex ratio towards females, although in this case the alteration was  
46 reversible. A transgenerational alteration in the germline via epigenetic mechanism  
47 from the previous generation is proposed as the most plausible explanation to this  
48 effect.

49

50 Keywords: organic pollutants; bioindicator; epigenetic; drinking water; zebrafish.

51

## 52 1. INTRODUCTION

53

54 Organic pollutants such as pharmaceutical and medical substances and persistent  
55 organic pollutants (POPs) have been dispersed worldwide and as a result are  
56 emerging in surface, groundwater and even in drinking waters, in this case due to  
57 inefficient removal treatments (Ikehata et al., 2008; Benner et al., 2013). The  
58 concentrations of these substances are low but increasingly numerous (year by year)  
59 and variable over time (Khetan and Collins, 2007; Rodil et al., 2012). These  
60 substances can exert toxicological but also epigenetic effects on many functions,  
61 operating on somatic cells and in the germ line, in this case promoting  
62 transgenerational effects (Rusiecki et al., 2008; Skinner, 2011).

63

64 In our previous work (Martínez-Sales et al., 2015), we defined and narrowed the  
65 most sensitive developmental and reproductive parameters in zebrafish, with the  
66 long-term aim of establishing the zebrafish as a bioindicator of the possible presence  
67 of environmental pollutants. Specifically, the assessment was carried out in three  
68 drinking waters from different tap water sources. The most sensitive parameters  
69 detected were: hatching rate, fertility rate and underdeveloped specimens. So, in the  
70 present work we focused on these parameters in order to study the possible  
71 cumulative effect and/or possible reversibility of the effects, between generations, of  
72 these environmental pollutants in the same three drinking waters (A, B and C) in  
73 both generations, despite the fact that there are other sensitive parameters, for  
74 example sex ratio.

75

76 2. MATERIAL AND METHODS

77

78 *Zebrafish maintenance*

79 Both F0 obtained from the original wild zebrafish colony and F1 generations were  
80 reared in the laboratory following the protocol described in Westerfield (1995).  
81 Briefly, adult zebrafish were kept in 20 L tanks at 28.5°C, in a 3:2 ratio (females:  
82 males) (Westerfield, 2007) and fed on granular food supplemented with recently  
83 defrosted hen egg yolk and shrimp meat (Simão et al., 2010 a). The light cycle was  
84 regulated at 14h light/ 10h dark (Matthews et al., 2002; Brand et al., 2002). The  
85 aquariums had water recirculation systems but without active carbon filters.  
86 According to the Westerfield (2007) recommendations, a quarter of the total  
87 aquarium water was removed weekly and replaced by clean water to avoid  
88 ammonium concentrations.

89

90 It must be stated that all environmental conditions were identical to all aquariums  
91 and the spatial distribution of the aquariums was randomized.

92

93 *Water sources*

94

95 The four different drinking waters used in the present study (the same than in our  
96 previous work) were classified depending on their source into: three waters from  
97 different tap water distribution networks (A, B and C) and one bottled spring water  
98 which was established as a control. Type A was tap water from a city located in a  
99 region with intensive farming activity, from the hydrological basin of the Túrria

100 river. Type B was from the tap water distribution network of a medium-sized city,  
101 supplied from the Túrria and Xúquer rivers. Finally, type C was tap water from a city  
102 also located in a region with intensive agricultural activity, but from the  
103 hydrological basin of the river Xúquer. Type A and C came from groundwater  
104 prospecting.

105

106 Before filling the aquariums with water, recipients (where the water was stored)  
107 were kept open for at least a week, with a large exchange surface to favour chlorine  
108 elimination (Westerfield, 1995).

109

110 It should be mentioned that all the waters are potable and also that the chemical  
111 parameters defined for tap water for human consumption in Royal Decree 140/2003  
112 of 7 February, which establishes the health criteria for the quality of water intended  
113 for human consumption, are suitable for zebrafish breeding and maintenance  
114 (Westerfield, 2007).

115

### 116 *Specimen management*

117

118 Fertilized embryos were obtained by siphoning. Batches of 20 fertilized embryos at  
119 the Mid Blastula Transition (MBT) stage with the chorion intact (Martinez-Sales et  
120 al., 2014; Martinez-Sales et al., 2015) were selected under a stereo microscope  
121 between those degenerated and those that initiated aberrant parthenogenetic  
122 development. These embryos were left in Petri dishes and cultured until 5 dpf (days  
123 post fertilization) at 28, 5°C in dishes with the same water type where their  
124 progenitors were reared (same water origin and water destination: A-A; B-B; C-C;

125 Control-Control) and, on the other hand, in dishes with control water (different  
126 water origin and water destination: A-control; B-control; C-control).

127

128 Next, from 5 dpf to complete adulthood (5 months post fertilization) larvae were left  
129 in aquariums (20 L) in the same type of water as that in which their progenitors were  
130 reared and in aquariums with control water, to assess either the possible cumulative  
131 effect when specimens are cultured in the same water or the possible reversibility  
132 effect when are cultured in control water. From these combinations, four replicates  
133 were established with a total of 28 aquariums.

134

135 After three months, marbles were placed in each aquarium with the aim of siphoning  
136 all aquariums 2 or 3 times a week throughout the 4<sup>th</sup> and the 5<sup>th</sup> month, to evaluate  
137 the onset of spawning and the fertility rate. Sex ratio of the surviving adults,  
138 underdeveloped specimens and survival and abnormality rates at 5 mpf were also  
139 evaluated. Moreover, in the F1 offspring (F2 larvae) we evaluated the survival and  
140 abnormality rates at 5 dpf and the hatching rate at 72 hpf (hours post fertilization).

141

142 The experimental procedures and animal care in this work fully comply with the  
143 standards for use of animals established by the Ethical Committee of the Polytechnic  
144 University of Valencia, which specifically approved this study.

145

### 146 *Experimental design*

147

148 Two different analyses were carried out on the most sensitive parameters obtained in  
149 our previous work: hatching rate, fertility rate and underdeveloped specimens. The

150 first analysis studied the possible cumulative effect between generations. To this  
151 end, fertility rate and underdeveloped specimens (runts) were compared in the F0  
152 and F1 generation. In turn, the hatching rate at 72 hpf was compared in the F1 and  
153 F2 generation. The second analysis studied the possible reversibility of the effects in  
154 fertility rate and in underdeveloped specimens in the F1 generation, and hatching  
155 rate in the F2 generation (see figure 1).

156

### 157 *Statistical analysis*

158

159 The possible cumulative and reversible effects in all parameters were analysed using  
160 Chi-square test (Statgraphics Plus 5.1). The Yates correction for continuity was used  
161 when a single degree of freedom was involved. Values were considered statistically  
162 different at  $P < 0.05$ .

163

## 164 3. RESULTS

165

166 As stated in material and methods, four replicates were performed in all water  
167 groups with a total of 28 aquariums at the outset. However, 8 aquariums were  
168 discarded due to total mortality of the larvae cultured in Petri dishes until 5dpf for  
169 reasons unknown and uncontrolled. This mortality cannot be associated to a water  
170 type, as the mortality was random between groups. So, the minimum number of  
171 replicates per group was two, with a total of 20 aquariums. In the first group  
172 (control-control) the final number of replicates was three, in the second group (A-A)  
173 the final number of replicates was two, in the third group (A-control) the final  
174 number of replicates was also two, in the fourth group (C-C) the final number was

175 three, in the fifth group (C- control) the final number was four, in the sixth group  
176 (B-B) the final number was two and in the seventh group (B-control) the final  
177 number was four.

178

### 179 3.1.- Hatching rate

180

181 Hatching rate was evaluated at 72 hpf (Martinez-Sales et al., 2015) in the F1 and F2  
182 generations during 4<sup>th</sup> and 5<sup>th</sup> mpf.

183

### 184 *Cumulative effect*

185

186 The analysis showed statistically significant differences ( $p<0.05$ ) between the F1  
187 and the F2 generations in all waters studied (see table 1). In all cases, the worst  
188 results were obtained in the second generation. These results reveal a cumulative  
189 effect in all waters, even in the control water. The negative cumulative effect in the  
190 case of water B should be highlighted.

191

### 192 *Reversible effect*

193

194 The analysis showed statistically significant differences ( $p<0.05$ ) between data from  
195 the specimens reared in waters with the same origin and destination and data from  
196 the specimens reared in control water in all waters studied (see tables 2, 3 and 4).  
197 The worst result was obtained in all waters with the same origin and destination.  
198 These results reveal that there was a reversible effect in all waters when specimens  
199 were cultured in control water.

200

201 3.2.- Fertility rate

202

203 Fertility rate was evaluated through 4<sup>th</sup> and 5<sup>th</sup> mpf in the F0 and F1 generations.

204

205 ***Cumulative effect***

206

207 The analysis showed statistically significant differences ( $p < 0.05$ ) between the F0  
208 and the F1 generations in all waters studied (see table 5). The worst results were  
209 obtained in the second generation (F1). These results reveal a cumulative effect in  
210 all waters, including the control water.

211

212 ***Reversible effect***

213

214 The analysis showed statistically significant differences ( $p < 0.05$ ) between data from  
215 specimens reared in waters with the same origin and destination and data from  
216 specimens reared in control water in all waters studied (see table 6, 7 and 8). In the  
217 case of waters A and B, the worst result was obtained in waters with the same origin  
218 and destination (A-A and B-B), whereas in water C the result did not improve when  
219 specimens were cultured in control water. These results revealed that there was a  
220 reversible effect in waters A and B when specimens were cultured in control water,  
221 but a non-reversible effect in water C.

222

223 3.3.- Underdeveloped specimens (runts)

224

225 In this second work, specimens evaluated at 5 mpf in the F1 generation were all  
226 sexes clearly identifiable, and morphologically were also similar. Hence, there were  
227 no underdeveloped specimens.

228

### 229 3.4.- Sex ratio

230

231 Even though in the previous work sex ratio was not a sensitive parameter, in the  
232 present work, water C displayed a feminization process. Therefore, sex ratio in  
233 water C was analysed at 5mpf in the F0 and in the F1 generations.

234

### 235 ***Cumulative effect***

236

237 The analysis showed statistically significant differences ( $p < 0.05$ ) between water C  
238 from F0 and water C from F1. The worst result was obtained in water C from F1,  
239 where the sex ratio was skewed towards females (males 25%: females 75%) (see  
240 table 9). No significant difference ( $p > 0.05$ ) was obtained in the other waters (A and  
241 B) whose sex ratio percentages were within the normal range in zebrafish in both  
242 generations (60 males: 40 females) (Fenske et al., 1999).

243

### 244 ***Reversible effect***

245

246 The feminization detected in specimens cultured in water C, disappeared when were  
247 reared in control water (see table 10).

248

## 249 4. DISCUSSION

250

251 Based upon results from our previous work (Martínez-Sales et al., 2015), hatching  
252 rate, fertility rate and underdeveloped specimens were the most sensitive parameters  
253 to detect the possible presence of environmental pollutants in drinking waters from  
254 different tap water distribution networks (A, B and C). These parameters were  
255 selected considering the full life-cycle (from development to reproduction) of  
256 zebrafish specimens.

257

258 The same waters were used in the present work, but it should be taken into account  
259 that although these waters have the same original source, the physical and chemical  
260 conditions of the water may have changed due to seasonal variations in quality at  
261 the water source (Ouyang et al., 2006), although in order to be drinkable it should  
262 meet legal strict limits. Nonetheless, differences between waters also appeared in the  
263 same parameters in this experiment, except in the rate of underdeveloped specimens.

264

265 The period around hatching is a critical stage during embryogenesis (Henn, 2011),  
266 which is why the hatching rate has been extensively used as a parameter in many  
267 toxicological studies (Han et al., 2011; Galus et al., 2013) as well as a parameter for  
268 reproductive toxicity assessment (Simon et al., 2011). Our results for hatching rate  
269 revealed that although the results were high in all waters in both generations, except  
270 in water B (86.47% in F1 and 37.5% in F2), there was a negative cumulative effect  
271 in the second generation in all waters tested, even in the control water. Surprisingly,  
272 water B reached the worst results in both generations compared to the control water,  
273 decreasing to 48.97% (86.47%-37.5%) in the second generation compared to the  
274 first. These outcomes may suggest either the possible increasing presence of the  
275 same pollutants in waters in both experiments (generations) which affect the

276 hatching process and/or the possible transmission of these negative effects to the  
277 next generation via epigenetic mechanisms (Skinner et al., 2010; Skinner, 2011).  
278 However, it should be stated that when specimens were cultured in control water,  
279 this cumulative effect disappeared, which rules out a possible transgenerational  
280 transmission via epigenetic mechanisms.

281

282 Fertility rate has also been used in many toxicological studies as a good parameter  
283 (Ankley and Johnson, 2004; Liu et al., 2014). Results from fertility show that there  
284 was a negative cumulative effect in the second generation compared to the first in all  
285 waters, even in the control water. The most pronounced reduction between  
286 generations was obtained in water A, 22.28% (42.60%-20.32%), as this water  
287 reached the lowest rate (20.32%), followed by water B (24.5%) in the second  
288 generation. These outcomes may suggest either the possible increasing presence of  
289 the same pollutants in waters in both experiments (generations), which affected the  
290 fertility rate and/or the possible transgenerational transmission of these negative  
291 effects to the next generation via epigenetic mechanisms (Skinner et al., 2010;  
292 Skinner, 2011). It should be noted that when specimens were cultured in control  
293 water, there was a reversible effect in waters A and B, which ruled out a possible  
294 transgenerational transmission via epigenetic mechanism in these waters, although  
295 the cumulative effect remained in water C, the fertility rate decreasing to 12.03%  
296 (43.03% -31%) when specimens were cultured in control water.

297

298 So, on the basis of these findings we posit the possible presence of environmental  
299 pollutants in water A and B that affect fertility rate in both generations without  
300 transgenerational transmission, due to the reversibility process in these waters.

301 Nevertheless, in water C the non-reversible effect also leads us to consider the  
302 possible presence of environmental pollutants in water C that affect fertility rate in  
303 both generations, but in this case with a possible transgenerational transmission due  
304 to the maintenance of the cumulative effects when specimens were cultured later in  
305 control water. This could be explained because early exposure during critical  
306 periods of development to environmental pollutants, such as endocrine disruptors  
307 (Braw-Tal, 2010), can promote an adult-onset alteration (in this case a reduction in  
308 fertility rate) long after the compound is removed, even in subsequent generations if  
309 the germ line is affected through epigenetic mechanisms (Skinner et al., 2010;  
310 Skinner, 2011).

311

312 Regarding the non-reversible effect of the fertility rate in water C, although we are  
313 unable to describe the mechanism of action behind this effect, a plausible  
314 explanation could be an early exposure to some pollutant in water C during a critical  
315 period of embryo development (Braw-Tal, 2010), such as the MBT stage in our  
316 case, without a germline alteration via epigenetic mechanism, as the crucial period  
317 for epigenetic regulation and modification of the germline is during the period of  
318 primordial germ cell migration and gonadal sex determination (Skinner et al., 2010),  
319 events that take place after the MBT stage (3 hpf) (Dahm, 2002), at the early  
320 gastrulation stage (from 6 hpf) (Yoshizaki et al., 2002). So, taking this argument  
321 into account, the most likely explanation could be an alteration in the germline  
322 transgenerational transmitted from the previous generation (parents) via epigenetic  
323 mechanisms to this generation.

324

325 Sex ratio is a relevant parameter used in many toxicological studies (Hill and Janz,  
326 2003; Baumann et al., 2013; Liu et al., 2014). However, in our previous work, it was  
327 not classified as a sensitive parameter because in all drinking waters tested sex ratios  
328 were within the normal ranges. Thus, all percentages of females were around 40%,  
329 which agreed with our current results and with other studies on zebrafish (60 males:  
330 40 females) (Fenske et al., 1999), (68:32) (Örn et al., 2003), (56:44) (Vaughan et al.,  
331 2001; Hsioa and Tsai, 2003). However, in this second experiment in water C there  
332 was an alteration of sex ratio towards females (75%), although this feminization  
333 changed towards normal values in zebrafish when specimens were cultured in  
334 control water.

335

336 These results suggests the possible presence of some environmental pollutants, only  
337 in water C, such as endocrine disrupting chemicals (17-ethinylestradiol, even at  
338 ng/l) that can disrupt sexual differentiation in fish (Larsen et al., 2009) and cause  
339 feminization and retardation of sexual maturation in zebrafish. These substances  
340 may trigger disruption of sex hormones during sexual development and alter female  
341 sex, male sex or even both sexes. In fish, the hormonal balance between estrogens  
342 and androgens appears to be an important factor in the course of sexual  
343 differentiation (Liu et al., 2014).

344

345 It must be highlighted that all environmental factors were rigorously controlled to  
346 avoid any external alteration of our sex differentiation in zebrafish, as this is known  
347 to be a difficult process in fish (Liew et al., 2014) that can be affected by several  
348 environmental factors in a very complex way (Baroiller et al., 1999).

349

350 Evidence from our results gathered to date corroborates that zebrafish is a suitable  
351 model for use as a bioindicator to detect environmental pollutants in drinking water.  
352 The complexity of detecting these substances in conventional potabilization  
353 treatments, due to their interactions and their variable and random presence even at  
354 low levels in drinking water, makes their routine chemical detection and control  
355 difficult or even impossible (Khetan and Collins, 2007; Benner et al., 2013). For this  
356 reason, bioindicators could be used as backup control measures to conventional  
357 potabilization treatments.

358

359 Finally, the detection in our previous (Martinez-Sales et al., 2015) and current works  
360 of the negative effects on reproductive parameters in zebrafish reared in drinkable  
361 water is cause for alarm, as the presence of these substances in drinking water may  
362 be one of the reasons behind the decline in human reproduction in metropolitan  
363 areas (Toft et al., 2006; Jurewicz et al., 2009; Braw-Tal, 2010; Vested et al., 2014).

364

365

366

367 5. DECLARATION OF INTEREST

368

369 The authors declare that there is no conflict of interest that could be perceived as  
370 prejudicing the impartiality of the research reported.

371

372 6. FUNDING

373

374 This research did not receive any specific grant from any funding agency in the  
375 public, commercial or not-for-profit sector.

376

377 7. ACKNOWLEDGMENTS

378

379 The authors would like to thank Mr. Javier Rubio Rubio for his valuable technical  
380 support and Mr. Neil Macowan for improving the English of this manuscript.

381

382

383 8. REFERENCES

384

385 Ankley GT & Johnson RD, 2004. Small fish models for identifying and assessing  
386 the effects of endocrine-disrupting chemicals. Institute for Laboratory Animal  
387 Research, 45: 469-483.

388 Baroiller JF, Guiguen Y & Fostier A, 1999. Endocrine and environmental aspects  
389 of sex differentiation in fish. Cellular and Molecular Life Sciences, 55: 910-931.

390 Baumann L, Holbech H, Keiter S, Kinnberg KL, Knörr S, Nagel T & Braunbeck T,  
391 2013. The maturity index as a tool to facilitate the interpretation of changes in  
392 vitellogenin production and sex ratio in the Fish Sexual Development Test.  
393 Aquatic Toxicology, 128-129: 34-42.

394 Benner J, Helbling DE, Kohler HPE, Wittebol J, Kaiser E, Prasse C, et al., 2013. Is  
395 biological treatment a viable alternative for micropollutant removal in drinking  
396 water treatment processes?. Water research, 47(16): 5955-5976.

397 Brand M, Granato M & Nüslein-Volhard C, 2002. Keeping and raising zebrafish.  
398 In: Zebrafish. A Practical Approach, pp. 16. Eds Nüslein-Volhard C and Dahm  
399 R. Oxford, England: Oxford University Press.

400 Braw-Tal R, 2010. Endocrine disruptors and timing of human exposure. Pediatric  
401 Endocrinology Reviews (PER), 8(1): 41-46.

402 Dahm R, 2002. Atlas of embryonic stages of development in the zebrafish  
403 (Appendix 2). In: Zebrafish. A Practical Approach, pp. 219-236. Eds Nüslein-  
404 Volhard C and Dahm R. Oxford, England: Oxford University Press.

405 Fenske M, Maack G, Ensenbach U & Segner H, 1999. Identification of estrogens-  
406 sensitive developmental stages of zebrafish, *Danio rerio*. Proceedings of the  
407 Ninth Annual Meeting of Society of Environmental Toxicology and Chemistry,  
408 25-29.

409 Galus M, Kirischian N, Higgins S, Purdy J, Chow J, Rangaranjan S, Li H, Metcalfe  
410 C & Wilson JY, 2013. Chronic, low concentration exposure to pharmaceuticals  
411 impacts multiple organ systems in zebrafish. Aquatic Toxicology, 132-133: 200-  
412 211.

413 Han Z, Jiao S, Shan Z & Zhang X, 2011. Effects of  $\beta$ -endosulfan on the growth and  
414 reproduction of zebrafish (*Danio rerio*). Environmental Toxicology and  
415 Chemistry, 30(11): 2525–2531.

416 Henn K, 2011. Limits of the embryo toxicity test with *Danio rerio* as an alternative  
417 to the acute fish toxicity test. Ph Thesis. University of Heidelberg, Germany.

418 Hill R & Janz D, 2003. Developmental estrogenic exposure in zebrafish (*Danio*  
419 *rerio*). I. Effects on sex ratio and breeding success. *Aquatic Toxicology*, 63: 417-  
420 429.

421 Hsiao CD & Tsai HJ, 2003. Transgenic zebrafish with fluorescent germ cell: a  
422 useful tool to visualize germ cell proliferation and juvenile hermaphroditism in  
423 vivo. *Developmental Biology*, 262: 313-323.

424 Ikehata K, Gamal El-Din M & Snyder SA, 2008. Ozonation and advanced  
425 oxidation treatment of emerging organic pollutants in water and wastewater.  
426 *Ozone: Science and Engineering*, 30(1): 21-26.

427 Jurewicz J, Hanke W, Radwan M & Bonde J, 2009. Environmental factors and  
428 semen quality. *International journal of occupational medicine and environmental*  
429 *health*, 22(4): 305-329.

430 Khetan SK & Collins TJ, 2007. Human Pharmaceuticals in the Aquatic  
431 Environment: A Challenge to Green Chemistry. *Chemical Reviews*, 107: 2319-  
432 2364.

433 Larsen MG, Bilberg K & Baatrup E, 2009. Reversibility of estrogenic sex changes  
434 in zebrafish (*Danio Rerio*). *Environmental Toxicology and Chemistry*, 28: 1789-  
435 1785.

436 Liew WC & Orbán L, 2014. Zebrafish sex: a complicated affair. *Briefings in*  
437 *functional genomics*, 13(2): 172-187.

438 Liu SY, Jin Q, Huang XH & Zhu GN, 2014. Disruption of zebrafish (*Danio rerio*)  
439 sexual development after full life-cycle exposure to environmental levels of  
440 triadimefon. *Environmental toxicology and pharmacology*, 37(1): 468-475.

441 Martinez-Sales M, García- Ximénez F & Espinós FJ, 2014. Zebrafish (*Danio rerio*)  
442 as a possible bioindicator of epigenetic factors present in drinking water that  
443 may affect reproductive function: Is chorion an issue? *Zygote*, 1-6.

444 Martinez-Sales M, García- Ximénez F & Espinós FJ, 2015. Zebrafish as a possible  
445 bioindicator of organic pollutants with effects on reproduction in drinking  
446 waters. *Journal of environmental sciences*. In press.

447 Matthews M, Trevarrow B & Matthews J, 2002. A virtual tour of the Guide for  
448 zebrafish users. *Laboratory Animal*, 31(3): 34-40.

449 Örn S, Holbech H, Madsen TH, Norrgren L & Petersen GI, 2003. Gonad  
450 development and vitellogenin production in zebrafish (*Danio rerio*) exposed to  
451 ethinylestradiol and methyl testosterone. *Aquatic Toxicology*, 65: 397–411.

452 Ouyang Y, Nkedi-Kizza P, Wu QT, Shinde D & Huang CH, 2006. Assessment of  
453 seasonal variations in surface water quality. *Water research*, 40(20): 3800-3810.

454 Rodil R, Quintana JB, Concha-Graña E, López-Mahía P, Muniategui-Lorenzo S &  
455 Prada-Rodríguez D, 2012. Emerging pollutants in sewage, surface and drinking  
456 water in Galicia (NW Spain). *Chemosphere*, 86(10): 1040-1049.

457 Rusiecki JA, Baccarelli A, Bollati V, Tarantini L, Moore LE & Bonfeld-Jorgensen  
458 EC, 2008. Global DNA hypomethylation is associated with high serum-  
459 persistent organic pollutants in Greenlandic Inuit. *Environ Health Perspect*,  
460 116(11): 1547-1552.

461 Simão M, Cardona-Costa J, Pérez Camps M & García-Ximénez F, 2010 a.  
462 Ultraviolet Radiation Dose to be Applied in Recipient Zebrafish Embryos for  
463 Germ-line Chimaerism is Strain Dependent. *Reproduction in Domestic Animals*,  
464 45: 1098-1103.

465 Simon O, Mottin E, Geffroy B & Hinton T, 2011. Effects of dietary uranium on  
466 reproductive endpoints—fecundity, survival, reproductive success—of the fish  
467 *Danio rerio*. *Environmental Toxicology and Chemistry*, 30(1): 220-225.

468 Skinner MK, Manikkam M & Guerrero-Bosagna C, 2010. Epigenetic  
469 transgenerational actions of environmental factors in disease etiology. *Trends in*  
470 *Endocrinology & Metabolism*, 21(4): 214-222.

471 Skinner MK, 2011. Role of epigenetics in developmental biology and  
472 transgenerational inheritance. *Birth Defects Research Part C: Embryo Today:*  
473 *Reviews*, 93(1): 51-55.

474 Toft G, Rignell-Hydbom A, Tyrkiel E, Shvets M, Giwercman A, Lindh CH, et al.,  
475 2006. Semen quality and exposure to persistent organochlorine pollutants.  
476 *Epidemiology*, 17(4): 450-458.

477 Vaughan M, Van Egmond R & Tyler CR, 2001. The effect of temperature on  
478 sexual differentiation and development in the zebrafish (*Danio rerio*).  
479 *Proceedings of the 11th Annual Meeting of the Society of Environmental*  
480 *Toxicology and Chemistry*, 6-10.

481 Vested A, Giwercman A, Bonde JP, & Toft G, 2014. Persistent organic pollutants  
482 and male reproductive health. *Asian journal of andrology*, 16(1): 71.

483 Westerfield M, 1995. The zebrafish book. University of Oregon, Eugene, OR.  
484 Westerfield M, 2007. The Zebrafish book: A guide for the laboratory use of  
485 zebrafish (*Danio rerio*). Edn 5. University of Oregon Press, Eugene, US.  
486 Yoshizaki G, Takeuchi Y, Kobayashi T, Ihara S & Takeuchi T, 2002. Primordial  
487 germ cells: the blueprint for a piscine life. *Fish Physiology and Biochemistry*,  
488 26(1): 3-12.  
489