

Document downloaded from:

<http://hdl.handle.net/10251/107410>

This paper must be cited as:

Albanese, A.; Bonet Solves, JA.; Ricker, WJ. (2018). The Fréchet space  $ces(p+)$ ,  $1 < p < \infty$ . *Journal of Mathematical Analysis and Applications*. 458(2):1314-1323.  
doi:10.1016/j.jmaa.2017.10.024



The final publication is available at

<http://doi.org/10.1016/j.jmaa.2017.10.024>

Copyright Elsevier

Additional Information

# THE FRÉCHET SPACES $\text{ces}(p+)$ , $1 < p < \infty$

ANGELA A. ALBANESE, JOSÉ BONET, WERNER J. RICKER

ABSTRACT. The Banach spaces  $\text{ces}(p)$ ,  $1 < p < \infty$ , were intensively studied by G. Bennett and others. The *largest* solid Banach lattice in  $\mathbb{C}^{\mathbb{N}}$  which contains  $\ell_p$  and which the Cesàro operator  $\mathbf{C} : \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$  maps into  $\ell_p$  is  $\text{ces}(p)$ . For each  $1 \leq p < \infty$ , the (positive) operator  $\mathbf{C}$  also maps the Fréchet space  $\ell_{p+} = \bigcap_{q>p} \ell_q$  into itself. It is shown that the *largest* solid Fréchet lattice in  $\mathbb{C}^{\mathbb{N}}$  which contains  $\ell_{p+}$  and which  $\mathbf{C}$  maps into  $\ell_{p+}$  is precisely  $\text{ces}(p+) := \bigcap_{q>p} \text{ces}(q)$ . Although the spaces  $\ell_{p+}$  are well understood, it seems that the spaces  $\text{ces}(p+)$  have not been considered at all. A detailed study of the Fréchet spaces  $\text{ces}(p+)$ ,  $1 \leq p < \infty$ , is undertaken. They are very different to the Fréchet spaces  $\ell_{p+}$  which generate them in the above sense. We prove that each  $\text{ces}(p+)$  is a power series space of finite type and order one, and that all the spaces  $\text{ces}(p+)$ ,  $1 \leq p < \infty$ , are isomorphic.

## 1. INTRODUCTION

Given an element  $x = (x_n)_n = (x_1, x_2, \dots)$  of  $\mathbb{C}^{\mathbb{N}}$  let  $|x| := (|x_n|)_n$  and write  $x \geq 0$  if  $x = |x|$ . By  $x \leq y$  we mean that  $(y - x) \geq 0$ . The sequence space  $\mathbb{C}^{\mathbb{N}}$  is a (locally convex) Fréchet space with respect to the coordinatewise convergence. For each  $1 < p < \infty$  define

$$\text{ces}(p) := \{x \in \mathbb{C}^{\mathbb{N}} : \|x\|_{\text{ces}(p)} := \left\| \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)_n \right\|_p < \infty\}, \quad (1.1)$$

where  $\|\cdot\|_p$  denotes the standard norm in  $\ell_p$ . An intensive study of the Banach spaces  $\text{ces}(p)$ ,  $1 < p < \infty$ , was undertaken in [6],[13]; see also the references therein. They are reflexive,  $p$ -concave Banach lattices (for the order induced by the positive cone of the Fréchet lattice  $\mathbb{C}^{\mathbb{N}}$ ) and the canonical vectors  $e_k := (\delta_{nk})_n$ , for  $k \in \mathbb{N}$ , form an unconditional basis, [6], [8]. For every pair  $1 < p, q < \infty$  the space  $\text{ces}(p)$  is *not* isomorphic to  $\ell_q$ , [6, Proposition 15.13], and is also *not* isomorphic to  $\text{ces}(q)$  if  $p \neq q$ , [4, Proposition 3.3].

The Cesàro operator  $\mathbf{C} : \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ , defined by

$$\mathbf{C}(x) := \left( x_1, \frac{x_1 + x_2}{2}, \dots, \frac{x_1 + x_2 + \dots + x_n}{n}, \dots \right), \quad x \in \mathbb{C}^{\mathbb{N}}, \quad (1.2)$$

---

*Key words and phrases.* Fréchet spaces, sequence spaces, power series spaces, Schwartz spaces, Fréchet lattices.

*Mathematics Subject Classification 2010:* Primary 46A45, 46A04; Secondary 46A11, 46A40, 46A63, 47B37.

satisfies  $|\mathbf{C}(x)| \leq \mathbf{C}(|x|)$  for  $x \in \mathbb{C}^{\mathbb{N}}$  and is a topological isomorphism of  $\mathbb{C}^{\mathbb{N}}$  onto itself. It is clear from (1.1) that  $\|x\|_{ces(p)} = \|\mathbf{C}(|x|)\|_p$  for  $x \in ces(p)$ . Hardy's inequality, [15, Theorem 326], ensures that  $\ell_p \subseteq ces(p)$  with  $\|x\|_{ces(p)} \leq p'\|x\|_p$  for  $x \in \ell_p$ , where  $\frac{1}{p} + \frac{1}{p'} = 1$ . Moreover,  $\ell_p \subseteq ces(p)$  is a *proper* containment, [8, Remark 2.2]. It is routine to verify that  $\mathbf{C}$  maps  $ces(p)$  continuously into  $\ell_p$ . The following remarkable fact (due to Bennett, [6, Theorem 20.31]) reveals a special feature of  $ces(p)$ .

**Proposition 1.1.** *Let  $1 < p < \infty$  and  $x \in \mathbb{C}^{\mathbb{N}}$ . Then*

$$x \in ces(p) \text{ if and only if } \mathbf{C}(|x|) \in ces(p). \quad (1.3)$$

The spaces  $ces(p)$  also arise in a different way. Fix  $1 < p < \infty$ . Since the Cesàro operator  $\mathbf{C}_p : \ell_p \rightarrow \ell_p$  (i.e.,  $\mathbf{C}$  restricted to  $\ell_p$ ) is a *positive* operator between Banach lattices, it is natural to seek continuous,  $\ell_p$ -valued extensions of  $\mathbf{C}_p$  to Banach lattices  $X \subseteq \mathbb{C}^{\mathbb{N}}$  which are larger than  $\ell_p$  and *solid* (i.e.,  $y \in \mathbb{C}^{\mathbb{N}}$  and  $|y| \leq |x|$  with  $x \in X$  implies that  $y \in X$ ). The *largest* of all those solid Banach lattices in  $\mathbb{C}^{\mathbb{N}}$  which contain  $\ell_p$  and for which such a continuous,  $\ell_p$ -valued extension of  $\mathbf{C}_p$  is possible is precisely  $ces(p)$ , [8, p.62]. Of course, this “largest extension”  $\mathbf{C} : ces(p) \rightarrow \ell_p$  is the restriction of  $\mathbf{C}$  from  $\mathbb{C}^{\mathbb{N}}$  to  $ces(p)$ .

For each  $1 \leq p < \infty$  define the vector space  $\ell_{p+} := \bigcap_{q>p} \ell_q$ ; it is a Fréchet space (and lattice for the order induced by the positive cone of  $\mathbb{C}^{\mathbb{N}}$ ) with respect to the increasing sequence of *lattice norms*

$$x \mapsto \|x\|_{p_k}, \quad x \in \ell_{p+}, \quad k \in \mathbb{N}, \quad (1.4)$$

for any sequence  $p < p_{k+1} < p_k$  with  $p_k \downarrow p$ . Moreover, each  $\ell_{p+} \subseteq \mathbb{C}^{\mathbb{N}}$  (with a continuous inclusion) is a reflexive, quasinormable, non-Montel, countably normed Fréchet space which is solid in  $\mathbb{C}^{\mathbb{N}}$  and contains no isomorphic copy of any infinite dimensional Banach space, [9], [18]. Clearly, for each  $1 < p < \infty$  the Banach space  $\ell_p \subseteq \ell_{p+}$  continuously and with a proper inclusion. Since  $\mathbf{C}_p$  is continuous for each  $1 < p < \infty$  (with operator norm  $p'$  where  $\frac{1}{p} + \frac{1}{p'} = 1$ , [15, Theorem 326]), it follows that  $\mathbf{C} : \ell_{p+} \rightarrow \ell_{p+}$  is also continuous, [3, Section 2]. The natural question is: To what extent do the properties and interrelations between the Banach spaces  $\ell_p$  and  $ces(p)$ ,  $1 < p < \infty$ , alluded to above reflect themselves in the connections between the corresponding Fréchet spaces  $\ell_{p+}$  and  $ces(p+) := \bigcap_{q>p} ces(q)$  which they generate? Although the Fréchet spaces  $\ell_{p+}$ ,  $1 \leq p < \infty$ , are well understood (see eg. [1], [9], [10], [18], [19] and the references therein), it seems that the Fréchet spaces  $ces(p+)$ ,  $1 \leq p < \infty$ , which are equipped with the *lattice norms*

$$x \mapsto \|x\|_{ces(p_k)}, \quad x \in ces(p+), \quad k \in \mathbb{N},$$

for any sequence  $p < p_{k+1} < p_k$  satisfying  $\lim_{k \rightarrow \infty} p_k = p$  (i.e.,  $ces(p+) = \text{proj}_k ces(p_k)$ ), have not been considered at all. The aim of this note is to make a detailed study of these spaces and to expose some of their striking features. Let us describe some sample results.

First, just like for  $C_p : \ell_p \longrightarrow \ell_p$ , for  $1 < p < \infty$ , the Cesàro operator  $C_{p+} : \ell_{p+} \longrightarrow \ell_{p+}$ , for  $1 \leq p < \infty$ , is also a *positive* operator, albeit now between Fréchet lattices. It turns out that the *largest* of all those solid Fréchet lattices in  $\mathbb{C}^{\mathbb{N}}$  which contain  $\ell_{p+}$  and  $C$  maps into  $\ell_{p+}$  (necessarily continuously) is precisely  $ces(p+)$ ; see Proposition 2.5. Although each Fréchet space  $\ell_{p+}$ , for  $1 \leq p < \infty$ , *fails* to have the property (1.3) of Proposition 1.1 (with  $\ell_{p+}$  in place of  $ces(p)$ ), the space  $ces(p+)$  that it generates in the above sense *does* have this remarkable property; see Propositions 2.2 and 2.4. A further contrast to  $\ell_{p+}$  is that each  $ces(p+)$ ,  $1 \leq p < \infty$ , is a Fréchet-Schwartz space (but, not nuclear) and the canonical vectors  $\{e_k : k \in \mathbb{N}\}$  form an unconditional basis (cf. Proposition 3.5). In particular,  $ces(p+)$  cannot be isomorphic to any of the non-Montel spaces  $\ell_{q+}$ ,  $1 \leq q < \infty$ . Since, for  $p \neq q$ , the spaces  $\ell_{p+}$  and  $\ell_{q+}$  are also not isomorphic (cf. Proposition 3.3), it is rather surprising that  $ces(p+)$  and  $ces(q+)$  *are* isomorphic Fréchet spaces for *all pairs*  $1 \leq p, q < \infty$ . These results are obtained as a consequence of the main result of this paper showing, remarkably, that  $ces(p+)$  coincides with the power series space of order one and finite type  $\Lambda_{-1/p'}(\alpha)$  with  $\alpha := (\log(k))_{k \in \mathbb{N}}$ ; see Theorem 3.1. Accordingly, all these spaces are diagonally isomorphic. We mention two further consequences. The Fréchet spaces  $\ell_{p+}$ , for  $1 \leq p < \infty$ , all *fail* to be (FBa)-spaces, [19], whereas every Fréchet space  $ces(p+)$  *is* an (FBa)-space, since it is a Köthe echelon space of order one; see Proposition 4.1. It is known that  $\ell_{p+}$  has the property that every  $\ell_{p+}$ -valued vector measure has relatively compact range if and only if  $1 \leq p < 2$ . This property also holds for  $ces(p+)$ , but for *every*  $1 \leq p < \infty$ .

## 2. OPTIMAL SOLID LATTICE PROPERTIES OF $ces(p+)$

We begin by noting, for each  $1 \leq p < \infty$ , that  $ces(p+)$  is reflexive, [17, Proposition 25.15], since each Banach space  $ces(q)$ ,  $q > p$ , is reflexive, [6, p.61].

**Lemma 2.1.** *For each  $1 \leq p < \infty$ , the space  $ces(p+)$  is a solid Fréchet lattice subspace of  $\mathbb{C}^{\mathbb{N}}$  and  $\ell_{p+} \subseteq ces(p+)$  with a continuous and proper inclusion.*

*Proof.* Clearly  $ces(p+)$  is a solid Fréchet lattice subspace of  $\mathbb{C}^{\mathbb{N}}$ . Since  $\ell_q \subseteq ces(q)$  with a continuous inclusion for each  $q > p > 1$ , it follows that  $\ell_{p+} \subseteq ces(p+)$  continuously.

Fix  $1 < p < \infty$ . By [8, Remark 2.2(ii)] there exists  $x \in ces(p) \setminus \ell_{\infty}$ . Since  $ces(p) \subseteq ces(p+)$  and  $\ell_{p+} \subseteq \ell_{\infty}$ , it follows that  $x \in ces(p+) \setminus \ell_{p+}$ .

For  $p = 1$  we know that  $\ell_{1+} \subseteq ces(1+)$ . If this containment was an equality, then the open mapping theorem for Fréchet spaces, [17, Theorem 24.30], implies that the identity map from  $\ell_{1+}$  onto  $ces(1+)$  is an isomorphism. This is impossible as  $\ell_{1+}$  is non-Montel whereas  $ces(1+)$  is Montel (see Proposition 3.5(ii) below). So,  $\ell_{1+} \subsetneq ces(1+)$ .  $\square$

The following observation is a direct consequence of the striking property of  $ces(q)$ ,  $1 < p < \infty$ , exhibited in Proposition 1.1 and the definition of  $ces(p+) = \bigcap_{q>p} ces(q)$ .

**Proposition 2.2.** *Let  $1 \leq p < \infty$  and  $x \in \mathbb{C}^{\mathbb{N}}$ . Then*

$$x \in ces(p+) \text{ if and only if } \mathbf{C}(|x|) \in ces(p+). \quad (2.1)$$

We will require the following fact.

**Lemma 2.3.** *For each  $1 \leq p < \infty$ , the Cesàro operator  $\mathbf{C} : ces(p+) \longrightarrow \ell_{p+}$  is continuous.*

*Proof.* Fix  $1 \leq p < \infty$ . If  $x \in ces(p+)$ , then  $|x| \in ces(q)$  for all  $q > p$  and so  $\mathbf{C}(|x|) \in \ell_q$  for all  $q > p$ . This is because  $\mathbf{C} : ces(q) \longrightarrow \ell_q$  is continuous as

$$\|\mathbf{C}(x)\|_q = \|\mathbf{C}(|x|)\|_q \leq \|\mathbf{C}(|x|)\|_q = \|x\|_{ces(q)}, \quad x \in ces(q).$$

Hence,  $\mathbf{C}(|x|) \in \ell_{p+}$ . This shows that  $\mathbf{C}$  maps  $ces(p+)$  into  $\ell_{p+}$ , necessarily continuously by the closed graph theorem for Fréchet spaces, [17, Theorem 24.31].  $\square$

The next result, in combination with Proposition 2.2, shows that  $ces(p+)$ ,  $1 \leq p < \infty$ , exhibits a very desirable property which  $\ell_{p+}$  fails to possess.

**Proposition 2.4.** *For each  $1 \leq p < \infty$ , the Fréchet space  $\ell_{p+}$  fails to have the property (2.1) in Proposition 2.2 (with  $\ell_{p+}$  in place of  $ces(p+)$ ).*

*Proof.* Fix  $1 \leq p < \infty$ . Assume that  $\ell_{p+}$  does have the property (2.1) in Proposition 2.2. By Lemma 2.1 there exists  $x \in ces(p+) \setminus \ell_{p+}$ . Hence, also  $|x| \in ces(p+) \setminus \ell_{p+}$ . Then Lemma 2.3 implies that  $\mathbf{C}(|x|) \in \ell_{p+}$  and hence, by the assumption on  $\ell_{p+}$ , also  $|x| \in \ell_{p+}$ ; contradiction. So,  $\ell_{p+}$  fails the property.  $\square$

The following result should be compared with its Banach lattice counterpart, [8, p.62].

**Proposition 2.5.** *The space  $ces(p+)$ ,  $1 \leq p < \infty$ , is the largest solid Fréchet lattice  $X$  in  $\mathbb{C}^{\mathbb{N}}$  which contains  $\ell_{p+}$  such that  $\mathbf{C}(X) \subseteq \ell_{p+}$ .*

*Proof.* Let  $[\mathbf{C}, \ell_{p+}]_s$  denote the largest solid Fréchet lattice  $X$  in  $\mathbb{C}^{\mathbb{N}}$  which contains  $\ell_{p+}$  such that  $\mathbf{C}(X) \subseteq \ell_{p+}$ . Since  $\mathbf{C}(ces(p+)) \subseteq \ell_{p+}$  (cf. Lemma 2.3), it follows that  $ces(p+)$  itself is a solid Fréchet lattice which contains  $\ell_{p+}$  such that  $\mathbf{C}(ces(p+)) \subseteq \ell_{p+}$ . Accordingly,  $ces(p+) \subseteq [\mathbf{C}, \ell_{p+}]_s$ .

Let  $X$  be any solid Fréchet lattice in  $\mathbb{C}^{\mathbb{N}}$  which contains  $\ell_{p+}$  such that  $\mathbf{C}(X) \subseteq \ell_{p+}$ . Given  $x \in X$  also  $|x| \in X$  and hence,  $\mathbf{C}(|x|) \in \ell_{p+} \subseteq ces(p+)$ . Proposition 2.2 implies that  $x \in ces(p+)$ . Accordingly,  $X \subseteq ces(p+)$ . This implies that  $[\mathbf{C}, \ell_{p+}]_s \subseteq ces(p+)$ .  $\square$

Since  $\ell_{p+} \subseteq ces(p+)$  continuously (cf. Lemma 2.1), in addition to  $\mathbf{C} : \ell_{p+} \longrightarrow \ell_{p+}$  one may also consider the positive Cesàro operator  $\mathbf{C} : ces(p+) \longrightarrow$

$ces(p+)$ . Even though the target space  $ces(p+)$  is now genuinely larger than  $\ell_{p+}$  (cf. Lemma 2.1), no further solid extension occurs!

**Proposition 2.6.** *The space  $ces(p+)$ ,  $1 \leq p < \infty$ , is also the largest solid Fréchet lattice  $X$  in  $\mathbb{C}^{\mathbb{N}}$  which contains  $ces(p+)$  such that  $\mathbf{C}(X) \subseteq ces(p+)$ .*

*Proof.* Denote by  $[\mathbf{C}, ces(p+)]_s$  the largest solid Fréchet lattice  $X$  in  $\mathbb{C}^{\mathbb{N}}$  which contains  $ces(p+)$  such that  $\mathbf{C}(X) \subseteq ces(p+)$ . Clearly  $ces(p+) \subseteq [\mathbf{C}, ces(p+)]_s$  because  $\mathbf{C}(ces(p+)) \subseteq \ell_{p+} \subseteq ces(p+)$ ; see Lemma 2.3.

Let  $X$  be any solid Fréchet lattice in  $\mathbb{C}^{\mathbb{N}}$  which contains  $ces(p+)$  such that  $\mathbf{C}(X) \subseteq ces(p+)$ . Given  $x \in X$  also  $|x| \in X$ . Hence,  $\mathbf{C}(|x|) \in ces(p+)$ . By Proposition 2.2,  $x \in ces(p+)$ . So,  $X \subseteq ces(p+)$ . This implies  $[\mathbf{C}, ces(p+)]_s \subseteq ces(p+)$ .  $\square$

For the Banach lattice counterpart of Proposition 2.6 see [8, Theorem 2.5].

### 3. $ces(p+)$ AS A POWER SERIES SPACE OF FINITE TYPE AND ORDER 1

A power series Fréchet space of *finite type*  $r \in \mathbb{R}$  and *order* 1 is defined, for any given strictly increasing sequence  $\alpha = (\alpha_k)_k \subseteq (0, \infty)$  satisfying  $\lim_{k \rightarrow \infty} \alpha_k = \infty$ , by

$$\Lambda_r(\alpha) := \{x \in \mathbb{C}^{\mathbb{N}} : \|x\|_t := \sum_{k=1}^{\infty} |x_k| e^{t\alpha_k} < \infty, \quad \forall t < r\};$$

see [17, Ch.29], also for the definition of the norms generating the Fréchet topology of  $\Lambda_r(\alpha)$ .

Our main result is rather remarkable and surprising. We require the following inequality

$$\frac{A_p}{k^{1/p'}} \leq \|e_k\|_{ces(p)} \leq \frac{B_p}{k^{1/p'}}, \quad k \in \mathbb{N}, \quad (3.1)$$

valid for strictly positive constants  $A_p, B_p$  and with  $\frac{1}{p} + \frac{1}{p'} = 1$ , [6, Lemma 4.7], where  $e_k := (\delta_{k,n})_n$  for each  $k \in \mathbb{N}$ .

**Theorem 3.1.** *The Fréchet space  $ces(p+)$ ,  $1 \leq p < \infty$ , is isomorphic to the power series space  $\Lambda_{-1/p'}(\alpha)$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ , of finite type  $-1/p'$  and order 1, where  $\alpha = (\log k)_k$ .*

*Proof.* Fix  $1 \leq p < \infty$ . Observe that

$$\Lambda_{-1/p'}((\log k)_k) = \{x \in \mathbb{C}^{\mathbb{N}} : \|x\|_t = \sum_{j=1}^{\infty} |x_j| j^t \text{ for all } t < -1/p'\}.$$

Let  $1 < q < \infty$ . For  $x \in ces(q)$  we have

$$\|x - \sum_{j=1}^m x_j e_j\|_{ces(q)} \leq \left( \sum_{n=m+1}^{\infty} \left( \frac{1}{n} \sum_{k=1}^n |x_k|^q \right)^{1/q} \right) \rightarrow 0, \quad \text{as } m \rightarrow \infty.$$

Therefore  $(e_k)_k$  are a basis of  $ces(q)$  for  $1 < q < \infty$ , [8, Proposition 2.1], hence, also of  $ces(p+)$  for each  $1 \leq p < \infty$ . Consequently, via (3.1) we have, with  $\frac{1}{q} + \frac{1}{q'} = 1$ , that

$$\|x\|_{ces(q)} \leq \sum_{j=1}^{\infty} |x_j| \cdot \|e_j\|_{ces(q)} \leq B_q \sum_{j=1}^{\infty} |x_j| j^{-\frac{1}{q'}}.$$

By [6, Lemma 4.7], for each  $\beta > 0$  we have

$$\frac{1}{\beta} \frac{1}{j^\beta} \leq \sum_{n=j}^{\infty} \frac{1}{n^{1+\beta}}, \quad j \in \mathbb{N}. \quad (3.2)$$

Let  $p < q < \infty$ . Given  $p < s < q$  it is clear that  $y := (n^{-\frac{1}{q'}})_n \in \ell_{s'}$ . So we can apply (3.2) with  $\beta = \frac{1}{q'}$  and Hölder's inequality to get

$$\begin{aligned} q' \sum_{j=1}^{\infty} |x_j| j^{-\frac{1}{q'}} &\leq \sum_{j=1}^{\infty} |x_j| \sum_{n=j}^{\infty} \frac{1}{n} n^{-\frac{1}{q'}} = \\ &= \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{j=1}^n |x_j| \right) n^{-\frac{1}{q'}} \leq \|y\|_{s'} \|x\|_{ces(s)}. \end{aligned}$$

This proves the result.  $\square$

**Corollary 3.2.** *Each of the Fréchet spaces  $ces(p+)$ , for  $1 \leq p < \infty$ , is isomorphic to the power series space  $\Lambda_0(\alpha)$  of finite type 0 and order 1, where  $\alpha = (\log k)_k$ .*

*Proof.* This follows directly from the fact that all finite type power series spaces  $\Lambda_r(\alpha)$ , with  $\alpha$  fixed, are diagonally isomorphic; see [17], page 358, lines 1-5.  $\square$

In view of Corollary 3.2 the following observation is relevant.

**Proposition 3.3.** *For every distinct pair  $1 \leq p, q < \infty$  the Fréchet spaces  $\ell_{p+}$  and  $\ell_{q+}$  are not isomorphic.*

*Proof.* We may assume that  $p < q$ . Suppose that there exists an isomorphism  $T : \ell_{q+} \rightarrow \ell_{p+}$ . Since the natural inclusion  $\ell_q \subseteq \ell_{q+}$  is continuous, the restricted operator  $T|_{\ell_q} : \ell_q \rightarrow \ell_{p+}$  is continuous. Consequently, since the inclusion  $\ell_{p+} \subseteq \ell_r$  is continuous for each  $r > p$ , it follows that  $T|_{\ell_q} : \ell_q \rightarrow \ell_r$  is continuous for each  $r \in (p, q)$ . By Pitt's theorem, [20],  $T : \ell_q \rightarrow \ell_r$  is compact (we denote  $T|_{\ell_q}$  simply by  $T$  again as no confusion can occur). Choose now any  $r \in (p, q)$ . Since  $\{e_j\}_{j=1}^{\infty}$  is a bounded set in  $\ell_q$  and  $T : \ell_q \rightarrow \ell_r$  is compact, the image  $\{T(e_j)\}_{j=1}^{\infty}$  is a relatively compact subset of  $\ell_r$ . Consequently, as  $\ell_{p+} = \bigcap_{p < r < q} \ell_r = \text{proj}_{p < r < q} \ell_r$ , it follows that  $\{T(e_j)\}_{j=1}^{\infty}$  is also a relatively compact subset of  $\ell_{p+}$ . Hence, there exists  $y \in \ell_{p+}$  and a subsequence  $\{T(e_{j(k)})\}_{k=1}^{\infty}$  of  $\{T(e_j)\}_{j=1}^{\infty}$  such

that  $T(e_{j(k)}) \rightarrow y$  in  $\ell_{p+}$  for  $k \rightarrow \infty$ . By continuity of the inverse operator  $T^{-1} : \ell_{p+} \rightarrow \ell_{q+}$  it follows that  $e_{j(k)} \rightarrow T^{-1}(y)$  in  $\ell_{q+}$ . Choose any  $s > q$ , in which case  $\ell_{q+} \subseteq \ell_s$  continuously, then also  $e_{j(k)} \rightarrow T^{-1}(y)$  in the Banach space  $\ell_s$ . This is impossible as  $\|e_{j(k)} - e_{j(l)}\|_{\ell_s} = 2^{1/s} \geq 1$  for all  $k \neq l$ . Hence, no such isomorphism  $T$  of  $\ell_{q+}$  onto  $\ell_{p+}$  can exist.  $\square$

**Remark 3.4.** For each pair  $1 \leq p < q < \infty$  it is clear that

$$ces(p+) \subseteq ces(q+). \quad (3.3)$$

Even though  $ces(p+)$  and  $ces(q+)$  are isomorphic as Fréchet spaces (cf. Corollary 3.2), the containment (3.3) is *proper*. Indeed, if it were an equality, then for any fixed  $r \in (p, q)$  it would follow from the (continuous) inclusions  $ces(p+) \subseteq ces(r) \subseteq ces(q) \subseteq ces(q+)$  that  $ces(r) = ces(q)$ . Consequently, the Banach spaces  $ces(r)$  and  $ces(q)$  would be isomorphic (with  $r < q$ ) which is not the case, [4, Proposition 3.3].

We now collect some further consequences of Theorem 3.1.

**Proposition 3.5.** *For each  $1 \leq p < \infty$  the following assertions hold.*

- (i) *The Fréchet space  $ces(p+)$  is a Köthe echelon space of order 1 and the canonical vectors  $(e_j)_{j \in \mathbb{N}}$  form an unconditional basis of  $ces(p+)$ .*
- (ii)  *$ces(p+)$  is a Fréchet-Schwartz space but, it is not nuclear.*
- (iii)  *$ces(p+)$  is not isomorphic to  $\ell_{q+}$  for each  $1 \leq q < \infty$ .*

*Proof.* (i) The space  $ces(p+)$  is a Köthe echelon space of order 1 (by Theorem 3.1). The canonical vectors are an unconditional basis for every Köthe echelon space of order 1. Even stronger, they form an absolute basis, [16, pp.314–315].

(ii) Every power series space is Schwartz by [17, Proposition 27.10]. The non-nuclearity of  $ces(p+) = \Lambda_{-1/p'}(\alpha)$ ,  $\alpha = (\log k)_k$  is a direct consequence of Corollary 3.2 and [17, Proposition 29.6 (2)].

(iii) The space  $\ell_{q+}$  is not Montel for each  $1 \leq q < \infty$ , [18], and hence, it cannot be isomorphic to  $ces(p+)$ , [17, Lemma 24.19].  $\square$

**Definition 3.6.** Let  $X$  be a Fréchet space whose topology is generated by a fundamental sequence  $(\|\cdot\|_n)_{n \in \mathbb{N}}$  of seminorms. The space  $X$  has the property  $(\overline{\Omega})$  if:  $\forall l \in \mathbb{N}$ ,  $c \in (0, 1) \exists k > l, \forall h > k, \exists C > 0$  :

$$\|y\|_k^{1+c} \leq C \|y\|_h' \|y\|_l^c, \quad \forall y \in X',$$

where  $\|y\|_k' := \sup\{|\langle x, y \rangle| : \|x\|_k \leq 1\}$  is the dual norm  $\|\cdot\|_k'$  of  $\|\cdot\|_k$  in  $X'$ .

It is a consequence of [17, Lemma 29.16] that the condition in Definition 3.6 coincides with property  $(\overline{\Omega})$ .

Every space  $\ell_{p+}$ , for  $1 \leq p < \infty$ , satisfies property  $(\overline{\Omega})$ , [18, proof of Proposition 2.4]. Since all power series spaces of finite type have property  $(\overline{\Omega})$ , [17, Proposition 29.12], we deduce the following result.

**Proposition 3.7.** *For each  $1 \leq p < \infty$  the Fréchet space  $ces(p+)$  has property  $(\overline{\Omega})$ .*



4. FURTHER PROPERTIES OF  $ces(p+)$ 

According to Taskinen, [23], a Fréchet space  $X$  is called an (FBa)-space if, for every Banach space  $Y$ , every bounded subset of the complete projective tensor product  $X\widehat{\otimes}Y$  is contained in the closed convex hull  $\overline{\text{co}}(C \otimes D)$  of bounded sets  $C \subseteq X$  and  $D \subseteq Y$ . In 1986 Taskinen constructed Fréchet spaces which are not (FBa)-spaces, thereby solving the “problem of topologies of Grothendieck”, [22]. Peris proved that the Fréchet spaces  $\ell_{p+}$ , for  $1 \leq p < \infty$ , are *not* (FBa) spaces, thus providing a natural and concrete class of spaces of this type, [19].

**Proposition 4.1.** *Each Fréchet space  $ces(p+)$ , for  $1 \leq p < \infty$ , is an (FBa)-space.*

*Proof.* Each space  $ces(p+)$ ,  $1 \leq p < \infty$ , is isomorphic to a Köthe echelon space of order 1 (cf. Proposition 3.5 (i)). Hence, it is necessarily an (FBa)-space, [14, p.70], [22, Section 3].  $\square$

Now we consider some features of  $ces(p+)$  of a somewhat different nature. First,  $ces(p+)$  is the complexification of the corresponding real Riesz space  $ces_{\mathbb{R}}(p+) := \{x \in ces(p+) : x = (x_n)_n \in \mathbb{R}^{\mathbb{N}}\}$ , in the sense of [26, pp.187–201]. Since  $ces_{\mathbb{R}}(p+)$  is solid in  $\mathbb{R}^{\mathbb{N}}$  it follows that  $ces_{\mathbb{R}}(p+)$  is *Dedekind complete* (i.e., every subset of  $ces_{\mathbb{R}}(p+)$  which is bounded from above in the order has a least upper bound) and hence, (per definition) also its complexification  $ces(p+)$  is Dedekind complete. Moreover, being reflexive, each of the (separable) Fréchet lattices  $ces(p+)$ ,  $1 \leq p < \infty$ , has a *Lebesgue topology*, [2, Theorems 10.3 and 10.9], that is, if  $x^{(\alpha)} \downarrow 0$  is a decreasing net in the order of  $ces(p+)$ , then  $\lim_{\alpha} x^{(\alpha)} = 0$  in the topology of  $ces(p+)$ .

For every  $1 \leq p < \infty$ , the Fréchet space  $\ell_{p+}$  has the property that *every*  $\ell_{p+}$ -valued vector measure (always assumed to be countably additive and defined on a  $\sigma$ -algebra) has relatively compact range if and only if  $p \in [1, 2)$ , [7, Proposition 2.8]. Once again the optimal solid extension  $ces(p+)$  of  $\ell_{p+}$  exhibits better behaviour in this regard.

**Proposition 4.2.** *Let  $p \in [1, \infty)$ . Then every  $ces(p+)$ -valued vector measure necessarily has relatively compact range.*

*Proof.* The range of every  $ces(p+)$ -valued vector measure is a relatively weakly compact set, [24], and hence, is also relatively compact as  $ces(p+)$  is a Fréchet-Montel space (by Proposition 3.5(ii)).  $\square$

Since  $ces(p+)$ ,  $1 \leq p < \infty$ , is *not* nuclear (cf. Proposition 3.5(ii)), there exist  $ces(p+)$ -valued vector measures which *fail* to have finite variation, [11, Corollary and Theorem 2].

A Fréchet space  $X$  is said to have the *Rybakov property*, [12], if for every  $X$ -valued vector measure  $\nu$  there exists  $x' \in X'$  such that  $\nu \ll |\langle \nu, x' \rangle|$  (i.e.,  $\nu(F) = 0$  for every measurable set  $F \subseteq E$  whenever  $|\langle \nu, x' \rangle|(E) = 0$ ). Here  $\langle \nu, x' \rangle$  is the complex measure  $E \mapsto \langle \nu(E), x' \rangle$  and  $|\langle \nu, x' \rangle|$  denotes its total

variation measure. A Fréchet space  $X$  has Rybakov's property if and only if it admits a continuous norm; see [12, Theorem 2.2] for real spaces and [21, Proposition 2.2] for complex spaces. Since both  $\ell_{p+}$  and  $ces(p+)$  admit a continuous norm, we have the following fact.

**Proposition 4.3.** *For each  $1 \leq p < \infty$  the Fréchet spaces  $\ell_{p+}$  and  $ces(p+)$  have the Rybakov property.*

A classical result of Bade, [5, Theorem 3.1], states: Given a  $\sigma$ -complete Boolean algebra of projections  $\mathcal{M}$  in a Banach space  $X$ , for each  $x_0 \in X$  there exists  $x' \in X'$  (called a Bade functional for  $x_0$  with respect to  $\mathcal{M}$ ) satisfying

- (i)  $\langle P(x_0), x' \rangle \geq 0$  for all  $P \in \mathcal{M}$ , and
- (ii) if  $\langle P(x_0), x' \rangle = 0$  for some  $P \in \mathcal{M}$ , then  $P(x_0) = 0$ .

A Fréchet space  $X$  is said to have the *Bade property*, [21], if every  $\sigma$ -complete Boolean algebra of projections  $\mathcal{M}$  in  $X$  satisfies (i), (ii) above (for every  $x_0 \in X$ ). This is the case if and only if  $X$  admits a continuous norm, [21, Corollary 2.1], which yields the following result.

**Proposition 4.4.** *For each  $1 \leq p < \infty$  the Fréchet spaces  $\ell_{p+}$  and  $ces(p+)$  have the Bade property.*

Our final result presents a description of the dual of  $ces(p+)$ .

Recall that the *Köthe dual*  $X^\times$  (or the associate space) of a Banach sequence space  $(X, \|\cdot\|_X)$ , with  $\varphi \subseteq X \subseteq \mathbb{C}^{\mathbb{N}}$ , is defined by

$$X^\times := \left\{ x \in \mathbb{C}^{\mathbb{N}} : \sum_{k=1}^{\infty} |x_k y_k| < \infty, \forall y \in X \right\},$$

endowed with the norm

$$\|x\|_{X^\times} := \sup \left\{ \sum_{k=1}^{\infty} |x_k y_k| : \|y\|_X \leq 1 \right\},$$

[17, Ch.27]. Here  $\varphi$  is the subspace of  $\mathbb{C}^{\mathbb{N}}$  consisting of those vectors having finite support. Every  $v \in X^\times$  defines an element of the dual Banach space  $X'$  of  $(X, \|\cdot\|_X)$  via  $u \rightarrow \sum_{n=1}^{\infty} u_n v_n$  for  $u \in X$ , and  $\|v\|_{X^\times} = \|v\|_{X'}$ .

For  $1 < q < \infty$  and  $\frac{1}{q} + \frac{1}{q'} = 1$ , the space  $d(q')$  is defined as

$$d(q') := \left\{ x \in \mathbb{C}^{\mathbb{N}} : \sum_{n=1}^{\infty} \sup_{k \geq n} (|x_k|^{q'}) < \infty \right\},$$

which is a Banach space when endowed with the norm

$$\|x\|_{d(q')} := \left( \sum_{n=1}^{\infty} \sup_{k \geq n} (|x_k|^{q'}) \right)^{1/q'}, \quad x \in d(q').$$

Observe that

$$x \in d(q') \text{ if and only if } \widehat{x} := (\sup_{k \geq n} |x_k|)_{n \in \mathbb{N}} \in \ell_{q'} \quad (4.1)$$

and that

$$\|x\|_{d(q')} = \|\widehat{x}\|_{\ell_{q'}};$$

see [6, p.3 & p.9]. We will require the following result of Bennett [6, p.61 & Corollary 12.17].

**Lemma 4.5.** *Let  $1 < q < \infty$  and  $\frac{1}{q} + \frac{1}{q'} = 1$ . The map  $\Phi_q: (ces(q))' \rightarrow d(q')$  defined by  $\Phi_q(f) := (\langle e_j, f \rangle)_{j \in \mathbb{N}}$ , for  $f \in (ces(q))'$ , is a linear isomorphism of the dual Banach space  $(ces(q))'$  onto the Banach space  $d(q')$  and*

$$\frac{1}{q'} \|\Phi_q(f)\|_{d(q')} \leq \|f\|_{(ces(q))'} \leq (q-1)^{1/q} \|\Phi_q(f)\|_{d(q')}, \quad f \in (ces(q))'.$$

Moreover,  $(ces(q))^\times = d(q')$  and  $(d(q'))^\times = ces(q)$ , with equivalent norms.

Fix  $1 \leq p < \infty$  and any sequence  $p < p_{n+1} < p_n$  for  $n \in \mathbb{N}$  satisfying  $\lim_{n \rightarrow \infty} p_n = p$ . Then  $p'_n < p'_{n+1} < p'$  for  $n \in \mathbb{N}$ . Since  $\|x\|_{d(r)} = \|\widehat{x}\|_{\ell_r}$  for all  $x \in d(r)$  and any  $1 < r < \infty$ , it follows that  $d(p'_n) \subseteq d(p'_{n+1})$  with a continuous inclusion for each  $n \in \mathbb{N}$ . We endow the vector space  $d(p'-) := \cup_{n \in \mathbb{N}} d(p'_n)$ , which is an increasing union, with the inductive limit topology, i.e.,  $d(p'-) := \text{ind}_n d(p'_n)$ , [17, Ch.24].

**Proposition 4.6.** *Let  $1 \leq p < \infty$ . The map  $\Lambda: (ces(p+))' \rightarrow d(p'-)$  given by  $\Lambda(f) := (\langle e_j, f \rangle)_{j \in \mathbb{N}}$ , for  $f \in (ces(p+))'$ , defines a linear bijection which is a topological isomorphism of the strong dual space  $(ces(p+))'_\beta$  onto  $d(p'-) = \text{ind}_n d(p'_n)$ .*

*Proof.* First observe that  $\varphi$  is dense in the Fréchet space  $ces(p+)$  as it is dense in each Banach space  $ces(q)$  for  $q > p$ .

Fix  $u \in (ces(p+))'$ . Select  $n \in \mathbb{N}$  and a constant  $K > 0$  such that

$$|\langle x, u \rangle| \leq K \|x\|_{ces(p_n)}, \quad x \in ces(p+).$$

So, there exists a unique  $\tilde{u} \in (ces(p_n))'$  whose restriction to  $ces(p+) \subseteq ces(p_n)$  coincides with  $u$ . By Lemma 4.5 the element  $(\langle e_j, u \rangle)_{j \in \mathbb{N}} = (\langle e_j, \tilde{u} \rangle)_{j \in \mathbb{N}}$  belongs to  $d(p'_n) \subseteq d(p'-)$ .

The previous argument implies that  $\Lambda$  is well defined. It is clearly linear. Moreover,  $\Lambda$  is injective by the density of  $\varphi = \text{span}\{e_j: j \in \mathbb{N}\}$  in  $ces(p+)$ . To show that  $\Lambda$  is also surjective let  $y = (y_j)_{j \in \mathbb{N}} \in d(p'-)$ . Then there exists  $m \in \mathbb{N}$  such that  $y \in d(p'_m)$ . Via Lemma 4.5 we can find  $f \in (ces(p_m))'$  with  $y = (\langle e_j, f \rangle)_{j \in \mathbb{N}}$ . Then the restriction  $v$  of  $f$  to  $ces(p+)$  belongs to  $(ces(p+))'$  and  $\Lambda(v) = y$ .

Define the injection  $J_n: (ces(p_n))' \rightarrow (ces(p+))'$ , for  $n \in \mathbb{N}$ , by setting  $J_n(f)$  to be the restriction of  $f \in (ces(p_n))'$  to  $ces(p+)$ . By the earlier part of the proof  $(ces(p+))' = \cup_{n \in \mathbb{N}} J_n((ces(p_n))')$  and so we may consider in  $(ces(p+))'$  the inductive limit topology  $\text{ind}_n (ces(p_n))'$ . Since  $ces(p+)$  is reflexive, its strong dual  $(ces(p+))'_\beta$  coincides with  $\text{ind}_n (ces(p_n))'$ .

By Lemma 4.5, for each  $n \in \mathbb{N}$  the restriction  $\Lambda_{p_n}$  of  $\Lambda$  to  $(ces(p_n))'$  is continuous from  $(ces(p_n))'$  onto  $d(p'_n)$ . This implies that  $\Lambda : (ces(p+))'_\beta \rightarrow \text{ind}_n d(p'_n)$  is a continuous bijection. By the closed graph theorem for (LB)-spaces, [17, Theorem 24.31 & Remark 24.36],  $\Lambda$  is also a topological isomorphism.  $\square$

**Acknowledgements.** The authors thank the referee for a considerable simplification of some proofs in Section 3.

The research of the first two authors was partially supported by the project MTM2016-76647-P (Spain). The second author thanks the Mathematics Department of the Katholische Universität Eichstätt-Ingolstadt (Germany) for its support and hospitality during his research visit in the period September 2016 - July 2017.

#### REFERENCES

- [1] T. Abdeljawad, M. Yurdakul, *The property of smallness up to a complemented Banach subspace*, Publ. Math. Debrecen **64** (2004), no. 3–4, 415–425.
- [2] C.D. Aliprantis, O. Burkinshaw, *Locally Solid Riesz Spaces*, Academic Press, New York San Francisco, 1978.
- [3] A.A. Albanese, J. Bonet, W.J. Ricker, *The Cesàro operator in the Fréchet spaces  $\ell^{p+}$  and  $L^{p-}$* , Glasgow Math. J. **59** (2017), 273–287.
- [4] A.A. Albanese, J. Bonet, W.J. Ricker, *Multiplier and averaging operators in the Banach spaces  $ces(p)$ ,  $1 < p < \infty$* , Preprint 2017.
- [5] W.G. Bade, *On Boolean algebras of projections and algebras of operators*, Trans. Amer. Math. Soc. **80** (1955), 345–360.
- [6] G. Bennett, *Factorizing the classical inequalities*, Mem. Amer. Math. Soc. **120** (1996), no 576, viii + 130 pp.
- [7] J. Bonet, W.J. Ricker, *The canonical spectral measure in Köthe echelon spaces*, Integral Equations Operator Theory, **53** (2005), 477–496.
- [8] G.P. Curbera, W.J. Ricker, *Solid extensions of the Cesàro operator on  $\ell^p$  and  $c_0$* , Integral Equations Operator Theory **80** (2014), 61–77.
- [9] J.-C. Díaz, *An example of a Fréchet space, not Montel, without infinite-dimensional normable subspaces*, Proc. Amer. Math. Soc. **96** (1986), 721.
- [10] M.M. Dragilev, P.A. Chalov, *On Fréchet spaces with an unconditional basis* (Russian. Russian summary), Mat. Zametki **80** (2006), no. 1, 29–32; translation in Math. Notes **80** (2006), no. 1-2, 27–30.
- [11] M. Duchoň, *Vector measures and nuclearity*, Math. Slovaca, **38** (1988), 79–83.
- [12] A. Fernández, F. Naranjo, *Rybakov's theorem for vector measures in Fréchet spaces*, Indag. Math. (N.S.), **8** (1997), 33–42.
- [13] K.-G. Grosse-Erdmann, *The Blocking Technique, Weighted Mean Operators and Hardy's Inequality*, Lecture Notes in Mathematics, **1679**, Springer Verlag, Berlin Heidelberg, 1998.
- [14] A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955).
- [15] G.H. Hardy, J.E. Littlewood, G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, reprinted, 1964.
- [16] H. Jarchow, *Locally Convex Spaces*, Teubner, Stuttgart, 1981.

- [17] R. Meise, D. Vogt, *Introduction to Functional Analysis*, Clarendon Press, Oxford, 1997.
- [18] G. Metafune, V.B. Moscatelli, *On the space  $\ell^{p+} = \cap_{q>p} \ell^q$* , Math. Nachr. **147** (1990), 7–12.
- [19] A. Peris, *Quasinormable spaces and the problem of topologies of Grothendieck*, Ann. Acad. Sci. Fenn. Ser. A I Math. **19** (1994), 167–203.
- [20] H.R. Pitt, *A note on bilinear forms*, J. London Math. Soc. **11** (1936), 171–174.
- [21] W.J. Ricker, *Existence of Bade functionals for complete Boolean algebras of projections in Fréchet spaces*, Proc. Amer. Math. Soc. **125** (1997), 2401–2407.
- [22] J. Taskinen, *Counterexamples to “Problème des topologies” of Grothendieck*, Ann. Acad. Sci. Fenn. Ser. A I Math. Diss. **63** (1986).
- [23] J. Taskinen, *(FBa) and (FBB)-spaces*, Math. Z. **198** (1988), 339–365.
- [24] I. Tweddle, *Weak compactness in locally convex spaces*, Glasgow Math. J. **9** (1968), 123–127.
- [25] D. Vogt, *Some results on continuous linear maps between Fréchet spaces*, Functional Analysis: Survey and Recent Results (eds. K.D. Bierstedt and B. Fuchssteiner), North-Holland Math. Stud. **90** (1984), 349–381.
- [26] A.C. Zaanen, *Riesz Spaces II*, North Holland, Amsterdam, 1983.

ANGELA A. ALBANESE, DIPARTIMENTO DI MATEMATICA E FISICA “E. DE GIORGI”,  
UNIVERSITÀ DEL SALENTO- C.P.193, I-73100 LECCE, ITALY  
*E-mail address:* `angela.albanese@unisalento.it`

JOSÉ BONET, INSTITUTO UNIVERSITARIO DE MATEMÁTICA PURA Y APLICADA  
IUMPA, UNIVERSITAT POLITÈCNICA DE VALÈNCIA, E-46071 VALENCIA, SPAIN  
*E-mail address:* `jbonet@mat.upv.es`

WERNER J. RICKER, MATH.-GEOGR. FAKULTÄT, KATHOLISCHE UNIVERSITÄT EICHSTÄTT-  
INGOLSTADT, D-85072 EICHSTÄTT, GERMANY  
*E-mail address:* `werner.ricker@ku-eichstaett.de`