

ÍNDICE DE CONTENIDOS

1. INTRODUCCIÓN GENERAL	1
1.1. Tecnologías embrionarias aplicadas en bovinos	2
1.2. Producción de embriones bovinos <i>in vitro</i>	7
1.2.1. Maduración <i>in vitro</i>	7
1.2.2. Fecundación <i>in vitro</i>	9
1.2.3. Cultivo <i>in vitro</i>	13
1.2.4. Características de los medios de cultivo <i>in vitro</i> para bovinos	16
1.2.5. Factores de crecimiento para el cultivo <i>in vitro</i> de embriones bovinos	20
1.2.6. Cocultivo de células endometriales y embriones	21
1.3. Criopreservación de embriones producidos <i>in vitro</i>	23
1.3.1. Congelación lenta	24
1.3.2. Vitrificación	25
1.4. Evaluación de la calidad y viabilidad de embriones bovinos producidos <i>in vitro</i>	27
1.4.1. Morfología	28
1.4.2. Recuento total y diferencial de células embrionarias	30
1.4.3. Apoptosis y daño celular	32
1.4.4. Contenido de lípidos intracelulares	33
1.4.5. Prueba de supervivencia a la criopreservación	35
1.4.6. Expresión génica	36
1.4.7. Dimorfismo sexual	37
1.5. La viabilidad del embrión en las receptoras	41
1.5.1. Establecimiento de la gestación	41
1.5.2. Pérdidas embrionarias tempranas y aborto	43

1.5.3. Resultados perinatales con embriones producidos <i>in vitro</i>	45
1.5.4. Salud perinatal de terneros producidos <i>in vitro</i>	47
1.6. Referencias	49
2. OBJETIVOS	70
3. CAPÍTULO I	
Short- and long-term outcomes of the absence of protein during bovine blastocyst formation <i>in vitro</i>	73
3.1. Introduction	75
3.2. Material and Methods	77
3.2.1. <i>In vitro</i> embryo production	77
3.2.2. Differential cell counts	79
3.2.3. Vitrification of embryos	79
3.2.4. Warming and survival of vitrified embryos	80
3.2.5. Oestrus synchronization of recipients, embryo transfer and pregnancy and birth monitoring	80
3.2.6. Gene expression analysis: RNA extraction, reverse transcription and quantification of mRNA transcript abundance	80
3.2.7. Embryo sexing by PCR	83
3.2.8. Statistical analysis	83
3.3. Results	84
3.3.1. Embryo development	84
3.3.2. Differential cell counts	85
3.3.3. <i>In vitro</i> survival of vitrification and warming	85
3.3.4. Embryo transfer to recipients and birth	86
3.3.5. Gene expression study	87

3.3.6. Sex analysis of Day-7 blastocysts	89
3.4. Discussion	89
3.5. References	96
4. CAPÍTULO II	
Protein in culture and endogenous lipid interact with embryonic stages <i>in vitro</i> to alter calf birthweight after embryo vitrification and warming	103
4.1. Introduction	105
4.2. Material and Methods	108
4.2.1. Embryo production	108
4.2.2. Embryo sexing by PCR	110
4.2.3. Vitrification of embryos for transfer	110
4.2.4. Lipid and cell nuclei staining	111
4.2.5. RNA analysis	112
4.2.6. Embryo transfer, pregnancy diagnosis and calf phenotyping	114
4.2.7. Experimental design	115
4.2.7.1. Experiment 1: Day-7 embryo development patterns and sex ratio after individual culture in the presence or absence of protein during 24h	115
4.2.7.2. Experiment 2: Analysis of gene expression after individual culture in the presence or absence of protein	115
4.2.7.3. Experiment 3: Long-term effects of protein removal and development patterns	116
4.2.8. Statistical analysis	116

4.3. Results	117
4.3.1. Embryo development in individual culture	117
4.3.2. Embryo sex ratio	119
4.3.3. Lipid content study	119
4.3.4. Gene expression study	122
4.3.5. Embryo transfer, pregnancy, birth and calf phenotypes	125
4.4. Discussion	128
4.5. References	134
5. CAPÍTULO III	
Low serum concentration in bovine embryo culture enhances early blastocyst rates on Day-6 with quality traits in the expanded blastocyst stage similar to BSA-cultured embryos	141
5.1. Introduction	143
5.2. Material and Methods	145
5.2.1. Oocyte collection and <i>in vitro</i> maturation (IVM)	145
5.2.2. Sperm preparation and <i>in vitro</i> fertilization (IVF)	146
5.2.3. <i>In vitro</i> culture (IVC)	146
5.2.4. Embryo vitrification, warming and <i>in vitro</i> survival	147
5.2.5. Apoptosis in blastocysts: TUNEL assay	148
5.2.6. Lipids and cell nuclei staining and quantification	149
5.2.7. Estrus synchronization of recipients, embryo transfer and pregnancy diagnosis	150
5.2.8. Experimental design	151
5.2.8.1. Experiment 1: FCS (0.1%) supplementation and embryo development	151
5.2.8.2. Experiment 2: FCS (0.1%) supplementation	

and <i>in vitro</i> survival to cryopreservation	152
5.2.8.3. Experiment 3: FCS (0.1%) supplementation and apoptosis rates of vitrified/warmed Day-7 expanded blastocysts	152
5.2.8.4. Experiment 4: FCS (0.1%) supplementation and lipid contents of Day-6 morulae and Day-7 blastocyst expanded	152
5.2.9. Statistical analysis	153
5.3. Results	153
5.3.1. Embryo development	153
5.3.2. <i>In vitro</i> survival to vitrification and warming	154
5.3.3. Apoptosis and total cell numbers	155
5.3.4. Lipid quantification study	157
5.3.5. Embryo transfer and pregnancies	159
5.4. Discussion	159
5.5. References	165
6. CAPÍTULO IV	
Hepatoma-derived growth factor: Protein quantification in uterine fluid, gene expression in endometrial-cell culture and effects on <i>in vitro</i> embryo development, pregnancy and birth	170
6.1. Introduction	172
6.2. Material and Methods	174
6.2.1. Embryo production	174
6.2.2. Vitrification of embryos for transfer	176
6.2.3. Animals and embryo transfer (ET) for uterine fluid collection	176
6.2.4. Animals and embryo transfer for pregnancy and birth	177

6.2.5. Multiple reaction monitoring (MRM) for HDGF contents in uterine fluid	178
6.2.6. Identification of HDGF in BSA supplements by western blot	179
6.2.7. Endometrial cell and embryo co-culture	180
6.2.8. Gene expression of HDGF in epithelial and stromal endometrial cell culture	181
6.2.9. Experimental design	183
6.2.9.1. Experiment 1: development of individually cultured morulae and early blastocysts in the presence of rHDGF	183
6.2.9.2. Experiment 2: targeted HDGF protein quantification in UF and BSA supplement	184
6.2.9.3. Experiment 3: influence of embryonic sex on gene expression of HDGF in cultured endometrial cells	184
6.2.9.4. Experiment 4: pregnancy rates and birth measurements after transfer of embryos produced with rHDGF	184
6.2.10. Statistical analysis	185
6.3. Results	186
6.3.1. Embryo development	186
6.3.2. HDGF contents in uterine fluid	186
6.3.3. HDGF identification in BSA extracts	188
6.3.4. Pregnancy and birth rates, and calf measurements, of embryos cultured with rHDGF	188
6.3.5. Gene expression of HDGF in endometrial cell cultures	189

6.4. Discussion	190
6.5. References	195
7. DISCUSIÓN GENERAL	200
7.1. Restricción de proteína y desarrollo de embriones	202
7.2. Morfología y recuento diferencial de células del blastocisto después de la restricción de proteína	205
7.3. Restricción de proteína y supervivencia a la criopreservación	206
7.4. Contenido de lípidos intracelulares después de la restricción de proteína	207
7.5. Expresión génica tras la restricción de proteína	208
7.6. Restricción de proteína y dimorfismo sexual	210
7.7. Gestación y resultados perinatales	212
7.8. Referencias	216
8. CONCLUSIONES	221

ÍNDICE DE FIGURAS

INTRODUCCIÓN GENERAL

Figura 1.1.	Producción mundial de embriones bovinos <i>in vivo</i> (MOET) e <i>in vitro</i> entre el año 1997 y 2016.	4
Figura 1.2.	Clasificación de ovocitos bovinos recuperados de ovarios antes del proceso de maduración <i>in vitro</i> .	7
Figura 1.3.	Esquema de los procesos intracelulares inducidos por la heparina durante la capacitación de espermatozoides bovinos.	11
Figura 1.4.	Desarrollo embrionario bovino y activación del genoma embrionario	15
Figura 1.5.	Esquema de enfriamiento y zonas críticas de daño celular durante la criopreservación.	27
Figura 1.6.	Diferentes estadios de embriones bovinos producidos <i>in vitro</i> e <i>in vivo</i>	29
Figura 1.7.	Acumulación de lípidos intracelulares en un blastocisto expandido bovino producido <i>in vitro</i> y representación esquemática de la composición de las gotas de lípidos	34

CAPÍTULO I

Figure 3.1.	Relative abundance of mRNA transcribed from genes representing metabolism, oxidative stress, imprinting mechanisms and endoplasmic reticulum stress performed on pools of Day-7 fresh expanded blastocysts and on pools of hatched blastocysts that survived 24 h <i>in vitro</i> after vitrification and warming of Day-7 expanded blastocysts derived from Day-6 embryos cultured individually in 12 μ L of synthetic oviduct fluid medium with protein or without protein	88
--------------------	--	-----------

CAPÍTULO II

Figure 4.1.	Lipid content per embryonic cell in Day-6 morulae and Day-7 expanded blastocysts cultured with or without protein from Day 6 to Day 7	120
Figure 4.2.	Lipid granules per embryonic cell counted in Day-7 expanded bovine blastocysts following single culture of Day-6 morulae or early blastocysts in SOF with protein or without protein	121
Figure 4.3.	Average cell abundance of lipid droplets <2 μ m in diameter in Day-7 expanded bovine blastocysts following single culture of Day-6 morulae or early blastocysts in SOF with	121

protein or without protein

- Figure 4.4.** Representative three-dimensional reconstructions of confocal datasets showing lipid droplets stained with Nile red in a Day-7 bovine expanded blastocyst **122**
- Figure 4.5.** Real-time PCR quantification of relative abundance of transcripts from genes involved in lipid fatty-acid metabolism in Day-7 fresh expanded blastocysts derived from Day-6 morulae cultured for 24 h with or without protein **124**
- Figure 4.6.** Real-time PCR quantification of relative abundance of transcripts from genes involved in metabolism, lipid oxidative metabolism, antioxidative and stress defence and apoptosis in Day-7 fresh expanded blastocysts derived from Day-6 morulae cultured for 24 h with or without protein **124**
- Figure 4.7.** Birthweight on a per bull basis of calves born after transfer of Day-7 *IVP* vitrified–warmed embryos whose development stages on Day 6 were monitored as compact morulae or early blastocysts **127**

CAPÍTULO III

- Figure 5.1.** Experimental design **151**
- Figure 5.2.** **(a)** Effects of serum (FCS) in culture up to Day-6 and developmental stage on Day-6, on total cell numbers and apoptosis percentages within hatched blastocysts surviving in culture after vitrification/warming of Day-7 expanded blastocysts. **(b)** Representative photomicrographs of single Z-axis confocal section of hatched blastocysts after vitrification/warming of expanded blastocysts produced in mSOF + 6 mg mL⁻¹ BSA with FCS up to Day-6 or without FCS up to Day-6, and after one-day single culture without protein from Day-6 morula or from Day-6 early blastocyst **156**
- Figure 5.3.** **(a)** Lipid droplets number and sizes per cell counted in fresh expanded blastocysts from culture with FCS up to Day-6 or without FCS up to Day-6, from Day-6 morula or from Day-6 early blastocyst **(b)** Representative three-dimensional reconstructions (frontal view) of confocal datasets showing lipid droplets stained with Nile Red in fresh expanded blastocysts produced in mSOF + 6 mg mL⁻¹ BSA with FCS up to Day-6 or without FCS up to Day-6, and after one-day single culture without protein from Day-6 morula or from Day-6 early blastocyst **158**

CAPÍTULO IV

Figure 6.1. Western blot analysis of HGDF in bovine cold-precipitated serum albumin and bovine fatty-acid-free serum albumin **188**

Figure 6.2. Quantitative real-time PCR analysis of HDGF relative expression in epithelial cells cultured for 48 h with one single male embryo, one single female embryo or no embryos **190**

ÍNDICE DE TABLAS

CAPÍTULO I

Table 3.1.	Details of primers used for qPCR	83
Table 3.2.	<i>In vitro</i> development of Day-6 embryos produced in group culture with SOFaaci with BSA and subsequently cultured individually in 12- μ L drops of SOFaaci with protein or without protein up to Day 8	85
Table 3.3.	Differential cell counts in Day-8 expanded blastocysts cultured individually from Day 6 to Day 8 in drops of SOFaaci with protein or without protein	85
Table 3.4.	<i>In vitro</i> survival after vitrification and warming of Day-7 and Day-8 bovine expanded blastocysts that were cultured individually in synthetic oviduct fluid medium with protein or without protein from Day 6 onwards	86
Table 3.5.	Bodyweight, calves with bodyweight larger than 50 kg, height at withers, thorax circumference, gestation length and average fetal daily weight gain of <i>IVP</i> calves born after transfer of bovine vitrified-warmed Day-7 expanded blastocysts derived from Day-6 embryos cultured individually in 12 μ L of synthetic oviduct fluid medium with protein or without protein	87
Table 3.6.	Embryo transfers performed per bull, with expression of Day-40 pregnancies and miscarriage attained	87

CAPÍTULO II

Table 4.1.	Oligonucleotides used for real-time PCR gene expression 1 analysis	113
Table 4.2.	<i>In vitro</i> blastocyst development rates of Day-6 morulae and early blastocysts cultured individually in 12 μ L drops of SOF containing amino-acids, citrate and myo-inositol with (+; 6mg mL ⁻¹ BSA) or without (-; 0.5 mg mL ⁻¹ PVA replacing BSA) protein up to Day 8	118
Table 4.3.	Expanded blastocyst sex ratio (percent males) according to their Day-6 origin from the morula or the early blastocyst stages and subsequent individual culture in 12 μ L drops of SOF with (+; 6 mg mL ⁻¹ BSA) or without (-; 0.5 mg mL ⁻¹ PVA replacing BSA) protein	119
Table 4.4.	Pregnancy, calving and miscarriage rates after transfer of bovine vitrified Day-7 expanded blastocysts in terms of their Day-6 origin from the morula or the early blastocyst stage	125

and subsequent individual culture in 12- μ L drops of SOF with (+; 6 mg mL⁻¹ BSA) or without (-; 0.5 mg mL⁻¹ PVA replacing BSA) protein

- Table 4.5.** Progesterone increases from Day-0 up to Day-7 in recipients transferred with Day-7 expanded blastocysts in terms of their Day-6 origin from the morula or the early blastocyst stage and their pregnancy status on Day-40 **125**
- Table 4.6.** Progesterone increases from Day-0 up to Day-7 in recipients transferred with Day-7 expanded blastocysts in terms their culture with or without protein from Day-6 onwards and their pregnancy status on Day-40 **125**
- Table 4.7.** Weight and morphometry of calves born after transfer of bovine vitrified–warmed Day-7 expanded blastocysts in terms of their Day-6 origin from the morula or the early blastocyst stage **126**
- Table 4.8.** Weight and morphometry of female calves born after transfer of bovine vitrified–warmed Day-7 expanded blastocysts according to their Day-6 origin from the morula or the early blastocyst stage **126**
- Table 4.9.** Weight and morphometry of calves born after transfer of bovine fresh vs. vitrified/warmed Day-7 expanded blastocysts in terms of their Day-6 origin from the morula or the early blastocyst stage and subsequent individual culture in 12- μ L of SOF containing protein (6 mg mL⁻¹ BSA) **128**

CAPÍTULO III

- Table 5.1.** Morula and early blastocyst rates on Day-6 after culture in mSOF + 6 mg mL⁻¹ BSA in groups with or without 0.1% fetal calf serum (FCS) **154**
- Table 5.2.** *In vitro* development of Day-6 morulae and early blastocysts produced in mSOF+ 6 mg mL⁻¹ BSA in groups with or without 0.1% FCS, and after single culture without protein up to Day-7 **154**
- Table 5.3.** *In vitro* survival after vitrification/warming of Day-7 bovine expanded blastocysts derived from morulae and early blastocysts produced in mSOF + 6 mg mL⁻¹ BSA with or without 0.1% FCS from Day-1 to Day-6 after one-day single culture without protein **155**
- Table 5.4.** Descriptive Day-40 and Day-60 pregnancy status of heifers **159**

transferred with embryos cultured from Day-1 to Day-6 in mSOF + 6 mg mL⁻¹ BSA, with or without 0.1% FCS and after one-day single culture without protein

CAPÍTULO IV

- Table 6.1.** Sequence of peptides from bovine hepatoma-derived growth factor detected in the multiple reaction monitoring study and their position in the protein **179**
- Table 6.2.** *In vitro* development of bovine morulae and early blastocysts cultured individually from day 6 to day 8 in droplets of synthetic oviduct fluid with BSA containing 100 ng mL⁻¹ recombinant HDGF (rHDGF) **187**
- Table 6.3.** *In vitro* development of bovine morulae and early blastocysts cultured individually from day 6 to day 8 in droplets of synthetic oviduct fluid with PVA containing 100 ng mL⁻¹ recombinant HDGF (rHDGF) **187**
- Table 6.4.** Pregnancy and calving rates of recipients transferred 1-2 times and 3-4 times with bovine vitrified Day-7 expanded blastocysts individually cultured from Day-6 to Day-7 in 12 µL of synthetic oviduct fluid medium + 0.5 mg mL⁻¹ PVA with or without rHDGF (100 ng mL⁻¹) **189**
- Table 6.5.** Weight and morphometry of calves born after transfer of bovine vitrified/warmed Day-7 expanded blastocysts cultured individually with or without r-HDGF (100 ng mL⁻¹) from Day-6 to Day-7 **189**