

Content

List of Figures	III
List of Tables	IV
Acknowledgements	VI
Abstract	VII
Resumen	IX
Resum	XI
1. Introduction	1
1.1. Toxoplasmosis	1
1.2. Bovine Babesiosis	3
1.3. Epidemiological mathematical models	4
2. Mathematical Model of Toxoplasmosis Disease in Varying Size Population	5
2.1. Introduction	5
2.2. Mathematical model	7
2.3. Scaling the model	8
2.4. Numerical simulation	11
2.5. Conclusions	14
3. Dynamics of a model of Toxoplasmosis disease in human and cat populations	15
3.1. Introduction	15
3.2. Mathematical model	18
3.3. Stability analysis of the model	20
3.3.1. Scaling model	20
3.3.2. Equilibrium points of the model	21
3.3.3. Stability analysis	22
3.4. Numerical simulations	25
3.5. Discussion and conclusions	27
4. Modeling dynamics of Babesiosis disease in bovines and ticks populations	30

4.1. Introduction	30
4.2. Mathematical model	31
4.3. Analysis of the model	34
4.3.1. Model equilibrium points	34
4.3.2. Stability analysis	35
4.4. Numerical simulations	37
4.5. Conclusions	41

Bibliography	43
---------------------	-----------

List of Figures

2.1.	Diagram of the Toxoplasmosis disease on the population.	8
2.2.	Dynamics of the different subpopulations $\beta= 0.0206055$ and $\gamma = 0.000232$	11
2.3.	Dynamics of the different subpopulations $\beta= 0.010327$ and $\gamma = 0.000232$	12
2.4.	Dynamics of the different subpopulations $\beta= 0.0206055$ and $\gamma = 0.1$	13
3.1.	Flow diagram of the Toxoplasmosis disease model for human and cat populations as defined in system (3.1).	19
3.2.	Dynamics of the different subpopulations when $\beta_c = 0.00066$, $\gamma = 0.000232$ and $\mathcal{R}_0 = 0.909$. ($X_1^* = 1, Y_1^* = 0$ and $A_1^* = 1$)	26
3.3.	Dynamics of the different subpopulations when $\beta_c = 0.0008$, $\gamma = 0.000232$ and $\mathcal{R}_0 = 1.2121$. ($X^* = 0.0603, Y^* = 0.9386$ and $A^* = 0.825$)	27
3.4.	Dynamics of the different subpopulations when $\beta_c = 0.0008$, $\gamma = 0.000232$, $p_c = 0.0001$ and $\mathcal{R}_0 = 121.21$. ($X^* = 0.0112, Y^* = 0.9878$ and $A^* = 0.00825$)	27
4.1.	Dynamics of the Babesiosis disease in bovine and tick populations	33
4.2.	Ecologic stability, when $\beta_B = 0.00061$, $\beta_T = 0.00048$, $\alpha_B = 0.001$ and $\mathcal{R}_0 = 6.8$	39
4.3.	Dynamics of the different subpopulations when $\beta_B = 0.0003$, $\lambda_B = 0.0265$, $p = 0.1$ and $\mathcal{R}_0 = 0.33$	39
4.4.	Dynamics of the different subpopulations when $\beta_B = 0.006$ and $p = 0.1$, with initial condition $S_B(0) = 0.3816$, $I_B(0) = 0.5184$, $S_T(0) = 0.60$ and $\mathcal{R}_0 = 67.54$	40
4.5.	Dynamics of the different subpopulations when $\beta_B = 0.006$ and $p = 0.1$, with initial condition $S_B(0) = 0.70$, $I_B(0) = 0.10$, $S_T(0) = 0.20$ and $\mathcal{R}_0 = 67.54$	40
4.6.	The dynamics of the subpopulations is sensitive to the parameters β_B and β_T	41

List of Tables

2.1.	Basic parameters of the mathematical model	8
2.2.	Parameters of the model	10
2.3.	Equilibrium point when $\beta = 0.0206055$ and $\gamma = 0.000232$ of system (2.11)	11
2.4.	Eigenvalues of the Jacobian $J(s^*, e^*)$ when $\beta=0.0206055$ and $\gamma=0.000232$ of system (2.11)	11
2.5.	Equilibrium point when $\beta = 0.010327$ and $\gamma = 0.000232$ of system (2.11)	12
2.6.	Eigenvalues of the Jacobian $J(s^*, e^*)$ when $\beta=0.010327$ and $\gamma=0.000232$ of system (2.11)	13
2.7.	Equilibrium point when $\beta= 0.0206055$ and $\gamma = 0.1$ of system (2.11)	13
2.8.	Eigenvalues of the Jacobian $J(s^*, e^*)$ when $\beta= 0.0206055$ and $\gamma = 0.1$ of system (2.11)	14
3.1.	Parameters of the model	26
4.1.	Parameters for simulation. The values β_B, β_T and α_B have been calculated assuming an ecologic stability.	38