




Article

Nondifferentiable G -Mond–Weir Type Multiobjective Symmetric Fractional Problem and Their Duality Theorems under Generalized Assumptions

Ramu Dubey ^{1,†} , Lakshmi Narayan Mishra ^{2,3,†}  and Luis Manuel Sánchez Ruiz ^{4,*,†} 

¹ Department of Mathematics, J C Bose University of Science and Technology, YMCA, Faridabad 121 006, Haryana, India; rdubeyjiya@gmail.com

² Department of Mathematics, School of Advanced Sciences, Vellore Institute of Technology (VIT) University, Vellore 632 014, Tamil Nadu, India; lakshminarayanmishra04@gmail.com

³ L. 1627 Awadh Puri Colony, Beniganj, Phase III, Opposite Industrial Training Institute (I.T.I.), Ayodhya main road, Faizabad 224 001, Uttar Pradesh, India

⁴ ETSID- Departamento de Matemática Aplicada & CITG, Universitat Politècnica de Valencia, E-46022 Valencia, Spain

* Correspondence: lmsr@mat.upv.es

† These authors contributed equally to this work.

Received: 29 September 2019; Accepted: 20 October 2019; Published: 1 November 2019



Abstract: In this article, a pair of nondifferentiable second-order symmetric fractional primal-dual model (G -Mond–Weir type model) in vector optimization problem is formulated over arbitrary cones. In addition, we construct a nontrivial numerical example, which helps to understand the existence of such type of functions. Finally, we prove weak, strong and converse duality theorems under aforesaid assumptions.

Keywords: multiobjective; symmetric duality; second-order; nondifferentiable; fractional programming; support function; G_f -bonvexity/ G_f -pseudobonvexity

1. Introduction

In multiobjective programming problems, convexity plays an important role in deriving optimality conditions and duality results. To relax convexity assumptions involved in sufficient optimality conditions and duality theorems, various generalized convexity notions have been proposed. Multiobjective type programming problem [1] is common in mathematical modeling of realistic phenomenon with a wide spectrum of utilization. Symmetric duality in nonlinear programming deals with the situation where dual of the dual is primal. Special dual problems of optimization are applied to many types of optimization problems. They are used for the proof of optimality of solutions, for designing and a theoretical justification of optimization algorithms, and for physical or economic interpretation of received solutions. Quite often dual problems introduce new meaning to modeled problems. For many interesting applications and developments of multiobjective optimization, we refer to the work of A. Chinchuluun and P.M. Pardalos [2] and the references cited therein.

In economics, we often come across a case where we have to maximize the efficiency of an economic system resulting optimization problems whose objective function is a ratio. Mangasarian [3] proposed the idea of second-order duality for nonlinear optimization problems. The perusal of second-order duality is important due to the computer simulation benefit over the first-order duality since this one supplies narrow ranges for the cost of the objectives when estimations are applied. Suneja et al. [1] and Kim et al. [4] extended the concept of symmetric duality to arbitrary cones.

Suneja et al. [5] considered a pair of multiobjective second order symmetric dual problems of Mond–Weir type without non-negativity constraints and established duality results under η -bonvexity and η -pseudobonvexity assumptions. Later, Khurana [6] defined cone-pseudoinvex and strongly cone-pseudoinvex functions and proved duality theorems for a pair of Mond–Weir type symmetric dual multiobjective programs over arbitrary cones. For more information on fractional programming, readers are advised to see [7–13].

The purpose of the present work is to study second order multiobjective fractional symmetric duality over arbitrary cones for nondifferentiable G -Mond–Weir type program under G_f -bonvexity/ G_f -pseudobonvexity assumptions. The paper is organized as follows. In Section 2, we present some relevant preliminaries. In Section 3, we consider a pair of G -Mond–Weir type nondifferentiable multiobjective second order fractional symmetric dual problems with cone constraints and establish appropriate duality theorems under aforesaid assumptions followed by conclusions.

2. Preliminaries and Definitions

Throughout this paper, R^n stands for the n -dimensional Euclidean space and R_+^n for its non-negative orthant. Consider the following vector minimization problem:

$$\begin{aligned} \text{(MP)} \quad & \text{Minimize } f(x) = \left\{ f_1(x), f_2(x), f_3(x), \dots, f_k(x) \right\}^T \\ & \text{Subject to } X^0 = \{x \in X \subset R^n : g_j(x) \leq 0, j = 1, 2, \dots, m\} \end{aligned}$$

where $f = \{f_1, f_2, \dots, f_k\} : X \rightarrow R^k$ and $g = \{g_1, g_2, \dots, g_m\} : X \rightarrow R^m$ are differentiable functions defined on X .

Definition 1. A point $\bar{x} \in X^0$ is said to be an efficient solution of (MP) if there exists no other $x \in X^0$ such that $f_r(x) < f_r(\bar{x})$, for some $r = 1, 2, \dots, k$ and $f_i(x) \leq f_i(\bar{x})$, for all $i = 1, 2, \dots, k$.

Definition 2. The positive polar cone S^* of a cone $S \subseteq R^s$ is defined by

$$S^* = \{y \in R^s : x^T y \geq 0, \text{ for all } x \in S\}.$$

Let $C_1 \subseteq R^n$ and $C_2 \subseteq R^m$ be closed convex cones with non-empty interiors and S_1 and S_2 be non-empty open sets in R^n and R^m , respectively, such that $C_1 \times C_2 \subseteq S_1 \times S_2$. Suppose $f = (f_1, f_2, \dots, f_k) : S_1 \times S_2 \rightarrow R^k$ is a vector-valued differentiable function.

Definition 3. The function f is said to be invex at $u \in S_1$ (with respect to η , where $\eta : S_1 \times S_2 \rightarrow R^n$), if $\forall x \in S_1$ and for fixed $v \in S_2$, we have

$$f_i(x, v) - f_i(u, v) \geq \eta^T(x, u) \nabla_x f_i(u, v), \text{ for all } i = 1, 2, \dots, k,$$

If the above inequality sign changes to \leq , then f is called incave at $u \in S_1$ with respect to η .

Definition 4. The function f is said to be pseudoinvex at $u \in S_1$ (with respect to η , where $\eta : S_1 \times S_2 \rightarrow R^n$), if $\forall x \in S_1$ and for fixed $v \in S_2$, we have

$$\eta^T(x, u) \nabla_x f_i(u, v) \geq 0 \Rightarrow f_i(x, v) - f_i(u, v) \geq 0, \text{ for all } i = 1, 2, \dots, k.$$

If the above inequality sign changes to \leq , then f is called pseudoincave at $u \in X$ with respect to η .

Definition 5. The function f is said to be G_f -invex at $u \in S_1$ (with respect to η), if there exists a differentiable function $G_f = (G_{f_1}, G_{f_2}, \dots, G_{f_k}) : R \rightarrow R^k$ such that each component $G_{f_i} : I_{f_i}(S_1 \times S_2) \rightarrow R$, where $I_{f_i}(S_1 \times S_2)$, $i = 1, 2, 3, \dots, k$ is the range of f_i , is strictly increasing on its domain and $\eta : S_1 \times S_2 \rightarrow R^n$, so that $\forall x \in S_1$, for fixed $v \in S_2$, we have

$$G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) \geq \eta^T(x, u)G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v), \text{ for all } i = 1, 2, \dots, k,$$

If the above inequality sign changes to \leq , then f is called G_f -incave at $u \in S_1$ with respect to η .

Definition 6. The function f is said to be G_f -pseudoinvex at $u \in S_1$ (with respect to η), if there exists a differentiable function $G_f = (G_{f_1}, G_{f_2}, \dots, G_{f_k}) : R \rightarrow R^k$ such that each component $G_{f_i} : I_{f_i}(S_1 \times S_2) \rightarrow R$, where $I_{f_i}(S_1 \times S_2)$, $i = 1, 2, 3, \dots, k$ is the range of f_i , is strictly increasing on its domain and $\eta : S_1 \times S_2 \rightarrow R^n$, so that $\forall x \in S_1$, for fixed $v \in S_2$, we have

$$\eta^T(x, u)G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v) \geq 0 \Rightarrow G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) \geq 0, \text{ for all } i = 1, 2, \dots, k.$$

If the above inequality sign changes to \leq , then f is called G_f -pseudoincave at $u \in X$ with respect to η .

Definition 7. The function f is said to be G_f -bonvex at $u \in S_1$ (with respect to η), if there exists a differentiable function $G_f = (G_{f_1}, G_{f_2}, \dots, G_{f_k}) : R \rightarrow R^k$ such that each component $G_{f_i} : I_{f_i}(S_1 \times S_2) \rightarrow R$, where $I_{f_i}(S_1 \times S_2)$, $i = 1, 2, 3, \dots, k$ is the range of f_i , is strictly increasing on its domain and $\eta : S_1 \times S_2 \rightarrow R^n$, so that $\forall x \in S_1$, for fixed $v \in S_2$ and $p_i \in R^n$, we have

$$\begin{aligned} G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) &\geq \eta^T(x, u)[G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v) + \{G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v)(\nabla_x f_i(u, v))^T \\ &+ G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v)\}p_i] - \frac{1}{2}p_i^T[G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v)(\nabla_x f_i(u, v))^T \\ &+ G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v)]p_i, \text{ for all } i = 1, 2, \dots, k. \end{aligned}$$

If the above inequality sign changes to \leq , then f is called G_f -boncave at $u \in S_1$ with respect to η .

Definition 8. The function f is said to be G_f -pseudobonvex at $u \in S_1$ (with respect to η), if there exists a differentiable function $G_f = (G_{f_1}, G_{f_2}, \dots, G_{f_k}) : R \rightarrow R^k$ such that each component $G_{f_i} : I_{f_i}(S_1 \times S_2) \rightarrow R$, where $I_{f_i}(S_1 \times S_2)$, $i = 1, 2, 3, \dots, k$ is the range of f_i , is strictly increasing on its domain and $\eta : S_1 \times S_2 \rightarrow R^n$, so that $\forall x \in S_1$, for fixed $v \in S_2$ and $p_i \in R^n$, $\eta^T(x, u)[G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v) + \{G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v)(\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v))$

$$\begin{aligned} \nabla_{xx} f_i(u, v)\}p_i] \geq 0 \Rightarrow G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) &+ \frac{1}{2}p_i^T[G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v)(\nabla_x f_i(u, v))^T \\ &+ G'_{f_i}(f_i(u, v))\nabla_{xx} f_i(u, v)]p_i \geq 0, \text{ for all } i = 1, 2, \dots, k. \end{aligned}$$

If the above inequality sign changes to \leq , then f is called G_f -pseudoboncave at $u \in S_1$ with respect to η .

We now give an example of G_f -bonvexity with respect to η , but not η -bonvex.

Example 1. Let $k = 4, n = 1, S_1 = S_2 = \left[-\frac{\pi}{6}, \frac{\pi}{6}\right], C_1 = C_2 = \left[\frac{-\pi}{6}, \frac{\pi}{6}\right]$.

Let $f : \left[-\frac{\pi}{6}, \frac{\pi}{6}\right] \times \left[-\frac{\pi}{6}, \frac{\pi}{6}\right] \rightarrow R^4$ be defined as

$$f(x, y) = \{f_1(x, y), f_2(x, y), f_3(x, y), f_4(x, y)\},$$

where $f_1(x, y) = e^y$, $f_2(x, y) = xe^y$, $f_3(x, y) = x^2 \sin^2 y$, $f_4(x, y) = y^2$ and $G_f = \{G_{f_1}, G_{f_2}, G_{f_3}, G_{f_4}\} : R \rightarrow R^4$ be defined as:

$$G_{f_1}(t) = t, G_{f_2}(t) = t^4, G_{f_3}(t) = t, G_{f_4}(t) = t^2.$$

Let $\eta : \left[-\frac{\pi}{6}, \frac{\pi}{6}\right] \times \left[-\frac{\pi}{6}, \frac{\pi}{6}\right] \rightarrow R$ be given as:

$$\eta(x, u) = xu.$$

To show that f is G_f -bonvex at $u = 0$ with respect to η , we have to claim that

$$\begin{aligned} \pi_i &= G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) - \eta^T(x, u)[G'_{f_i}(f_i(u, v))\nabla_x f_i(u, v) + \{G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v) \\ &(\nabla_x f_i(u, v))^T\} + G'_{f_i}(f_i(u, v))\nabla_{xx} f_i(u, v)]p_i + \frac{1}{2}p_i^T[G''_{f_i}(f_i(u, v))\nabla_x f_i(u, v) \\ &(\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v))\nabla_{xx} f_i(u, v)]p_i \geq 0, \quad i = 1, 2, 3, 4. \end{aligned}$$

Putting the values of $f_1, f_2, f_3, f_4, G_{f_1}, G_{f_2}, G_{f_3}, G_{f_4}$ and $u = 0$ in the above expressions, we have

$$\begin{aligned} \pi_1 &= 0, \quad \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right], \\ \pi_2 &= x^4 e^{4v}, \quad \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right], \\ \pi_3 &= x^2 \sin^2 v, \quad \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right], \end{aligned}$$

and

$$\pi_4 = 0, \quad \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right].$$

Hence, $\pi_1 \geq 0, \pi_2 \geq 0$ (from Figure 1), $\pi_3 \geq 0$ (in Figure 2) and $\pi_4 \geq 0, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$ and $\forall p$.

Therefore, f is G_f -bonvex at $u = 0$ with respect to η and p .

Next, we claim that function f is not η -bonvex. For this, it is sufficient to prove that at least one f'_i s is not η -bonvex.

Let

$$\xi = f_3(x) - f_3(u) - \eta^T(x, u)[\nabla_x f_3(u) - \nabla_{xx} f_3(u)]p_3 + \frac{1}{2}p_3^T[\nabla_{xx} f_3(u)]p_3$$

or

$$\xi = xe^v - ue^v - 0, \quad \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right],$$

$$\xi = xe^v \text{ at } u = 0 \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right].$$




Figure 1. The function $\pi_2 = x^4 e^{4v}, \forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$ is non-negative.

It follows that $\xi \not\geq 0, u \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$ and $\forall p$ (in Figure 3). Therefore, f_3 is not η -bonvex at $u = 0$ with respect to p_3 . Hence, $f = (f_1, f_2, f_3, f_4)$ is not η -bonvex at $u = 0$ with respect to p .

Definition 9. Let C be a compact convex set in R^n . The support function of C is defined by

$$s(x|C) = \max\{x^T y : y \in C\}.$$

The subdifferential of $s(x|C)$ is given by

$$\partial s(x|C) = \{z \in C : z^T x = s(x|C)\}.$$

For any convex set $S \subset \mathbb{R}^n$, the normal cone to S at a point $x \in S$ is defined by

$$N_S(x) = \{y \in \mathbb{R}^n : y^T(z - x) \leq 0 \text{ for all } z \in S\}.$$

It is readily verified that for a compact convex set S , y is in $N_S(x)$ if and only if

$$s(y|S) = x^T y.$$

Suppose that $S_1 \subseteq \mathbb{R}^n$ and $S_2 \subseteq \mathbb{R}^m$ are open sets such that $C_1 \times C_2 \subset S_1 \times S_2$.




Figure 2. The function $\pi_3 = x^2 \sin^2 v$, $\forall p, \forall x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$ is non-negative.




Figure 3. The function $\zeta = xe^v$ becomes negative at some $x, v \in \left[-\frac{\pi}{6}, \frac{\pi}{6}\right]$.

3. Second-Order Nondifferentiable Multiobjective Symmetric Fractional Programming Problem Over Arbitrary Cones

Now, we consider the following pair of a nondifferentiable multiobjective second-order fractional symmetric dual program over arbitrary cones

$$(GMFP) \text{ Minimize } U(x, y, z, r, p) = (U_1(x, y, z_1, r_1, p_1), U_2(x, y, z_2, r_2, p_2), \dots, U_k(x, y, z_k, r_k, p_k))^T$$

subject to

$$-\sum_{i=1}^k \lambda_i [G'_{f_i}(f_i(x, y)) \nabla_y f_i(x, y) - z_i + \{G''_{f_i}(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T + G'_{f_i}(f_i(x, y))$$

$$\nabla_{yy} f_i(x, y) \} p_i - U_i(x, y, p_i) \{ G'_{g_i}(g_i(x, y)) \nabla_y g_i(x, y) + r_i + \{ G''_{g_i}(g_i(x, y)) \nabla_y g_i(x, y) \} (\nabla_y g_i(x, y))^T + G'_{g_i}(g_i(x, y)) \nabla_{yy} g_i(x, y) \} p_i \} \in C_2^*,$$

$$y^T \left[\sum_{i=1}^k \lambda_i \left\{ G'_{f_i}(f_i(x, y)) \nabla_y f_i(x, y) - z_i + \{ G''_{f_i}(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T + G'_{f_i}(f_i(x, y)) \nabla_{yy} f_i(x, y) \} p_i - U_i(x, y, p_i) \{ G'_{g_i}(g_i(x, y)) \nabla_y g_i(x, y) + r_i + \{ G''_{g_i}(g_i(x, y)) \nabla_y g_i(x, y) \} (\nabla_y g_i(x, y))^T + G'_{g_i}(g_i(x, y)) \nabla_{yy} g_i(x, y) \} p_i \right\} \right] \geq 0,$$

$x \in C_1, \lambda > 0, z_i \in D_i, r_i \in F_i, i = 1, 2, \dots, k.$

(GMFD) Maximize $T(u, v, w, t, q) = (T_1(u, v, w_1, t_1, q_1), (T_2(u, v, w_2, t_2, q_2), \dots, T_k(u, v, w_k, t_k, q_k))^T$

subject to

$$\sum_{i=1}^k \lambda_i [G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + w_i + G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)] q_i - T_i(u, v, q_i) [G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) - t_i + \{ G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) \} (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)] q_i \in C_1^*,$$

$$u^T \left[\sum_{i=1}^k \lambda_i \left\{ G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) - w_i + G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v) \right\} q_i - T_i(u, v, q_i) [G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) - t_i + \{ G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) \} (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)] q_i \right] \leq 0,$$

$v \in C_1, \lambda > 0, w_i \in Q_i, t_i \in E_i, i = 1, 2, \dots, k.$

where

$$U_i(x, y, z_i, r_i, p_i) = \frac{G_{f_i}(f_i(x, y)) + s(x|Q_i) - y^T z_i - \frac{1}{2} p_i^T [G''_{f_i}(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T]}{G_{g_i}(g_i(x, y)) - s(x|E_i) + y^T r_i - \frac{1}{2} p_i^T [G''_{g_i}(g_i(x, y)) \nabla_y g_i(x, y) (\nabla_y g_i(x, y))^T]} + \frac{G'_{f_i}(f_i(x, y)) \nabla_{yy} f_i(x, y) p_i}{G'_{g_i}(g_i(x, y)) \nabla_{yy} g_i(x, y) p_i}$$

and

$$T_i(u, v, w_i, t_i, q_i) = \frac{G_{f_i}(f_i(u, v)) - s(v|D_i) + u^T w_i - \frac{1}{2} q_i^T [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T]}{G_{g_i}(g_i(u, v)) + s(v|F_i) - u^T t_i - \frac{1}{2} q_i^T [G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T]} + \frac{G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v) q_i}{G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v) q_i};$$

and

$S_1 \subseteq R^n$ and $S_2 \subseteq R^m$; C_1 and C_2 are arbitrary cones in R^n and R^m , respectively, such that $C_1 \times C_2 \subseteq S_1 \times S_2$; $f_i : S_1 \times S_2 \rightarrow R$ and $g_i : S_1 \times S_2 \rightarrow R$ are differentiable functions; $G_{f_i} : I_{f_i} \rightarrow R$ and $G_{g_i} : I_{g_i} \rightarrow R$ are differentiable strictly increasing functions on their domains; Q_i, E_i are compact

convex sets in R^n ; and D_i, F_i are compact convex sets in $R^m, i = 1, 2, 3, \dots, k$. C_1^* and C_2^* are positive polar cones of C_1 and C_2 , respectively. It is assumed that in the feasible regions, the numerators are nonnegative and denominators are positive. p_i and q_i are vectors in R^m and R^n , respectively, $\lambda \in R^k$.

Equivalently, the above problem is reduced in the given form:

(EGMFP) Min $R(x, y, z, r, p) = (R_1(x, y, z_1, r_1, p_1), R_2(x, y, z_2, r_2, p_2), \dots, R_k(x, y, z_k, r_k, p_k))$
 subject to

$$G_{f_i}(f_i(x, y)) + s(x|Q_i) - y^T z_i - \frac{1}{2} p_i^T [G_{f_i}''(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T + G_{f_i}'(f_i(x, y)) \nabla_{yy} f_i(x, y)] p_i - R_i(x, y, z_i, r_i, p_i) [G_{g_i}(g_i(u, v)) - s(x|E_i) + y^T r_i - \frac{1}{2} q_i^T [G_{g_i}''(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G_{g_i}'(g_i(u, v)) \nabla_{xx} g_i(u, v)] q_i] = 0, i = 1, 2, \dots, k, \tag{1}$$

$$-\sum_{i=1}^k \lambda_i [G_{f_i}'(f_i(x, y)) \nabla_y f_i(x, y) - z_i + [G_{f_i}''(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T + G_{f_i}'(f_i(x, y)) \nabla_{yy} f_i(x, y)] p_i - R_i(x, y, z_i, r_i, p_i) \{G_{g_i}'(g_i(x, y)) + r_i \nabla_y g_i(x, y) + (G_{g_i}''(g_i(x, y)) \nabla_y g_i(x, y) (\nabla_y g_i(x, y))^T + G_{g_i}'(g_i(x, y)) \nabla_{yy} g_i(x, y)) p_i\}] \in C_2^*, \tag{2}$$

$$y^T \sum_{i=1}^k \lambda_i [G_{f_i}'(f_i(x, y)) \nabla_y f_i(x, y) - z_i + [G_{f_i}''(f_i(x, y)) \nabla_y f_i(x, y) (\nabla_y f_i(x, y))^T + G_{f_i}'(f_i(x, y)) \nabla_{yy} f_i(x, y)] p_i - R_i(x, y, z_i, r_i, p_i) \{G_{g_i}'(g_i(x, y)) \nabla_y g_i(x, y) + r_i + (G_{g_i}''(g_i(x, y)) \nabla_y g_i(x, y) (\nabla_y g_i(x, y))^T + G_{g_i}'(g_i(x, y)) \nabla_{yy} g_i(x, y)) p_i\}] \geq 0, \tag{3}$$

$$x \in C_1, \lambda > 0, z_i \in D_i, r_i \in F_i, i = 1, 2, \dots, k. \tag{4}$$

(EGMFD) Maximize $S(u, v, w, t, q) = [S_1(u, v, w_1, t_1, q_1), S_2(u, v, w_2, t_2, q_2), \dots, S_k(u, v, w_k, t_k, q_k)]$

subject to

$$G_{f_i}'(f_i(u, v)) - s(v|D_i) + u^T w_i - \frac{1}{2} q_i^T \{G_{f_i}''(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G_{f_i}'(f_i(u, v)) \nabla_{xx} f_i(u, v)\} q_i - S_i(u, v, w_i, t_i, q_i) [G_{g_i}'(g_i(u, v)) + s(v|E_i) - u^T t_i - \frac{1}{2} q_i^T \{G_{g_i}''(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G_{g_i}'(g_i(u, v)) \nabla_{xx} g_i(u, v)\} q_i] = 0, i = 1, 2, \dots, k. \tag{5}$$

$$\sum_{i=1}^k \lambda_i [G_{f_i}'(f_i(u, v)) \nabla_x f_i(u, v) + w_i + \{G_{f_i}''(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G_{f_i}'(f_i(u, v)) \nabla_{xx} f_i(u, v)\} q_i - T_i(u, v, w_i, t_i, q_i) \{G_{g_i}'(g_i(u, v)) \nabla_x g_i(u, v) - t_i + G_{g_i}''(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G_{g_i}'(g_i(u, v)) \nabla_{xx} g_i(u, v)\} q_i] \in C_1^*, \tag{6}$$

$$\begin{aligned}
& u^T \sum_{i=1}^k \lambda_i [G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + w_i + \{G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + \\
& G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)\} q_i - T_i(u, v, w_i, t_i, q_i) \{G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) - t_i + G''_{g_i}(g_i(u, v)) \\
& \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)\} q_i] \leq 0, \tag{7}
\end{aligned}$$

$$v \in C_1, \lambda > 0, w_i \in B_i, t_i \in E_i, i = 1, 2, \dots, k. \tag{8}$$

Let Z^0 and W^0 be the sets of feasible solutions of (EGMFP) and (EGMFD), respectively. Next, we prove duality theorems for (EGMFP) and (EGMFD), which equally apply to (GMFP) and (GMFD), respectively. Let $z = (z_1, z_2, \dots, z_k)$, $r = (r_1, r_2, \dots, r_k)$, $w = (w_1, w_2, \dots, w_k)$, $t = (t_1, t_2, \dots, t_k)$ and $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$.

Theorem 1. (Weak Duality). Let $(x, y, R, z, r, \lambda, p) \in Z^0$ and $(u, v, S, w, t, \lambda, q) \in W^0$. Assume that for $i = 1, 2, 3, \dots, k$:

- (i) $f_i(\cdot, v)$ is G_{f_i} -convex and $(\cdot)^T w_i$ is invex at u for fixed v with respect to η_1 .
- (ii) $g_i(\cdot, v)$ is a G_{g_i} -concave and $(\cdot)^T t_i$ is invex at u for fixed v with respect to η_1 .
- (iii) $f_i(x, \cdot)$ is a G_{f_i} -concave and $(\cdot)^T z_i$ is invex at y for fixed x with respect to η_2 .
- (iv) $g_i(x, \cdot)$ is a G_{g_i} -convex and $(\cdot)^T r_i$ is invex at y for fixed x with respect to η_2 .
- (v) $\eta_1(x, u) + u \in C_1$ and $\eta_2(v, y) + y \in C_2$.
- (vi) $G_{g_i}((x, v)) + v^T r_i - x^T t_i > 0$.

Then, the following can not hold simultaneously:

$$R_i \leq S_i, \text{ for all } i = 1, 2, 3, \dots, k \text{ and } R_j < S_j, \text{ for some } j = 1, 2, 3, \dots, m.$$

Proof. From Assumption (v) and Equation (6), we get

$$\begin{aligned}
& (\eta_1(x, u) + u)^T \sum_{i=1}^k \lambda_i [G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + w_i + [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T \\
& + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)] q_i - T_i(u, v, q_i) \{G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) - t_i \\
& + (G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v) q_i\} \geq 0. \tag{9}
\end{aligned}$$

Using Equations (7) and (9), we obtain,

$$\begin{aligned}
& \eta_1^T(x, u) \left[\sum_{i=1}^k \lambda_i (G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + w_i + [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T \right. \\
& + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)] q_i - T_i(u, v, q_i) \{G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) - t_i \\
& \left. + (G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v) q_i) \right] \geq 0. \tag{10}
\end{aligned}$$

From Assumption (i), we have

$$\begin{aligned}
& G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) \geq n_1^T(x, u) G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) \\
& (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)] p_i - \frac{1}{2} p_i^T [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) \\
& (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v) p_i], \quad i = 1, 2, \dots, k. \tag{11}
\end{aligned}$$

and

$$x^T w_i - u^T w_i \geq \eta_1^T(x, u) w_i, \quad i = 1, 2, \dots, k. \tag{12}$$

Since $\lambda > 0$ and combining above inequalities, it follows that

$$\begin{aligned} \sum_{i=1}^k [G_{f_i}(f_i(x, v)) + x^T w_i - G_{f_i}(f_i(u, v)) - u^T w_i] &\geq \eta_1^T(x, u) \sum_{i=1}^k \lambda_i [G'_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) + w_i \\ &+ [G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)] p_i \\ &- \frac{1}{2} p_i^T \{G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T + G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v)\} p_i]. \end{aligned} \tag{13}$$

Similarly, from Assumption (ii), we get

$$\begin{aligned} -G_{g_i}(g_i(x, v)) + G_{g_i}(g_i(u, v)) &\geq -\eta_1^T(x, u) [G'_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) + [G''_{g_i}(g_i(u, v)) \\ \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)] p_i + \frac{1}{2} p_i^T \{G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) \\ (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)\} p_i], \quad i = 1, 2, \dots, k, \end{aligned} \tag{14}$$

and

$$x^T t_i - u^T t_i \geq \eta_1^T(x, u) t_i, \quad i = 1, 2, \dots, k. \tag{15}$$

Multiplying by $\lambda_i T_i$ in above inequalities and taking summation over $i = 1, 2, 3, \dots, k$, it follows that

$$\begin{aligned} \sum_{i=1}^k \lambda_i T_i [-G_{g_i}(g_i(x, v)) + x^T t_i + G_{g_i}(g_i(u, v)) - u^T t_i] &\geq -\eta_1^T(x, u) \sum_{i=1}^k \lambda_i T_i [[G'_{g_i}(g_i(u, v)) - t_i + \\ \nabla_x g_i(u, v) [G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)] p_i \\ - \frac{1}{2} p_i^T \{G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T + G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v)\} p_i]. \end{aligned} \tag{16}$$

Adding the inequalities in Equations (13) and (16), we get

$$\begin{aligned} \sum_{i=1}^k \lambda_i [G_{f_i}(f_i(x, v)) - G_{f_i}(f_i(u, v)) - T_i (G_{g_i}(g_i(x, v)) - G_{g_i}(g_i(u, v)))] \\ \geq -\sum_{i=1}^k \frac{\lambda_i q_i^T}{2} [G'_{f_i}(f_i(u, v)) \nabla_{xx} f_i(u, v) + G''_{f_i}(f_i(u, v)) \nabla_x f_i(u, v) (\nabla_x f_i(u, v))^T \\ - T_i \{G'_{g_i}(g_i(u, v)) \nabla_{xx} g_i(u, v) + G''_{g_i}(g_i(u, v)) \nabla_x g_i(u, v) (\nabla_x g_i(u, v))^T\} p_i]. \end{aligned} \tag{17}$$

Since $v^T r_i \leq s(v|F_i)$, from Equations (17) and (5), we get

$$\sum_{i=1}^k \lambda_i [G_{f_i}(f_i(x, v)) + x^T w_i - s(v|D_i) + T_i (x^T t_i - v^T r_i - G_{g_i}(g_i(x, v)))] \geq 0. \tag{18}$$

Similarly, using Hypotheses (iii)–(v) and the primal constraints in Equations (1)–(4), we have

$$\sum_{i=1}^k \lambda_i [-G_{f_i}(f_i(x, v)) + v^T z_i - s(x|Q_i) + R_i (-x^T t_i + v^T r_i + G_{g_i}(g_i(x, v)))] \geq 0. \tag{19}$$

On adding the inequalities in Equations (18) and (19), we get

$$\sum_{i=1}^k \lambda_i [v^T z_i - s(v|D_i) + x^T w_i - G_{f_i}(f_i(x, v)) - s(x|Q_i) + (R_i - S_i)(-x^T t_i + v^T r_i + G_{g_i}(g_i(x, v)))] \geq 0. \quad (20)$$

Since $\lambda_i > 0$, $v^T z_i - s(v|D_i) + x^T w_i - s(x|C_i) \leq 0$, $i = 1, 2, 3, \dots, k$, it yields

$$\sum_{i=1}^k \lambda_i (R_i - T_i)(G_{g_i}(g_i(x, v)) + v^T r_i - x^T t_i) \geq 0.$$

From Assumption (vi), we have, $G_{g_i}((x, v)) + v^T r_i - x^T t_i >$, $i = 1, 2, 3, \dots, k$. Since $\lambda > 0$, it follows that $R \not\leq S$, hence the result. \square

Remark 1. Since every convex function is pseudoconvex, the above weak duality theorem for the symmetric dual pair (EGMFP) and (EGMFD) can also be obtained under pseudobonvexity assumptions.

Theorem 2. (Weak Duality). Let $(x, y, R, z, r, \lambda, p) \in Z^0$ and $(u, v, S, w, t, \lambda, q) \in W^0$. Assume that for $i = 1, 2, 3, \dots, k$:

- (i) $f_i(\cdot, v)$ is G_{f_i} -pseudobonvex and $(\cdot)^T w_i$ is pseudoinvex at u for fixed v with respect to η_1 .
- (ii) $g_i(\cdot, v)$ is a G_{g_i} -pseudobonconcave and $(\cdot)^T t_i$ is pseudoinvex at u for fixed v with respect to η_1 .
- (iii) $f_i(x, \cdot)$ is a G_{f_i} -pseudobonconcave and $(\cdot)^T z_i$ is pseudoinvex at y for fixed x with respect to η_2 .
- (iv) $g_i(x, \cdot)$ is a G_{g_i} -pseudobonvex and $(\cdot)^T r_i$ is pseudoinvex at y for fixed x with respect to η_2 .
- (v) $\eta_1(x, u) + u \in C_1$ and $\eta_2(v, y) + y \in C_2$.
- (vi) $G_{g_i}((x, v)) + v^T r_i - x^T t_i > 0$.

Then, the following cannot hold simultaneously:

$$R_i \leq S_i, \text{ for all } i = 1, 2, 3, \dots, k \text{ and } R_j < S_j, \text{ for some } j = 1, 2, 3, \dots, m.$$

Proof. The proof follows on the lines of Theorem 1. \square

Theorem 3. (Strong Duality). Let $(\bar{x}, \bar{y}, \bar{R}, \bar{z}, \bar{r}, \bar{\lambda}, \bar{p})$ be an efficient solution to (EGMFP), fix $\lambda = \bar{\lambda}$ in (EGMFD). Further, assume that

- (i) $\{G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i \{G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T\}\}$ is positive definite

and

$$p_i^T [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + [G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i \{G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T\}] \geq 0, \text{ for all } i = 1, 2, 3, \dots, k.$$

- (ii) The matrix $\left\{ G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i \{G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T\} \right\}$ is positive definite for $i = 1, 2, 3, \dots, k$.

- (iii) For $\beta > 0$ and $\bar{p}_i \in \mathbb{R}^m$, $\bar{p}_i \neq 0$, $i = 1, 2, \dots, k$ implies that

$$\sum_{i=1}^k \beta_i \bar{p}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + [G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i [G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T] \neq 0.$$

(iv) $[G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + \{G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T)\}]_{i=1}^k$ is linearly independent.

(iv) $\bar{R}_i > 0, i = 1, 2, 3, \dots, k.$

Then, there exist $\bar{w}_i \in Q$ and $\bar{t}_i \in E_i, i = 1, 2, 3, \dots, k$ such that $(\bar{x}, \bar{y}, \bar{R}, \bar{w}, \bar{\lambda}, \bar{t}, \bar{q} = 0)$ is feasible for (EGMFD). Furthermore, if the assumptions of Theorem 3.1 or 3.2 are satisfied, then $(\bar{x}, \bar{y}, \bar{R}, \bar{w}, \bar{\lambda}, \bar{t}, \bar{q} = 0)$ is an efficient solution to (EGMFD).

Proof. Since $(\bar{x}, \bar{y}, \bar{R}, \bar{w}, \bar{\lambda}, \bar{t}, \bar{q} = 0)$ is an efficient solution of (EMFP), by Fritz John necessary conditions [14], there exists $\alpha \in \mathbb{R}^k, \beta \in \mathbb{R}_+, \gamma \in \mathbb{C}_{\neq}, \delta \in \mathbb{R}$ and $\zeta \in \mathbb{R}^k$ such that

$$\begin{aligned} & (x - \bar{x})^T \sum_{i=1}^k \beta_i \left[G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i - \frac{1}{2} \bar{p}_i^T \nabla_x \{ [G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T \right. \\ & + G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) \} \bar{p}_i - \bar{R}_i \left(G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) + \bar{t}_i - \frac{1}{2} \bar{p}_i^T \nabla_x \{ [G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) \right. \\ & \left. \left. (\nabla_y g_i(\bar{x}, \bar{y}))^T + G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) \} \bar{p}_i \right) \right] + (\gamma - \delta \bar{y})^T \sum_{i=1}^k \bar{\lambda}_i \left[G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) \nabla_y f_i(\bar{x}, \bar{y}) \right. \\ & + G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{xy} f_i(\bar{x}, \bar{y}) + \nabla_x \{ [G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T + G'_{f_i}(f_i(\bar{x}, \bar{y})) \\ & \nabla_{yy} f_i(\bar{x}, \bar{y}) \} \bar{p}_i - \bar{R}_i \left(G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) \nabla_y g_i(\bar{x}, \bar{y}) + G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{xy} g_i(\bar{x}, \bar{y}) \right. \\ & \left. \left. + \nabla_x \{ [G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T + G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) \} \bar{p}_i \right) \right] \geq 0, \forall x \in C_1, \end{aligned} \tag{21}$$

$$\begin{aligned} & \sum_{i=1}^k \left[(\beta_i - \delta \bar{\lambda}_i) \{ (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) - \bar{z}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + (G''_{f_i}(f_i(\bar{x}, \bar{y})) \right. \\ & \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i ((G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) + \bar{r}_i + (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) \\ & + (G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T) \bar{p}_i) \} + ((\gamma - \delta \bar{y}) \bar{\lambda}_i - \beta_i \bar{p}_i) \{ (G'_{f_i}(f_i(\bar{x}, \bar{y})) \\ & \nabla_{yy} f_i(\bar{x}, \bar{y}) + (G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T) - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) \\ & + (G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T) \} + \left((\gamma - \delta \bar{y}) \bar{\lambda}_i - \frac{\beta_i \bar{p}_i}{2} \right) \{ \nabla_y ((G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) \\ & + (G''_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i (\nabla_y ((G'_{g_i}(g_i(\bar{x}, \bar{y})) \\ & \nabla_{yy} g_i(\bar{x}, \bar{y}) + (G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y})(\bar{x}, \bar{y}))^T) \bar{p}_i) \} \right] = 0, \end{aligned} \tag{22}$$

$$\begin{aligned}
& (\gamma - \delta\bar{y})\{(G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) - \bar{z}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + (G''_{f_i}(f_i(\bar{x}, \bar{y})) \\
& \nabla_y f_i(\bar{x}, \bar{y})(\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i) - \bar{R}_i((G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) + \bar{r}_i + (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) \\
& g_i(\bar{x}, \bar{y}) + (G''_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y})(\nabla_y g_i(\bar{x}, \bar{y}))^T) \bar{p}_i)\} = 0, i = 1, 2, 3, \dots, k, \tag{23}
\end{aligned}$$

$$\begin{aligned}
& (\bar{\lambda}_i(\gamma - \delta\bar{y}) - \beta_i \bar{p}_i)^T [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + G''_{f_i}(f_i(\bar{x}, \bar{y}))(\nabla_y f_i(\bar{x}, \bar{y})) \\
& (\nabla_y f_i(\bar{x}, \bar{y}))^T - \bar{R}_i[G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y})) \\
& (\nabla_y g_i(\bar{x}, \bar{y}))^T]] = 0, i = 1, 2, 3, \dots, k. \tag{24}
\end{aligned}$$

$$\begin{aligned}
& \alpha_i - \beta_i [G_{g_i}(g_i(\bar{x}, \bar{y})) - s(\bar{x}|E_i) + \bar{y}^T \bar{r}_i - \frac{1}{2} \bar{p}_i^T [G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y})) \\
& (\nabla_y g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))^T] p_i] - (\gamma - \delta\bar{y})[\bar{\lambda}_i(G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) + \bar{r}_i + \\
& (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))^T) \bar{p}_i] = 0, i = 1, 2, \dots, k. \tag{25}
\end{aligned}$$

$$\beta_i \bar{y} + (\gamma - \delta\bar{y}) \bar{\lambda}_i \in N_{D_i}(\bar{z}_i), i = 1, 2, \dots, K, \tag{26}$$

$$\beta_i \bar{R}_i \bar{y} + (\gamma - \delta\bar{y}) \bar{R}_i \bar{\lambda}_i \in N_{F_i}(\bar{r}_i), i = 1, 2, 3, \dots, k, \tag{27}$$

$$\begin{aligned}
& \bar{y}^T \sum_{i=1}^k \bar{\lambda}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) - \bar{z}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + G''_{f_i}(f_i(\bar{x}, \bar{y})) \\
& (\nabla_y f_i(\bar{x}, \bar{y}))(\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) + \bar{r}_i + \{G'_{g_i}(g_i(\bar{x}, \bar{y})) \\
& \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))^T) \bar{p}_i]] = 0. \tag{28}
\end{aligned}$$

$$\begin{aligned}
& \delta \bar{y}^T \sum_{i=1}^k \bar{\lambda}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) - \bar{z}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + G''_{f_i}(f_i(\bar{x}, \bar{y})) \\
& (\nabla_y f_i(\bar{x}, \bar{y}))(\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) + \bar{r}_i + \{G'_{g_i}(g_i(\bar{x}, \bar{y})) \\
& \nabla_{yy} g_i(\bar{x}, \bar{y}) + G''_{g_i}(g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))(\nabla_y g_i(\bar{x}, \bar{y}))^T) \bar{p}_i]] = 0. \tag{29}
\end{aligned}$$

$$\bar{\lambda}^T \bar{\zeta} = 0, \tag{30}$$

$$\bar{w}_i \in Q_i, \bar{t}_i \in E_i, \bar{x}^T \bar{t}_i = S(\bar{x}|E_i), \bar{x}^T \bar{w}_i = S(\bar{x}|Q_i), i = 1, 2, 3, \dots, k, \tag{31}$$

$$(\alpha, \delta, \bar{\zeta}) \geq 0, (\alpha, \beta, \gamma, \delta, \bar{\zeta}) \neq 0. \tag{32}$$

From Assumption (i) and Equation (24), we have

$$\gamma \bar{\lambda}_i - \beta_i \bar{p}_i - \bar{\lambda}_i \delta \bar{y} = 0. \tag{33}$$

We claim that $\beta_i \neq 0, \forall i$. The proof is by contradiction. Let $\beta_i = 0$ for some i . Since $\bar{\lambda} > 0$, the relation in Equation (33) yields

$$\gamma = \delta \bar{y}. \quad (34)$$

From the relation in Equations (22), (33) and (34), we obtain

$$\begin{aligned} & \sum_{i=1}^k (\beta_i - \delta \bar{\lambda}_i) [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) - \bar{z}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + \\ & G''_{f_i}(f_i(\bar{x}, \bar{y})) (\nabla_y f_i(\bar{x}, \bar{y})) (\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i [G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) + \bar{r}_i \\ & + \{G''_{g_i}(g_i(\bar{x}, \bar{y})) (\nabla_y g_i(\bar{x}, \bar{y})) (\nabla_y g_i(\bar{x}, \bar{y}))^T + G'_{g_i}(\nabla_{yy} g_i(\bar{x}, \bar{y}))\} \bar{p}_i]] = 0. \end{aligned} \quad (35)$$

On using Assumption (iv), this gives

$$\beta_i - \delta \bar{\lambda}_i = 0, i = 1, 2, \dots, k. \quad (36)$$

Since $\beta_i = 0$, we obtain $\delta \bar{\lambda}_i = 0$ but $\bar{\lambda}_i > 0, i = 1, 2, \dots, k$ and thus the relation in Equation (36) implies $\delta = 0$. Thus, from the relation in Equations (25), (34) and (36), we get $\alpha_i = 0, i = 1, 2, \dots, k$. In addition, from the relation in Equation (34), we get $\gamma = 0$, which is a contradiction, since $(\alpha, \beta, \gamma, \delta) \neq 0$. Hence, we get $\beta_i \neq 0, i = 1, 2, \dots, k$.

Since $\bar{\lambda} > 0$, using Equations (22) and (33), we get

$$\begin{aligned} & \sum_{i=1}^k \beta_i \bar{p}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) + \bar{w}_i + (G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_{yy} f_i(\bar{x}, \bar{y}) + \\ & G''_{f_i}(f_i(\bar{x}, \bar{y})) (\nabla_y f_i(\bar{x}, \bar{y})) (\nabla_y f_i(\bar{x}, \bar{y}))^T) \bar{p}_i - \bar{R}_i [G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_y g_i(\bar{x}, \bar{y}) - \bar{t}_i \\ & + \{G''_{g_i}(g_i(\bar{x}, \bar{y})) (\nabla_y g_i(\bar{x}, \bar{y})) (\nabla_y g_i(\bar{x}, \bar{y}))^T + G'_{g_i}(\nabla_{yy} g_i(\bar{x}, \bar{y}))\} \bar{p}_i]] = 0. \end{aligned} \quad (37)$$

Hence, from Assumption (iii), we get $\bar{p}_i = 0, i = 1, 2, \dots, k$. From the relation in Equation (33), $\bar{p}_i = 0, i = 1, 2, \dots, k$ and $\bar{\lambda} > 0$, we have $\gamma = \delta \bar{y}$, from Equations (21) and (22), we have

$$\sum_{i=1}^k \bar{\lambda}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i - \bar{R}_i [G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) - \bar{t}_i]] = 0. \quad (38)$$

$$\sum_{i=1}^k (\beta_i - \delta \bar{\lambda}_i) [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_y f_i(\bar{x}, \bar{y}) + \bar{w}_i - \bar{R}_i G'_{g_i} \nabla_y g_i(\bar{x}, \bar{y}) - \bar{t}_i] = 0. \quad (39)$$

By Assumptions (i) and (iii), we have

$$\beta_i = \delta \bar{\lambda}_i, i = 1, 2, \dots, k. \quad (40)$$

Since $\beta_i > 0$ and $\bar{\lambda}_i > 0, i = 1, 2, \dots, k$, the relation in Equation (40) implies that $\delta > 0$, and the relation in Equation (38) reduces to

$$(x - \bar{x})^T \sum_{i=1}^k \bar{\lambda}_i [G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) - \bar{t}_i)] \geq 0, \forall x \in C_1. \quad (41)$$

Let $x \in C_1$. Then, $x + \bar{x} \in C_1$ as C_1 is a closed convex cone. On substituting $x + \bar{x}$ into the place of x in Equation (41), we get

$$x^T \sum_{i=1}^k \bar{\lambda}_i [(G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i) - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) - \bar{t}_i)] \geq 0.$$

Hence,

$$\sum_{i=1}^k \bar{\lambda}_i [(G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i) - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) - \bar{t}_i)] \in C_1^*. \tag{42}$$

In addition, by letting $x = 0$ and $x = 2\bar{x}$ simultaneously in Equation (41), we have

$$x^T \sum_{i=1}^k \bar{\lambda}_i [(G'_{f_i}(f_i(\bar{x}, \bar{y})) \nabla_x f_i(\bar{x}, \bar{y}) + \bar{w}_i) - \bar{R}_i (G'_{g_i}(g_i(\bar{x}, \bar{y})) \nabla_x g_i(\bar{x}, \bar{y}) - \bar{t}_i)] = 0. \tag{43}$$

Since $\gamma = \delta \bar{y}$ and $\delta > 0$, we have

$$\bar{y} = \frac{\gamma}{\delta} \in C_2. \tag{44}$$

From Equations (26) and (34) and using $\beta > 0$, we get $\bar{y} \in N_{D_i}(\bar{z}_i)$, $i = 1, 2, 3, \dots, k$. This implies

$$\bar{y}^T \bar{z}_i = S(\bar{y}|D_i), \quad i = 1, 2, 3, \dots, k. \tag{45}$$

Similarly, by Equation (27) and Assumption (iii), $\bar{y} \in N_{F_i}(\bar{r}_i)$, $i = 1, 2, 3, \dots, k$, we obtain

$$\bar{y}^T \bar{r}_i = S(\bar{y}|F_i), \quad i = 1, 2, 3, \dots, k. \tag{46}$$

Combining Equations (31), (45), (46) and (31), it follows that

$$(G_{f_i}(f_i(\bar{x}, \bar{y})) - S(\bar{y}|D_i) + \bar{x}^T \bar{w}_i) - \bar{R}_i (G_{g_i}(g_i(\bar{x}, \bar{y})) + S(\bar{y}|F_i) - \bar{x}^T \bar{t}_i) = 0, \quad i = 1, 2, 3, \dots, k. \tag{47}$$

This together with Equations (42)–(43) and (47) shows that $(\bar{x}, \bar{y}, \bar{R}, \bar{\lambda}, \bar{w}, \bar{t}) \in W^0$. Now, let $(\bar{x}, \bar{y}, \bar{R}, \bar{\lambda}, \bar{w}, \bar{t})$ be not an efficient solution of (EGMFD). Then, there exists other $(u, v, R, \lambda, w, t) \in W^0$ such that $\bar{R}_i \leq S_i, \forall i = 1, 2, \dots, k$ and $\bar{R}_j < S_j$, for some $j = 1, 2, \dots, m$. This contradicts the result of the Theorems 1 and 2. Hence, the proof is complete. \square

Remark 2. In the case of symmetric programming problem, the proof of converse duality theorem remains same as Theorem 3.

Theorem 4. (Converse duality theorem). Let $(\bar{u}, \bar{v}, \bar{S}, \bar{t}, \bar{w}, \bar{\lambda}, \bar{q})$ be an efficient solution to (EGMFD), fix $\lambda = \bar{\lambda}$ in (EGMFP). Further, assume that

- (i) $\{G'_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_{xx} f_i(\bar{u}, \bar{v}) + G''_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_x f_i(\bar{u}, \bar{v})(\nabla_x f_i(\bar{u}, \bar{v}))^T - \bar{S}_i \{G'_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_{xx} g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T\} \}$ is positive definite and
- $q_i^T [G'_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_{xx} f_i(\bar{u}, \bar{v}) + [G''_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_x f_i(\bar{u}, \bar{v})(\nabla_x f_i(\bar{u}, \bar{v}))^T - \bar{S}_i [G'_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_{xx} g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T] \geq 0, \text{ for all } i = 1, 2, 3, \dots, k.$

- (ii) The matrix $\left\{ G'_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_{xx} f_i(\bar{u}, \bar{v}) + [G''_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_x f_i(\bar{u}, \bar{v})(\nabla_x f_i(\bar{u}, \bar{v}))^T - \bar{S}_i [G'_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_{xx} g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T] \right\}$

$\left. \nabla_{xx}g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T \right\}$ is positive definite for $i = 1, 2, 3, \dots, k$.

(iii) For $\beta > 0$ and $\bar{q}_i \in \mathbb{R}^n$, $\bar{q}_i \neq 0$, $i = 1, 2, \dots, k$ implies that

$$\sum_{i=1}^k \beta_i \bar{q}_i [G'_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_{xx} f_i(\bar{u}, \bar{v}) + [G''_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_x f_i(\bar{u}, \bar{v})(\nabla_x f_i(\bar{u}, \bar{v}))^T - \bar{S}_i [G'_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_{xx} g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T] \neq 0,$$

(iv) $[G'_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_{xx} f_i(\bar{u}, \bar{v}) + \{G''_{f_i}(f_i(\bar{u}, \bar{v})) \nabla_x f_i(\bar{u}, \bar{v})(\nabla_x f_i(\bar{u}, \bar{v}))^T - \bar{S}_i (G'_{g_i}(g_i(\bar{u}, \bar{v}))$

$\nabla_{xx}g_i(\bar{u}, \bar{v}) + G''_{g_i}(g_i(\bar{u}, \bar{v})) \nabla_x g_i(\bar{u}, \bar{v})(\nabla_x g_i(\bar{u}, \bar{v}))^T\}]_{i=1}^k$ is linearly independent.

(iv) $\bar{S}_i > 0$, $i = 1, 2, 3, \dots, k$. Then, there exist $\bar{z}_i \in D_i$ and $\bar{r}_i \in E_i$, $i = 1, 2, 3, \dots, k$ such that $(\bar{u}, \bar{v}, \bar{S}, \bar{z}, \bar{\lambda}, \bar{r}, \bar{p} = 0)$ is feasible for (EGMFP). Furthermore, if the assumptions of Theorem 3.1 or 3.2 are satisfied, then $(\bar{u}, \bar{v}, \bar{S}, \bar{z}, \bar{\lambda}, \bar{r}, \bar{p} = 0)$ is an efficient solution to (EGMFP).

Proof. The results can be obtained on the lines of Theorem 3. \square

4. Conclusions

In this paper, we use the concept of G_f -bonvex/ G_f -pseudobonvex functions to establish duality results for G -Mond–Weir type dual model related to multiobjective nondifferentiable second-order symmetric fractional programming problem over arbitrary cones. Numerical examples are also illustrated to justify the existence of such type of functions. The present work can be further extended to nondifferentiable higher-order symmetric fractional programming over cones. This will orient the future task for the researcher working in this area.

Author Contributions: All authors contributed equally in writing this article. All authors read and approved the final manuscript.

Acknowledgments: The authors wish to thank the referees for useful comments.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Suneja, S.K.; Aggarwal, S.; Davar, S. Multiobjective symmetric duality involving cones. *Eur. J. Oper. Res.* **2002**, *141*, 471–479.
2. Chinchuluun, A.; Pardalos, P.M. A survey of multiobjective optimization. *Ann. Oper. Res.* **2007**, *154*, 29–50.
3. Mangasarian, O.L. Second and higher-order duality in nonlinear programming. *J. Math. Anal. Appl.* **1975**, *51*, 607–620.
4. Kim, D.S.; Yun, Y.B.; Lee, W.J. Multiobjective symmetric duality with cone constraints. *Eur. J. Oper. Res.* **1998**, *107*, 686–691.
5. Suneja, S.K.; Lalitha, C.S.; Khurana, S. Second order symmetric duality in multiobjective programming. *Eur. J. Oper. Res.* **2003**, *144*, 492–500.
6. Khurana, S. Symmetric duality in multiobjective programming involving generalized cone-invex functions. *Eur. J. Oper. Res.* **2005**, *165*, 592–597.
7. Antcza, T. New optimality conditions and duality results of G -type in differentiable mathematical programming. *Nonlinear Anal.* **2007**, *66*, 1617–1632.
8. Dubey, R.; Mishra, L.N.; Mishra, V.N. Duality relations for a class of a multiobjective fractional programming problem involving support functions. *Am. J. Oper. Res.* **2018**, *8*, 294–311.
9. Dubey, R.; Mishra, V.N. Symmetric duality results for a nondifferentiable multiobjective programming problem with support function under strongly assumptions. *RAIRO-Operation Res.* **2019**, *53*, 539–558.
10. Dubey, R.; Mishra, L.N.; Ali, R. Special class of second-order non-differentiable symmetric duality problems with (G, α_f) -pseudobonvexity assumptions. *Mathematics* **2019**, *7*, 763.

11. Antczak, T. On G -invex multiobjective programming. *Part I. Optim. J. Glob. Optim.* **2009**, *43*, 97–109.
12. Kang, Y.M.; Kim, D.S.; Kim, M.H. Optimality conditions of G -type in locally Lipschitz multiobjective programming. *Viet. J. Math.* **2014**, *40*, 275–285.
13. Gao, X. Sufficiency in multiobjective programming under second-order $B - (p, r) - V$ -type Ifunctions. *J. Interdiscip. Math.* **2014**, *17*, 385–402.
14. Brumelle, S. Duality for multiple objective convex programs. *Math. Oper. Res.* **1981**, *6*, 159–172.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).