

Document downloaded from:

<http://hdl.handle.net/10251/141653>

This paper must be cited as:

Llorens Rodríguez, R.; Noé, E.; Alcañiz Raya, ML.; Deutsch, JE. (12-2). Time since injury limits but does not prevent improvement and maintenance of gains in balance in chronic stroke. *Brain Injury*. 32(3):303-309. <https://doi.org/10.1080/02699052.2017.1418905>



The final publication is available at

<https://doi.org/10.1080/02699052.2017.1418905>

Copyright Taylor & Francis

Additional Information

Time since injury limits but does not prevent improvement and maintenance of gains in balance in chronic stroke

Roberto Llorens, PhD^{1,2}, Enrique Noé, MD PhD², Mariano Alcañiz¹, PhD, Judith E.
Deutsch, PhD.³

¹Neurorehabilitation and Brain Research Group, Instituto de Investigación e Innovación
en Bioingeniería, Universitat Politècnica de València, Camino de Vera s/n, 46022
Valencia, Spain.

²Servicio de Neurorehabilitación y Daño Cerebral de los Hospitales NISA, Fundación
Hospitales NISA, Río Tajo 1, 46011 Valencia, Spain.

³Rivers Lab, Department of Movement and Rehabilitation Sciences, Rutgers University-
School of Professions, 65 Bergen Street, Newark, USA.

Corresponding author

Name: Roberto Llorens

Email: rllorens@i3b.upv.es

Tel: +34963877518

Running head: Effect of chronicity on rehabilitation of balance

Keywords: stroke, time since injury, chronicity, posturography, balance, virtual reality.

Abstract

Objective

To determine the influence of time since injury on the efficacy and maintenance of gains of rehabilitation of balance after stroke.

Method

Forty-seven participants were assigned to a least (6-12 months), a moderate (12-24 months), or a most chronic (>24 months) group. Participants trained for 20 one-hour sessions, administered three to five times a week, combining conventional physical therapy and visual feedback-based exercises that trained the ankle and hip strategies. Participants were assessed before, after the intervention, and one month later with a posturography test (Sway Speed and Limits of Stability) and clinical scales.

Results

In contrast to other subjects, the most chronic participants failed to improve their sway and to maintain the benefits detected in the Limits of Stability after the intervention. Although all the participants improved in those clinical tests that better matched the trained skills, time since injury limited the improvement, and over all, the maintenance of gains.

Conclusion

Time since injury limits but does not prevent improvement in chronic stages post-stroke, and this effect appears to be more pronounced with maintaining gains. These findings support that training duration and intensity as well as type of therapy may need to be adjusted based on time post-stroke.

Keywords

Stroke; Chronic Brain Injury; Postural Balance; Posture; Rehabilitation; Virtual Reality
Therapy.

Introduction

In the last decade, vast resources have been invested to understand brain mechanisms after stroke. Current knowledge on pathophysiology has formed the basis of evidence-based neurorehabilitation.¹ The existing evidence supports that endogenous neurological recovery occurs with peak intensity within the first three months post-stroke and then declines and ends at six months.² Traditionally, the idea of recovery plateau has strongly influenced the way that rehabilitation has been administered.³ There is a general agreement among clinicians that the earlier the rehabilitation starts, the greater the benefits that patients can obtain.² Efficacy of early interventions has been supported by longitudinal studies that demonstrate greater improvement at earlier stages post-stroke.⁴ In addition, different attempts to determine the ideal timing for rehabilitation have shown that early interventions may provide greater benefits to functional independence for persons post-stroke in the acute⁵ and sub-acute phase.⁶

However, the boundaries of the window of opportunity for rehabilitation is still a matter of debate with clinical, economical, and ethical consequences.³ Recent literature has questioned the existence of a plateau, evidencing instead the lifelong potential of the neurobiological systems to adapt and reorganize. In accordance with this, it has been shown that functional improvements still occur after six months, supported by brain mechanisms that, although possible, must be externally driven.⁷ Functional and brain recovery is well documented after rehabilitation of upper limb function in the chronic phase.^{8,9} Different interventions have also shown to improve balance and postural control at this phase. One-on-one strength training interventions have reported benefits to individuals post-stroke more than one year after the onset.¹⁰ Group programs have also provided similar results after more than one¹¹ and two years.¹² Task-oriented

training involving reaches, steps, and raises has been reported to improve balance after one year since injury.¹³ Similar intense mobility training focusing on obstacle avoidance, walking, sit-to-stand, stair-climbing, and coordination tasks has also provided benefits to balance six¹⁴ and nine months after the onset.¹⁵ Provision of visual feedback on force plate has been also reported to promote benefits not only six months¹⁶ but also years after stroke.¹⁷ In addition virtual reality-based training of the ankle and hip strategies^{18,19} and the stepping strategies^{20,21} at this stage have been also shown to provide benefits in balance to individuals with stroke.

All these results support the evidence that, first, greater improvements are expected in early stages, but, second, rehabilitation of balance is possible in chronic stages. Unfortunately, differences in the intervention, dosage, and participants of the previous studies prevent accurate characterization of the effect of time since injury in the rehabilitation of balance in this phase.^{22,23} Characterization of this effect would help to better define rehabilitation interventions, over all, in the chronic stages.

According to the existing evidence, we hypothesized that time since injury would limit, but not nullify, the effect of an intensive training on balance function and postural control and the maintenance of gains after the intervention. The objective of this study was to determine the efficacy of a visual feedback-based intervention in groups of chronic post-stroke subjects with different time since injury. The intervention focused on the training of the ankle and hip strategies using force plates, and was previously included in the standard practice of both units, motivated by the positive effects of this^{18,19} and similar interventions^{16,17} reported in chronic subjects with variable time since injury.

Materials and methods

Participants

All the stroke outpatients who were attending a rehabilitation program in two specialized neurorehabilitation centers were candidates to participate in this study. The inclusion criteria were: 1) age ≥ 50 and ≤ 65 years old; 2) time since onset $>$ six months; 3) time from admission $>$ three months; 4) Mini-Mental State Examination²⁴ score above 23; and 5) ability to walk 10 meters indoors with or without assistive devices or orthotics. Subjects were excluded if they presented: 1) inability to follow instructions defined by Mississippi Aphasia Screening Test²⁵ score below 45; 2) unilateral spatial neglect; 3) ataxia or any other cerebellar symptom; and 4) severe auditory or visual impairments.

The study was approved by the Institutional Review Board of both medical centers. Written consent was obtained from all of the subjects who satisfied the inclusion criteria and accepted to participate in the study.

Intervention Protocol

Immediately prior to the intervention all the participants were enrolled in a long-term neurorehabilitation program in one of both centers that included conventional physical therapy three to five days a week. Participants were assigned to one of three groups according whether their time since injury was less than one year, from one to two years, and greater than two years. These artificial cut-offs were chosen as being representative of the literature.^{10-14,16} The intervention involved 20 sessions, one-hour long, administered three to five days a week. Training sessions combined 20 minutes of conventional physical therapy with 40 minutes of specific training of the ankle and hip strategies through virtual reality-based customized exercises on a balance board,^{18,19} with two-minute breaks each ten minutes. After the intervention, participants returned to the conventional physical therapy program. The conventional balance training consisted of one-on-one exercises including: 1) static standing exercises in different positions

(Romberg position, tandem stance, single stance, etc.) using verbal, visual, and perceptual cues to increase weight bearing to the affected lower limb; 2) stepping tasks to increase weight transfer and improve the stepping strategy; 3) static and dynamic balance exercises including arm activities during functional tasks to improve balance self-confidence in daily activities; and 4) walking exercises under different conditions (obstacle course, indoor and outdoor walking, stair climbing, etc.). The virtual reality-based training of the ankle and hip strategies consisted of six different exercises that promoted the training of the ankle and hip strategies (See Supplemental File for further information). Exercises provided audiovisual information of the center of pressure and required participants to displace it towards different targets in a virtual environment through weight transferences and postural readjustments.

The difficulty of the conventional physical therapy, including that prior and subsequent to the intervention, and the customized training of the ankle and hip strategies were established every two weeks by an experienced physical therapist according to the particular needs, condition, and progress of each participant. The maximum excursion of the center of pressure in the training of the ankle and hip strategies was defined as the 80% of the limits of stability of each participant.

All the participants were assessed before and after the intervention, and one month after, with a posturography test that studied Sway Speed during the modified Clinical Test of Sensory Interaction on Balance and Limits of Stability,²⁶ and with a battery of clinical scales that included the Berg Balance Scale (BBS),²⁷ the Functional Reaches Test (FRT),²⁸ the 30-Second Sit-to-Stand Test (30CST),²⁹ the Stepping Test in the paretic and non-paretic side,³⁰ the Timed Up-and-Down Stairs Test,³¹ the Timed Up-and-Go Test,³² and the 10-meter Walking Test.³³

Statistical Analysis

Demographical and clinical comparisons between groups at baseline were performed with analysis of variance (ANOVA) and Chi-squared or Fisher exact tests, as appropriate. Repeated-measures ANOVA with time as the within-subjects factor and chronicity as between-subjects factor were performed for all the tests and scales. The main effects of time, chronicity, and the time-chronicity interaction effects were evaluated. ANOVA findings that violated the sphericity assumption were accommodated by Greenhouse and Geisser's conservative degrees of freedom adjustment. For each repeated-measures ANOVA, we present the partial eta squared (η^2_p) as a measure of effect size; values may range between 0 and 1, with higher values representing higher proportions of variance explained by the independent variable.

The α level was set at 0.05 for all analyses (two-sided). All analyses were computed with SPSS Statistics, version 22 (IBM®, Armonk, NY, USA).

Results

Participants

A total pool of 167 outpatients were screened. Of those, 51 subjects (30.5%) met inclusion criteria and agreed to participate in the study. As explained above, participants were classified according to their time since stroke. Two participants of the moderately chronic group and one of the most chronic group were discharged and one participant of the least chronic group suffered a second stroke and their data were, consequently, excluded. Final analysis included data of 47 participants (Table 1). No significant differences in demographical or clinical data at inclusion were detected between the groups.

Posturographic Assessment

Different responses of the groups were detected in the Sway Speed ($p=0.030$, $\eta^2_p=0.11$). The least and moderately chronic participants similarly reduced their sway after the intervention, and continued improving at the follow-up examination. However, no change in this variable was detected in subjects with greatest chronicity. The differences between groups were more pronounced for the Limits of Stability. All the participants increased their limits after the intervention but only the least chronic subjects continued improving at follow-up, while the moderately chronic subjects maintained the gains, and the most chronic subjects failed to retain those gains ($p=0.000$, $\eta^2_p=0.12$). Posturographic data are shown in Table 2 and depicted in Figure 1.

Clinical Scales and Tests

Significant differences between the three groups were detected in the BBS ($p=0.005$, $\eta^2_p=0.15$), the FRT ($p=0.044$, $\eta^2_p=0.12$), and the 30CST ($p=0.017$, $\eta^2_p=0.10$). All the participants improved their scores in these tests after the intervention, even though more limited changes were reached by the most chronic group. Examination at follow-up evidenced that only the least chronic subjects continued improving in the three tests, while the moderately chronic subjects improved in the BBS and the FRT and maintained the gains in the 30CST, and the most chronic subjects maintained the gains in the BBS, but slightly worsened their scores in the FRT and 30CST. Clinical data are shown in Table 3 and significant measures are depicted in Figure 2.

Discussion

This paper reports on the efficacy of a visual feedback-based intervention that focused on postural adjustments through the use of the ankle and hip strategies in subjects post-stroke with different times since injury who were enrolled in a conventional physical therapy program immediately prior to the study. Our results support that this

intervention improved balance and postural control for persons at the chronic stages post-stroke, and more importantly, that time since injury modulated improvement as well as the ability to maintain gains.

Time Since Injury does Not Prevent Improvement

The improvement in posturographic data after the intervention in all groups, evidenced by increased Limits of Stability and reduced Sway Speed, supports previous reports in chronic populations.¹⁶ Even though a decrease in postural sway does not necessarily reflect improvement of the balance ability,³⁴ results in the clinical variables confirmed this relationship in our study. As proof, our data showed improvement in all groups in the BBS, the FRT, and the 30CST, like preliminary studies with similar interventions.^{18,19}

It is important to highlight that the significant clinical changes were detected in those tests that involved bilateral symmetrical lower extremity loading and consequently better match the trained skills. No significant changes were detected in more dynamic and gait tests, which may be explained by specificity of the training. However, this hypothesis should be further studied. Our results are in agreement with previous evidence¹⁰⁻²¹ that documents that it is possible to improve balance and postural control months and even years after the onset.

Time Since Injury Limits the Improvement

The different responses to training of the three groups in the sway speed indicated a negative effect of time since injury on the efficacy of the intervention in postural control. This effect was not reflected in the Limits of Stability, where the least chronic group showed the lowest increase. This finding might be interpreted as a ceiling effect on this measure, since this group had a mean baseline score of 92.8, more than 10 points higher than the other groups.

Results in the clinical outcomes are in line with the existing evidence. Greater improvement has been reported in functional independence in individuals with less time since injury,^{5,6} although the effect of this variable in the efficacy of an intervention could be less evident in the acute phase³⁵ and more pronounced over time. This relationship has been also confirmed in the motor function of upper extremities, where greater motor improvement has been detected at more acute stages³⁶ in comparison to that at chronic stages³⁷ after similar interventions. Our results support previous evidence that endorses greater improvement at early stages after stroke.

Time Since Injury Limits the Maintenance of Gains

The responses of the participants of the least chronic groups confirm previous findings after a similar intervention, where participants with similar characteristics were able to maintain the gains and even improve after the intervention.¹⁹ The return to the conventional physical therapy program after the intervention could have provided these participants with a sufficient dose and intensity of rehabilitation to maintain the benefits and even to be a solid background for further improvements. In contrast, the change of intervention could have interrupted the progress of the most chronic participants, who might need more time to show comparable improvements and to maintain part of the benefits obtained during the intervention. This effect could be especially detrimental in those skills that better matched the training, as reflected by the difficulty to maintain the gains in the Limits of Stability and the FRT, both tests require specific control of the ankle and hip strategies. A large number of studies have reported the progress of individuals post-stroke months and even years after the onset. Although some contradictory reports exist, a great body of research reveals that clinical deterioration is common after discharge of rehabilitation, not only with regards to dependency,³⁸ but also to function.³⁹ However, literature about the maintenance of gains when returning to

a physical therapy program after a specific intervention is more limited, especially in virtual reality-based interventions, where few empirical studies exist.^{21,40} Our results might reflect an overall increased difficulty in maintaining specific body function and structural improvements in the very chronic stages, while evidencing a preserved capability to retain the improvement in global balance.

Limitations

First, although the number of participants is comparable and even higher to similar interventions, it can be considered small. However, the similarity of the three groups on other variables, both demographic or clinical, support that time since injury could be a determining factor for improvement and maintenance of gains. Second, the specificity of the visual feedback-based intervention could restrict extrapolation to the results to other interventions. Third, follow-up assessment was at one month after the intervention making long-term effects of the intervention unknown. Finally, the design of our study precludes specific identification of the source of improvement. Consequently, it is not possible to discern whether the improvement was caused by the intervention itself, by the change of intervention, or by both.

Clinical Implications

The results of this study may have several implications for the clinical practice. First, the data provide evidence that persons post-stroke are able to improve their balance even at chronic stages. In line with previous work, this questions the existence of a ceiling effect in the rehabilitation of functional deficits after stroke.³ As a great percentage of stroke survivors will present motor deficits throughout their lifetime, and reduced balance is common following stroke, negatively impacting independence and safety,⁴¹ the findings presented here raise the important question of when patients should be discharged from neurorehabilitation programs. Our results presented as well

as previous literature,¹⁰⁻²¹ argue that a serious proportion of the discharges are in response to economic factors or resources rather than to rehabilitation criteria, resulting in some individuals not be receiving enough rehabilitation to reach their full recovery potential.⁴²

Second, as persons in the most chronic phase post-stroke did not maintain the gains from the program, one might speculate that they may need training of greater duration or greater dose than those who are less chronic. Third, although it is not possible to determine if the change from conventional to a visual feedback-based program was the source of improvement in our study, the change of intervention has been suggested to promote motor improvement in individuals who had plateaued.³ Different studies have corroborated the efficacy of novel rehabilitation interventions in this condition.^{43,44} Our results support the temporary use of intensive virtual reality-based balance training to overcome the plateau in the motor recovery promoted by conventional physical therapy in chronic individuals.

Finally, our results suggest that the chronic state post-stroke should be understood as a continuum rather than a single unchanging phase. Research should focus on scrutinizing the range of time post-stroke as there may be groups of chronic subjects (as was the case in this study) that respond differently.

Conclusions

Our results suggest that time since injury restricts, but does not prevent, the efficacy of physical interventions on balance and postural control and, over all, the maintenance of functional improvements in specific skills. However, despite these limitations, the improvement experienced by all the participants and the maintenance of gains in the global balance condition support the inclusion of individuals with chronic stroke in balance rehabilitation programs.

Acknowledgements

The authors wish to thank the staff and patients of the Hospital NISA Valencia al Mar (Valencia, Spain) and Instituto Valenciano de Acción Social (Valencia, Spain). This work was supported by Universitat Politècnica de València (Grant PAID-10-16).

Declaration of interest

The authors report no declarations of interest

References

1. Pekna M, Pekny M, Nilsson M. Modulation of neural plasticity as a basis for stroke rehabilitation. *Stroke* 2012;43(10):2819-2828.
2. Teasell R, Hussein N. General concepts: Therapies for rehabilitation and recovery. In: Ovbiagele B, editor. *Ischemic stroke therapeutics: A comprehensive guide*. Cham: Springer International Publishing; 2016. p 195-201.
3. Page SJ, Gater DR, Bach-y-Rita P. Reconsidering the motor recovery plateau in stroke rehabilitation. *Archives of physical medicine and rehabilitation* 2004;85(8):1377-1381.
4. Balasch IBM, Balasch IPS, Noe Sebastian E, Duenas Moscardo L, Ferri Campos J, Lopez-Bueno L. Study of the recovery patterns of elderly subacute stroke patients in an interdisciplinary neurorehabilitation unit. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association* 2015;24(10):2213-8.
5. Maulden SA, Gassaway J, Horn SD, Smout RJ, DeJong G. Timing of initiation of rehabilitation after stroke. *Archives of physical medicine and rehabilitation* 2005;86(12 Suppl 2):S34-S40.
6. Paolucci S, Antonucci G, Grasso MG, Morelli D, Troisi E, Coiro P, Bragoni M. Early versus delayed inpatient stroke rehabilitation: A matched comparison conducted in Italy. *Archives of physical medicine and rehabilitation* 2000;81(6):695-700.
7. Cassidy JM, Cramer SC. Spontaneous and therapeutic-induced mechanisms of functional recovery after stroke. *Translational stroke research* 2016.

8. Gauthier LV, Taub E, Perkins C, Ortmann M, Mark VW, Uswatte G. Remodeling the brain: Plastic structural brain changes produced by different motor therapies after stroke. *Stroke* 2008;39(5):1520-5.
9. Yin D, Luo Y, Song F, Xu D, Peterson BS, Sun L, Men W, Yan X, Fan M. Functional reorganization associated with outcome in hand function after stroke revealed by regional homogeneity. *Neuroradiology* 2013;55(6):761-70.
10. Weiss A, Suzuki T, Bean J, Fielding RA. High intensity strength training improves strength and functional performance after stroke. *American journal of physical medicine & rehabilitation / Association of Academic Physiatrists* 2000;79(4):369-76; quiz 391-4.
11. Eng JJ, Chu KS, Kim CM, Dawson AS, Carswell A, Hepburn KE. A community-based group exercise program for persons with chronic stroke. *Medicine and science in sports and exercise* 2003;35(8):1271-8.
12. Mount J, Bolton M, Cesari M, Guzzardo K, Tarsi J, Jr. Group balance skills class for people with chronic stroke: A case series. *Journal of neurologic physical therapy : JNPT* 2005;29(1):24-33.
13. Yang YR, Wang RY, Lin KH, Chu MY, Chan RC. Task-oriented progressive resistance strength training improves muscle strength and functional performance in individuals with stroke. *Clinical rehabilitation* 2006;20(10):860-70.
14. Fritz SL, Pittman AL, Robinson AC, Orton SC, Rivers ED. An intense intervention for improving gait, balance, and mobility for individuals with chronic stroke: A pilot study. *Journal of neurologic physical therapy : JNPT* 2007;31(2):71-6.

15. Macko RF, Benvenuti F, Stanhope S, Macellari V, Taviani A, Nesi B, Weinrich M, Stuart M. Adaptive physical activity improves mobility function and quality of life in chronic hemiparesis. *Journal of rehabilitation research and development* 2008;45(2):323-8.
16. Lee SW, Shin DC, Song CH. The effects of visual feedback training on sitting balance ability and visual perception of patients with chronic stroke. *Journal of physical therapy science* 2013;25(5):635-9.
17. Tsaklis PV, Grooten WJ, Franzen E. Effects of weight-shift training on balance control and weight distribution in chronic stroke: A pilot study. *Topics in stroke rehabilitation* 2012;19(1):23-31.
18. Gil-Gomez JA, Llorens R, Alcaniz M, Colomer C. Effectiveness of a wii balance board-based system (ebavir) for balance rehabilitation: A pilot randomized clinical trial in patients with acquired brain injury. *Journal of neuroengineering and rehabilitation* 2011;8:30.
19. Llorens R, Albiol S, Gil-Gómez J-A, Alcañiz M, Colomer C, Noé E. Balance rehabilitation using custom-made wii balance board exercises: Clinical effectiveness and maintenance of gains in an acquired brain injury population. *Int J Disabil Hum Dev* 2014;13(3):327-332.
20. Llorens R, Gil-Gomez JA, Alcaniz M, Colomer C, Noe E. Improvement in balance using a virtual reality-based stepping exercise: A randomized controlled trial involving individuals with chronic stroke. *Clinical rehabilitation* 2015;29(3):261-8.
21. Llorens R, Noe E, Colomer C, Alcaniz M. Effectiveness, usability, and cost-benefit of a virtual reality-based telerehabilitation program for balance recovery

- after stroke: A randomized controlled trial. *Archives of physical medicine and rehabilitation* 2015;96(3):418-425 e2.
22. Dromerick AW, Edwardson MA, Edwards DF, Giannetti ML, Barth J, Brady KP, Chan E, Tan MT, Tamboli I, Chia R and others. Critical periods after stroke study: Translating animal stroke recovery experiments into a clinical trial. *Frontiers in human neuroscience* 2015;9:231.
 23. Geiger RA, Allen JB, O'Keefe J, Hicks RR. Balance and mobility following stroke: Effects of physical therapy interventions with and without biofeedback/forceplate training. *Phys Ther* 2001;81(4):995-1005.
 24. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research* 1975;12(3):189-98.
 25. Romero M, Sanchez A, Marin C, Navarro MD, Ferri J, Noe E. Clinical usefulness of the spanish version of the mississippi aphasia screening test (mastsp): Validation in stroke patients. *Neurologia* 2012;27(4):216-24.
 26. Llorens R, Latorre J, Noe E, Keshner EA. Posturography using the wii balance board: A feasibility study with healthy adults and adults post-stroke. *Gait & posture* 2016;43:228-32.
 27. Berg K, Wood-Dauphinee S, Williams JI. The balance scale: Reliability assessment with elderly residents and patients with an acute stroke. *Scandinavian journal of rehabilitation medicine* 1995;27(1):27-36.
 28. Duncan PW, Weiner DK, Chandler J, Studenski S. Functional reach: A new clinical measure of balance. *Journal of gerontology* 1990;45(6):M192-7.

29. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exercise Sport* 1999;70(2):113-119.
30. Hill KD, Bernhardt J, McGann AM, Maltese D, Berkovits D. A new test of dynamic standing balance for stroke patients: Reliability, validity and comparison with healthy elderly. *Physiother Can* 1996;48(4):257-262.
31. Zaino CA, Marchese VG, Westcott SL. Timed up and down stairs test: Preliminary reliability and validity of a new measure of functional mobility. *Pediatric physical therapy : the official publication of the Section on Pediatrics of the American Physical Therapy Association* 2004;16(2):90-8.
32. Podsiadlo D, Richardson S. The timed "up & go": A test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society* 1991;39(2):142-8.
33. Bohannon RW, Andrews AW, Thomas MW. Walking speed: Reference values and correlates for older adults. *The Journal of orthopaedic and sports physical therapy* 1996;24(2):86-90.
34. Cho K, Lee K, Lee B, Lee H, Lee W. Relationship between postural sway and dynamic balance in stroke patients. *Journal of physical therapy science* 2014;26(12):1989-1992.
35. Gagnon D, Nadeau S, Tam V. Ideal timing to transfer from an acute care hospital to an interdisciplinary inpatient rehabilitation program following a stroke: An exploratory study. *BMC health services research* 2006;6:151.
36. Myint JM, Yuen GF, Yu TK, Kng CP, Wong AM, Chow KK, Li HC, Chun Por W. A study of constraint-induced movement therapy in subacute stroke patients in hong kong. *Clinical rehabilitation* 2008;22(2):112-24.

37. Page SJ, Sisto S, Levine P, McGrath RE. Efficacy of modified constraint-induced movement therapy in chronic stroke: A single-blinded randomized controlled trial. *Archives of physical medicine and rehabilitation* 2004;85(1):14-8.
38. Ullberg T, Zia E, Petersson J, Norrving B. Changes in functional outcome over the first year after stroke: An observational study from the swedish stroke register. *Stroke* 2015;46(2):389-94.
39. Clanchy KM, Tweedy SM, Trost SG. Evaluation of a physical activity intervention for adults with brain impairment: A controlled clinical trial. *Neurorehabilitation and neural repair* 2016.
40. Colomer C, Llorens R, Noe E, Alcaniz M. Effect of a mixed reality-based intervention on arm, hand, and finger function on chronic stroke. *Journal of neuroengineering and rehabilitation* 2016;13(1):45.
41. Geurts AC, de Haart M, van Nes IJ, Duysens J. A review of standing balance recovery from stroke. *Gait & posture* 2005;22(3):267-81.
42. Lang CE, MacDonald JR, Gnip C. Counting repetitions: An observational study of outpatient therapy for people with hemiparesis post-stroke. *Journal of neurologic physical therapy : JNPT* 2007;31(1):3-10.
43. Whitall J, McCombe Waller S, Silver KH, Macko RF. Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. *Stroke* 2000;31(10):2390-5.
44. Hesse S, Schulte-Tigges G, Konrad M, Bardeleben A, Werner C. Robot-assisted arm trainer for the passive and active practice of bilateral forearm and wrist movements in hemiparetic subjects. *Archives of physical medicine and rehabilitation* 2003;84(6):915-20.

Tables

Table 1. Characteristics of the Participants.

Age, chronicity, and scores in the Berg Balance Scale are defined in terms of mean and standard deviation. Gender, etiology, and hemiparesis are expressed as a number and percentage of total number of participants. TACI: total anterior circulation infarct; PACI: partial anterior circulation infarct; LACI: lacunar infarct; POCI: posterior circulation infarct; NS: non-significant.

Table 2. Posturographic Data.

Data are given in terms of mean and standard deviation. T: time effect. GxT: group by time effect. *: $p < 0.05$, **: $p < 0.01$. NS: non-significant.

Table 3. Clinical Data.

Data are given in terms of mean and standard deviation. T: time effect. GxT: group by time effect. *: $p < 0.05$, **: $p < 0.01$. NS: non-significant.

Figures

Figure 1. Progress on Posturographic Data

Statistical significance is provided for each group in those time intervals with a significant time effect. The significant time effects and group-by-time effects detected in each time interval and their effect size are provided beneath them.

Figure 2. Progress on Significant Clinical Data

Statistical significance is provided for each group in those time intervals with a significant time effect. The significant time effects and group-by-time effects detected in each time interval and their effect size are provided beneath them.