

Document downloaded from:

<http://hdl.handle.net/10251/147781>

This paper must be cited as:

Manzo, E.; Ciavatta, ML.; López-Gresa, MP.; Gavagnin, M.; Villani, G.; Naik, CG.; Cimino, G. (2007). New bioactive hydrogenated linderazulene-derivatives from the gorgonian *Echinogorgia complexa*. *Tetrahedron Letters*. 48(14):2569-2571.
<https://doi.org/10.1016/j.tetlet.2007.02.020>



The final publication is available at

<https://doi.org/10.1016/j.tetlet.2007.02.020>

Copyright Elsevier

Additional Information

New Bioactive Hydrogenated Linderazulene-Derivatives from the Gorgonian *Echinogorgia complexa*

Emiliano Manzo,* Maria Letizia Ciavatta, Maria Pilar Lopez Gresa, Margherita Gavagnin, Guido Villani, Chandrakant Govind Naik and Guido Cimino



Iso-echinofuran (**3**) and 8,9-dihydro-linderazulene (**4**) are inhibitors of mitochondrial respiratory chain.

New Bioactive Hydrogenated Linderazulene-Derivatives from the Gorgonian *Echinogorgia complexa*

Emiliano Manzo,^{a*} Maria Letizia Ciavatta,^a Maria Pilar Lopez Gresa,^b Margherita Gavagnin,^a Guido Villani,^a Chandrakant Govind Naik^c and Guido Cimino^a

^a Istituto di Chimica Biomolecolare, CNR, Via Campi Flegrei 34, I 80078-Pozzuoli (Naples), Italy

^b Centro Ecologia Química Agrícola, Universidad Politécnica Valencia, Campus de Vera, Valencia, Spain

^c National Institute of Oceanography, CSIR, Dona Paula, 403 004 Goa, India

Abstract: Chemical analysis of the secondary metabolite pattern of the gorgonian *Echinogorgia complexa*, collected along South Indian coasts, showed the presence of two new tricyclic guaiane furanosesquiterpenes **3** and **4** co-occurring with the known echinofuran **5**, which are structurally related to the pigment linderazulene **1**. The unprecedented molecules **3** and **4** were characterized by spectral methods, mainly by NMR techniques. Compounds **3-5** displayed moderate activity in the mitochondrial respiratory chain inhibition assay.

Keywords: gorgonian, pigments, linderazulene, echinofuran, furanosesquiterpenes, MRC.

*Author to whom correspondence should be addressed; E-mail: emanzo@icmib.na.cnr.it

Azulene sesquiterpenes are typical secondary metabolites of marine octocorals of the order *Gorgonacea*.¹⁻¹¹ Most of them are responsible of the brilliant colors of gorgonians and in addition various bioactivities including cytotoxic, antimicrobial, antifungal, immunostimulatory and cell division inhibitory properties have been reported for azulene sesquiterpenes.^{7,8,10} The two members of this class that have been isolated from marine organisms are linderazulene (**1**)^{1,3,5} and guaiazulene (**2**),^{2,3,10,12} previously reported from terrestrial sources.^{13,14} Subsequently, several related sesquiterpenes have been found in gorgonian species belonging to different genera (*Paramuricea*, *Acalycigorgia*, *Bebryce*, *Echinogorgia*, *Pseudopterogorgia*).^{3-11,15,16} Guaiane azulenes have been found to be quite labile and easily decomposable on exposure to air and light during the work-up. A photo artifact from linderazulene (**1**) has been also reported.¹⁷

In this paper we describe the isolation, structure elucidation and biological activity of two new hydrogenated linderazulene derivatives designated as tricyclic furanosesquiterpenes **3** and **4**, co-occurring with the known echinofuran **5**⁸ in the gorgonian *Echinogorgia complexa* Nutting collected from subtidal regions of West Indian coasts (Goa).

Although previous studies on *E. complexa* showed a strong antifouling activity of the alcoholic extract prepared from this species,^{18,19} but no active compounds were reported so far.

INSERT FORMULAE 1-6

A sample of the gorgonian *E. complexa* (dry weight 54 g) was collected in December 2005, immediately frozen at -20 °C and subsequently transferred to ICB for the chemical studies. The frozen animal was extracted exhaustively with acetone (4x200 mL). The Et₂O soluble

fraction of the acetone extract was concentrated under reduced pressure to furnish a residual orange oil (600 mg), which was analyzed by TLC chromatography (light petroleum ether/diethyl ether in different ratio) showing mainly the presence of sterols and lipids. The less polar part of the extract was found to contain three yellow spots between Rf 0.35-0.55 (*n*-hexane). The extract was then purified on a silica-gel column (petroleum light ether with increasing amount of diethyl ether) to obtain the mixture of yellow components (18 mg). A preliminary NMR analysis of this mixture suggested the presence of furano-sesquiterpenes, and subsequent comparison with available literature data indicated that these three compounds were related to linderazulene type of structural framework. This fraction was further purified on a semi-preparative silica-gel plate, using *n*-hexane as eluent, to afford echinofuran (**5**, 1.0 mg) and two novel compounds named as *iso*-echinofuran (**3**, 0.7 mg) and 8,9-dihydro-linderazulene (**4**, 0.7 mg) in order of increasing polarity.

(-)-Echinofuran (**5**) was identified by comparison of ¹H-NMR, mass spectral data and [α]_D²⁵ with the literature values.^{8,20} The ¹³C-NMR values of **5**, not previously assigned, were also assigned by performing 2D-NMR experiments.²¹ Due to high susceptibility of these molecules in chloroform, NMR analysis of **5** was carried out in deuterated benzene. Compounds **3** and **4**, structurally related to **5**, were also analyzed in the same manner.

Compound **3**²² had a molecular formula C₁₅H₁₈O, which was determined by HR-APCIMS giving a pseudo-molecular ion [M+H]⁺ *m/z* 215.1431, and ¹³C-NMR spectrum (Table 1). The ¹H-NMR spectrum of **3** showed strong analogies with that of echinofuran (**5**), differing only from this are the presence of two additional signals, an aliphatic methine proton (δ 2.35, m, H-7a) and an olefinic proton (δ 6.33, bs, H-9), which replaced the methylene group H₂-9 resonating at δ 3.59 (1H, bd, *J*=17.8 Hz, H-9) and 3.02 (1H, d, *J*=17.8 Hz, H-9) in **5**. The ¹³C-NMR spectrum showed signals attributable to seven sp³ carbons (two vinyl methyls at δ 7.8 and δ 22.2; three methylenes at δ 28.6, 29.8 and 32.0; two methine carbons at δ 50.9 and 45.9) and eight sp² carbons attributable to four double bonds: two tri-substituted (δ 136.0, 121.8, 137.3 and 116.8), a tetra-substituted (δ 149.0 and 120.0), and an exocyclic double bond (δ 105.6 and 156.1). These data indicated that compound **3** was an isomer of **5**, with a double bond at C-8/C-9 position instead of a double bond at C-7a/C-8. The relative stereochemistry of A,B-ring junction of **3** was suggested to be *cis*, the same as the related bebryazulene (**6**) by comparing the carbon values of 5-membered ring of both compounds,¹¹ that were similar. Unfortunately, **3** degraded before recording nOe difference experiments, which had been planned to confirm this suggestion. The proton and carbon values of **3** were assigned by 2D-NMR experiments (Table 1).

INSERT TABLE 1

The molecular formula C₁₅H₁₆O of compound **4**²³ was determined by HR-APCIMS which gave a pseudo-molecular ion [M+H]⁺ *m/z* 213.1520 and ¹³C-NMR values (Table 1). The ¹H-NMR spectrum displayed olefinic signals at δ 6.75 (1H, s, H-4) and 6.02 (2H, bs, H-6 and H-7), a singlet at δ 6.77 due an α-furan proton (H-2), two double doublets at δ 2.82 (1H, dd, *J*=16.4, 2.9 Hz, H-9) and 2.66 (1H, dd, *J*=16.4, 9.4 Hz, H-9) attributed to a methylene coupled

to a methine at δ 2.83 (1H, m, H-8), which was further correlated to a secondary methyl signal resonating at δ 1.08 (3H, d, $J=6.8$ Hz, H₃-12), and finally two methyl broad singlets occurring at δ 1.73 (H₃-10) and 2.06 (H₃-11). The ¹³C-NMR spectrum contained signals due to ten sp² carbons according to the presence of an additional unsaturation with respect to compound **3** (Table 1). Careful comparison of spectral data of **4** with those of linderazulene-related compounds^{1,5,7,8,11} suggested that it was a dihydro-linderazulene. A series of 2D-NMR experiments (¹H-¹H COSY, HMQC, HMBC) led to the assignment of structure **4** and fully satisfying with assigned values of proton and carbon signals (Table 1).

Interesting biological activities have been reported in the literature for several guaianoid azulene metabolites.^{7,8,10} In particular, echinofuran (**5**) was found to inhibit cell division of fertilized sea urchin eggs.⁸ Compounds **3-5**, isolated in the present study, have been tested in the mitochondrial respiratory chain assay. They displayed a moderate activity for the inhibition of the integrated electron transfer chain (NADH oxidase activity) in beef heart sub-mitochondrial particles (SMP).²⁴ IC₅₀ values were 4.3 ± 0.15 μ M, 2.5 ± 0.04 μ M and 2.2 ± 0.31 μ M for compounds **3**, **4** and **5** respectively, whereas full inhibition of rotenone sensitive NADH oxidase activity was achieved at approximately 2 μ M.²⁵

Acknowledgements

We thank ICB Mass Service and ICB NMR Service Centre (Mr. S. Zambardino is kindly acknowledged), Mr. C. Iodice for spectrophotometric measurements and Mr. R. Turco for graphical work. Biological assays were performed by Prof. Estornell D. and Dr. Romero V. that the authors kindly acknowledge. This work was partially supported by a bilateral CNR(Italy)-CSIR(India) programme. We are grateful to Dr. P.A.Thomas (Central Marine Fisheries Research Institute-CMFRI, Vizhijam, Kerala, India) for the identification of *E.complexa*.

References and notes

1. Imre, S.; Thomson, R.H.; Yalhi, B. *Experientia* **1981**, *37*, 442-443.
2. Fusetani, N.; Matsunaga, S.; Konosu, S. *Experientia* **1981**, *37*, 680-681.
3. Okuda, R. K.; Klein, D.; Kinnel, R. B.; Li, M.; Scheuer, P. J. *Pure Appl. Chem.* **1982**, *54*, 1907-1914.
4. Li, M. K. W.; Scheuer, P. J. *Tetrahedron Lett.* **1984**, *25*, 587-590.
5. Li, M. K. W.; Scheuer, P. J. *Tetrahedron Lett.* **1984**, *25*, 2109-2110.
6. Li, M. K. W.; Scheuer, P. J. *Tetrahedron Lett.* **1984**, *25*, 4707-4708.
7. Sakemi, S.; Higa, T. *Experientia* **1987**, *43*, 624-625.
8. Tanaka, J.I.; Miki, H.; Higa, T. *J. Nat. Prod.* **1992**, *55*, 1522-1524.
9. Ochi, M.; Kataoka, K.; Tatsukawa, A.; Kotsuhi, H.; Shibata, K. *Chem. Lett.* **1993**, 2003-2006.
10. Seo, Y.; Rho, J.-R.; Geum, N.; Yoon, J. B.; Shin, J. *J. Nat. Prod.* **1996**, *59*, 985-986.
11. Aknin, M.; Rudi, A.; Kashman Y.; Gaydou, E. M. *J. Nat. Prod.* **1998**, *61*, 1286-1287.
12. Howard, B. M.; Fenical, W. *Tetrahedron Lett.* **1978**, 2453-2456.
13. Haagen-Smit, A. J. *Fortschr. Chem. Org. Naturst.* **1948**, *5*, 40-71.
14. Nozoe, T.; Ito, S. *Fortschr. Chem. Org. Naturst.* **1961**, *19*, 32-119.
15. Rodriguez, A.D.; Boulanger, A. *J. Nat. Prod.* **1996**, *59*, 653-657.
16. Rodriguez, A.D.; Boulanger, A. *J. Nat. Prod.* **1997**, *60*, 207-211.
17. Alpertunga, B.; Imre, S.; Cowe, H.; Cox, P. J.; Thomson, R. H. *Tetrahedron Lett.* **1983**, *24*, 4461-4462.
18. Wilsanand, V.; Wagh, A.B.; Bapuji, M. *Indian J. Mar. Sci.* **1999**, *28*, 280-284.
19. Wilsanand, V.; Wagh, A.B.; Bapuji, M. *Microbios* **2001**, *104*, 131-140.
20. Due to the easy degradation of compound **5** it is conceivable that the optical rotation value may be inaccurate. ¹H-NMR spectrum was also recorded in CDCl₃ as reported in ref. 8.
21. *Compound 5*: R_f = 0.50 (*n*-hexane); [α]_D²⁵ = -14° (*c* = 1.0 mg, *n*-hexane); lit. [8] [α]_D²⁰ = -91° (CHCl₃); APCIMS: *m/z* 215.1440 (calculated for C₁₅H₁₈O⁺: 215.1436); ¹H-NMR (400M Hz, C₆D₆): δ 6.93 (1H, bs, H-2), δ 4.98 (1H, bs, H-11a), δ 4.90 (1H, bs, H-11b), δ 3.59 (1H, bd, $J=17.8$ Hz, H-9), δ 3.26 (1H, m, H-4), δ 3.02 (1H, d, $J=17.8$ Hz, H-9), δ 2.52 (1H, dt, $J=3.42, 3.42, 15.7$ Hz, H-7a), δ 2.38 (2H, m, H-6), δ 2.28 (2H, m, H-4), δ 2.24 (1H, m, H-7b), δ 1.72 (3H,

bs, H₃-10), δ 1.52 (3H, bs, H₃-12). ¹³C-NMR (C₆D₆): δ 156.3 (C-5, C), δ 149.2 (C-9a, C), δ 140.5 (C-7a, C), δ 135.7 (C-2, CH), δ 124.5 (C-8, C), δ 121.1 (C-3, C), δ 119.3 (C-3a, C), δ 105.6 (C-11, CH₂), δ 46.0 (C-4a, CH), δ 33.7 (C-9, CH₂), δ 32.8 (C-4, CH₂), δ 30.2 (C-6, CH₂), δ 29.9 (C-7, CH₂), δ 21.3 (C-12, CH₃), δ 8.8 (C-10, CH₃).

22. **Compound 3**: $R_f = 0.45$ (*n*-hexane); $[\alpha]_D^{25} = -2.8^\circ$ ($c = 0.07$, *n*-hexane); APCIMS: m/z 215.1431 (calculated for C₁₅H₁₈OH⁺: 215.1436); ¹H- and ¹³C-NMR data are given in Table 1.

23. **Compound 4**: $R_f = 0.40$ (*n*-hexane); $[\alpha]_D^{25} = -38.2^\circ$ ($c = 0.07$, *n*-hexane); APCIMS: m/z 213.1520 (calculated for C₁₅H₁₆OH⁺: 213.1517); ¹H- and ¹³C-NMR data are given in Table 1.

24. Fato, R.; Estornell, D.; Bernardo, S.; Pallotti, F.; Parenti-Castelli, G.; Lenaz, G. *Biochemistry* **1996**, *35*, 2705-2716.

25. Fontana, A.; Cimino, G.; Gavagnin M.; Gonzalez M.C.; Estronell, E.; *J. Med. Chem.* **2001**, *44*, 2362-2365.

Table 1 NMR Spectroscopic Data^a (400 MHz, C₆D₆) for compound (**3**) and compound (**4**)

Position	3		4	
	δ_C mult.	δ_H (J in Hz)	δ_C mult.	δ_H (J in Hz)
2	136.0 CH	6.94 bs	137.3 CH	6.77 bs
3	121.8 qC	-	120.6 qC	-
3a	120.0 qC	-	119.5 qC	-
4	29.8 CH ₂	2.80 dd (16.4, 3.9) 2.39 dd (16.4, 3.1)	122.4 CH	6.75 s
4a	45.9 CH	2.56 m	133.2 qC	-
5	156.1 qC	-	144.1 qC	-
6	32.0 CH ₂	2.28 m 2.11 m	125.6 CH	-6.02 bs
7	28.6 CH ₂	1.67 m 1.32 m	125.6 CH	-6.02 bs
7a	50.9 CH	2.35 m	138.1 qC	-
8	137.3 qC	-	31.5 CH	2.83 m
9	116.8 CH	6.33 bs	34.1 CH ₂	2.82 dd (16.4, 2.9) 2.66 dd (16.4, 9.4)
9a	149.0 qC	-	158.1 qC	-
10	7.8 CH ₃	1.73 bs	7.4 CH ₃	1.73 bs
11	105.6 CH ₂	5.00 bs 4.91 bs	12.2 CH ₃	2.06 bs
12	22.2 CH ₃	1.67 bs	19.8 CH ₃	1.08 d (6.8)

^aAssignments aided by COSY, HSQC, HMBC

