

Assessing the effects of *in vitro* imposed water stress on pineapple growth in relation to biochemical stress indicators using polynomial regression analysis

Daviel GÓMEZ¹, Doris ESCALANTE², Elliasha HAJARI³,
Oscar VICENTE^{4*}, SERSHEN⁵, José C. LORENZO²

¹Universidad de Concepción, Facultad de Ingeniería, Departamento de Ingeniería Química, Laboratorio de CarboCat, Región del Bio-Bio, Chile; davielga26@gmail.com

²University of Ciego de Avila, Bioplant Center, Laboratory for Plant Breeding and Conservation of Genetic Resources, Ciego de Avila, 69450, Cuba; descalante@bioplantacuba.cu; jclorenzo@bioplantacuba.cu

³Plant Improvement; Agricultural Research Council-Tropical and Subtropical Crops; Private Bag X11208, Nelspruit, 1200, South Africa; HajariE@arc.agric.za

⁴Universitat Politècnica de València, Institute for the Preservation and Improvement of Valencian Agrobiodiversity (COMAV), Valencia, Spain; ovicente@upvnet.upv.es

⁵University of the Western Cape, Department for Biodiversity and Conservation Biology, Bellville, 7535, South Africa; sersbenn@gmail.com

Abstract

Knowing the mechanisms that operate under water stress in commercial crops, particularly those that can affect productivity, such as phenolic or cell wall metabolism, is becoming increasingly important in a scenario of global climate change. However, our understanding of how to analyse statistically the relationships between these commonly used biochemical markers of water stress and growth in crops like pineapple, needs to be improved. In the present work, we have addressed the question of whether polynomial regression analysis can be used to describe the influence of selected plant metabolites (chlorophylls, carotenoids, phenolics and aldehydes) on shoot biomass, in response to a mannitol-induced water stress in temporary immersion bioreactors (TIBs). Polynomial regression analysis has been applied to investigate plant stress responses in many species but is very seldom used in *in vitro* screening studies. Here, the relationship between biochemical indicators (x ; independent variable) and shoot growth (y ; dependent variable) has been characterised, with y modelled as an n^{th} degree polynomial in x . This statistical approach accommodated for the non-linear (curvilinear) relationships between variables, and the results showed that shoot biomass was negatively, and significantly correlated with soluble phenolics, cell wall-linked phenolics and other aldehydes (characterised by “High” R^2 values).

Keywords: *Ananas comosus* (L.) Merr.; biostatistics; drought; *in vitro* osmotic stress; mannitol; plant metabolites; temporary immersion bioreactors (TIBs)

Introduction

Statistical tools can outline trends in biological data sets. Regression analysis is used as a predictive tool to investigate the relationship between a dependent and an independent variable, with linear regressions being the most common type of analysis. However, when the collected data are non-linear (or curvilinear), polynomial regressions are performed. This is often the case for many biological processes in plants (see references below). For polynomial regressions, the relationship between the independent variable x and the dependent variable y is modelled as an n^{th} degree polynomial in x . In order to validate the strength of the model, the coefficient of determination is calculated. The coefficient of determination is the square of the correlation coefficient, also known as “R”. The R^2 values range from 0 (indicating a poor fit) to 1 (indicative of the best fit, or relationship, between the two factors) (Ivanov, 1989).

Polynomial regression models have a wide range of applications and have been used in engineering, computer science, human resource management, urban planning, health sciences, and many other fields (Ivanov, 1989). In the context of plant sciences, polynomial regression analysis has been extensively applied in studies investigating the response of plants to varied stress conditions. Fernández *et al.* (2002) used polynomial adjustment of log-transformed data to show that water stress altered the developmental response of two desert grass species, with leaf area ratio being the most prominent factor affected. Similarly, Vieira *et al.* (2019) used polynomial regression analysis in modelling the gas exchange behaviour of melon exposed to water stress. Wijewardana *et al.* (2019) showed that polynomial regressions could be employed to model the germination of soybean seeds exposed to water stress. A novel application was described by Runeckles (1982), where polynomial regressions were applied to define the relative death rate of plants exposed to stresses. Polynomial regressions were also used to investigate the effect of sub-acute high temperature on ovule and pollen development in tomato (Peet *et al.*, 1997), the effect of high temperature on photosynthetic processes in a forest ecosystem (Peet *et al.*, 1997; Hüve *et al.*, 2019), or to model the effect of excess ammonium fertiliser in tomato (Barker, 1999). However, to the best of our knowledge, there are no published reports on the use of polynomial regression analysis to identify suitable biochemical indicators of stress tolerance in *in vitro* screen systems such as Temporary Immersion Bioreactors (TIBs). Hence, the current study reports on the characterisation of pineapple plants responses to mannitol (to mimic water stress) in TIBs, in relation to established stress biomarkers, such as the levels of chlorophyll a and b, carotenoids, phenolics (soluble, cell wall-linked and exuded into the medium) and aldehydes, and the fresh mass of the shoots produced. The empirical data used for the analyses emanates from a previously published study by our group (Gómez *et al.*, 2017).

Materials and Methods

Plant material and in vitro culture conditions

Pineapple crowns were collected from plants grown in the field (cv. ‘MD2’) as per Daquinta and Benegas (1997). Upon receipt in the laboratory, the crown leaves were removed, and the stem was washed with detergent and tap water to remove soil and other particles. The crown stem was then decontaminated by immersion in 1% (w/v) calcium hypochlorite for 10 min. Axillary buds, including a portion of basal tissue, were excised and placed onto initiation medium. Explants were cultured in 300 ml glass containers containing 5 ml of liquid culture medium per explant. The culture medium was comprised of MS (Murashige and Skoog, 1962) salts, 100 mg l⁻¹ *myo*-inositol, 0.1 mg l⁻¹ thiamine-HCl, 30 g l⁻¹ sucrose, 4.4 µM 6-benzyladenine (BA), and 5.3 µM naphthaleneacetic acid (NAA). After 45 d of growth, shoots were subcultured to promote multiplication of explants, in the same medium described above, except that it contained 9.3 µM BA and 1.6 µM NAA. Shoots were multiplied for six months, with plants being transferred to fresh medium every 45 d. Once a sufficient number of shoots were obtained, they were placed in TIBs according to Escalona *et al.* (1999), supplemented

with 3.0 μM paclobutrazol. The TIBs were programmed to allow for immersion of plantlets into the culture medium for 2 min every 3 h for 30 d. Each TIB contained 200 ml of liquid medium with five explants per bioreactor.

Plants were exposed to varying levels of mannitol in TIBs (0, 50, 100, 150, and 200 mM). Each treatment was comprised of 3 bioreactors (5 explants per bioreactor). Cultures were maintained at $25 \pm 1^\circ\text{C}$; $80 \mu\text{mol m}^{-2} \text{s}^{-1}$ with fluorescent light and an 8-hour photoperiod.

Measurement of stress indicators

Following 30 days of culture, shoot cluster fresh weight, and the levels of different biochemical stress markers were determined in the plant material. Total chlorophylls and carotenoids were extracted from plants with 5 ml acetone (80%, v/v). The samples were then centrifuged (12,000 rpm at 4°C for 15 min). The absorbance of the supernatant was measured at 646.6 and 663.6 nm, and chlorophyll a and b contents were calculated as described in Porra (2002), whereas total carotenoid concentrations were determined from the absorbance of the samples at 470 nm, according to the method of Lichtenthaler (1987). A colourimetric method based on the Folin Ciocalteu reagent was used for extraction and quantification of phenolics (Gurr *et al.*, 1992), and the results were expressed as mg chlorogenic acid equivalents per g fresh weight. Malondialdehyde (MDA) and other aldehydes were quantified based on the reaction with thiobarbituric acid (Heath and Packer, 1968). Phenolics excreted into the medium were also determined, using a modification of the Hoagland (1990) procedure: 0.5 ml of culture medium was mixed with 4.5 ml of distilled water and 0.5 ml of Folin Ciocalteu reagent (50% v/v). The mixture was stirred, the reaction was allowed to proceed for 5 min and 1 ml of a saturated sodium carbonate solution was added. The mixture was stirred again, left for 60 min, and the optical density was measured at 725 nm. The concentration of phenolics was determined using a calibration curve with chlorogenic acid as the standard.

Data analysis

Polynomial regression analyses were performed using Microsoft Excel. Each polynomial regression involved 15 (x, y) pairs (i.e. 5 mannitol concentrations; 3 TIBs per treatment). Determination coefficients (R^2) obtained for $y = ax^2 + bx + c$ used the *shoot fresh weight* as the dependent variable (y), whereas biochemical indicators represented the independent variables (x). The R^2 values were classified as “*Low*” from 0.2184 to 0.3870; “*Medium*” from 0.3870 to 0.5556; and “*High*” from 0.5556 to 0.7243. The following formula was used to define the three R^2 categories: $(\text{Max } R^2 \text{ value observed} - \text{Min } R^2 \text{ value observed}) / 3$.

Results and Discussion

The current study investigated the application of polynomial regression analysis as a tool to define the response of pineapple plants to osmotic stress, applied in TIBs. The R^2 value represents a numerical statistical index that describes the goodness of fit of a regression model. In the present work, R^2 was used to indicate what fraction of y variations occurred as a consequence of x variations (Ivanov, 1989). Figure 1 shows the analysis and determination coefficient (R^2) values corresponding to the independent variables chlorophyll a (panel A), chlorophyll b (B), carotenoids (C), soluble phenolics (D), and Figure 2 shows cell wall-linked phenolics (A), soluble phenolics in the culture medium (B), MDA (C) and other aldehydes (D). The results indicated that the soluble and cell wall-linked phenolics, and other aldehydes affected the changes in shoot biomass observed in response to osmotic stress (Figure 1D, Figure 2A, D). In this regard, 72.11% of the variations of pineapple shoot cluster fresh weight under mannitol stress correlated with changes in soluble phenolic levels ($R^2 = 0.7211$; Figure 1D). The determination coefficient for this relationship was classified as “*High*”.




Figure 1. Determination coefficients (R^2) obtained in polynomial regressions analyses (Order 2: $y = ax^2 + bx + c$; Microsoft Excel). The fresh weight of shoot clusters was the dependent variable (y), and chlorophyll a (A), chlorophyll b (B), carotenoids (C) and soluble phenolics (D), the independent variables. R^2 were classified as "Low" from 0.2184 to 0.3870; "Medium" from 0.3870 to 0.5556; and "High" from 0.5556 to 0.7243

The results showed that as the soluble phenolic content of shoots increased, shoot weight declined, in agreement with the trend detected in the empirical data. A similar trend was observed for cell wall-linked phenolics and other aldehydes, i.e. as the levels of these two metabolites increased, shoot mass was adversely affected. These two factors also generated "High" R^2 values: 68.17% of shoot cluster weight variations were explained by modifications of cell wall-linked phenolic contents (Figure 2A), and 72.43% by increases in the levels of other aldehydes (Figure 2D).

"Medium" R^2 were recorded for carotenoid levels ($R^2 = 50.95\%$, Figure 1C) and MDA contents ($R^2 = 51.61\%$, Figure 2C). Changes in levels of chlorophylls a and b (Figure 1A, B), and soluble phenolics excreted into the culture medium (Figure 2B) did not appear to (mathematically) influence biomass of pineapple shoots in TIBs under osmotic stress, as the corresponding R^2 values were classified as "Low".

From our previous work (Gómez *et al.*, 2017), it was established that exposure of pineapple shoots to mannitol above 50 mM for 30 days reduced the fresh weight of shoots with respect to the non-stressed controls; for example, the highest mannitol concentration tested (200 mM) decreased the multiplication rate by 59.7% and cluster fresh weight by 59.8%. From that study, several significant differences were found between the control and mannitol treatments using basic statistical analyses, namely one-way ANOVA and determination of 'overall coefficients of variation' (OCVs).




Figure 2. Determination coefficients (R^2) obtained in polynomial regressions analyses (Order 2: $y = ax^2 + bx + c$; Microsoft Excel). The fresh weight of shoot clusters was the dependent variable (y), and cell wall-linked phenolics (A), soluble phenolics in the culture medium (B), MDA (C) and other aldehydes (D), the independent variables. R^2 were classified as "Low" from 0.2184 to 0.3870; "Medium" from 0.3870 to 0.5556; and "High" from 0.5556 to 0.7243.

Applying the more precise and robust polynomial regression analysis, those data have been reanalysed, which allowed refining the conclusions of the previous study. To give just some examples, both chlorophyll b and shoot soluble phenolics contents increased in response to the osmotic stress applied; comparing the controls with the 200 mM mannitol treatment, this increase amounted to 1.5-fold for chlorophyll b contents (from 36.93 to 54.29 $\mu\text{g g}^{-1}$ fresh weight), and to 2.1-fold for soluble phenolics (from 430.15 to 904.03 $\mu\text{g g}^{-1}$ fresh weight). In both cases, the increments were statistically significant (one-way ANOVA, Tukey, $p > 0.05$), with 'medium' OCVs (Gómez *et al.*, 2017). However, polynomial regression analysis clearly distinguished the two putative stress indicators, ruling out the use of chlorophyll b (with a 'Low' R^2 value) as a reliable biochemical marker of osmotic stress in this specific experimental system, as compared to soluble phenolics, which showed a 'High' R^2 value. Similarly, the significant increase in cell wall-linked phenolics and 'other aldehydes' in response to mannitol-induced osmotic stress, showing 'low' OCVs (Gómez *et al.*, 2017), were nevertheless strongly correlated ('High' determination coefficients, R^2) with the variations of pineapple shoot cluster fresh weight.

The impacts of climate change are already causing detrimental effects on agricultural productivity as well as in natural ecosystems. Warmer temperatures accompanied by longer dry spells and changing rainfall patterns affect all phases of plant development. Access to sufficient amounts of water (of good quality) at the appropriate developmental stage represents one of the most critical factors determining plant growth (Hernández *et al.*, 2015; Naik and Al-Khayri, 2016).

Plants have evolved complex mechanisms to mitigate the adverse effects caused by drought, with a general reduction in growth being the typical response (Avramova *et al.*, 2015). This could occur as a result of

changes in CO₂ assimilation (Gindaba *et al.*, 2004) and effects on water relations, such as relative water content (Galle *et al.*, 2007), leaf turgor pressure (Schachtman and Goodger, 2008) or osmotic potential (Silva *et al.*, 2010). Accompanying these physiological changes, alterations in the concentration of specific plant metabolites with protective functions have been detected. Since salt stress includes an osmotic component, as water deficit, similar groups of metabolites are affected by drought and salinity, including carbohydrate levels, starch content (Chao *et al.*, 2006; Zaher-Ara *et al.*, 2016), sugars and oligosaccharides (Anderson and Kohorn, 2001), antioxidants and enzymes (Avramova *et al.*, 2016; Avramova *et al.*, 2017). In particular, several compounds with protective properties have been identified, including proline (AbdElgawad *et al.*, 2015; Zaher-Ara *et al.*, 2016), genkwanin (Awasthi *et al.*, 2016), tanshinone IIA (Zaker *et al.*, 2015), steviol glycosides (Gupta *et al.*, 2015), *myo*-inositol (Díaz-López *et al.*, 2012), glycine-betaine (Quan *et al.*, 2004), anthocyanin (Awasthi *et al.*, 2016), and abscisic acid (Gurmani *et al.*, 2007).

From our previous work (Gómez *et al.*, 2017) and the present analysis of pineapple plants micropropagated in TIBs, the levels of soluble phenolics and other aldehydes can be defined as the most precise biomarkers of mannitol-induced osmotic stress; these two groups of compounds are known to play a role in plant responses to stress conditions. These results are in agreement with other published reports of studies conducted in different culture systems and with different plant species (Winkel-Shirley, 2002; Haghghi *et al.*, 2012; Rivelli *et al.*, 2012; Selmar and Kleinwächter, 2013; Boestfleisch *et al.*, 2014; Hernández *et al.*, 2015; Boestfleisch and Papenbrock, 2017). This observation provides evidence indicating that TIBs, in combination with regression analyses of biochemical stress indicators, are suitable systems to investigate plant responses to stress conditions.

Conclusions

The current study supports the use of polynomial regression analysis for comparing the relative importance of biochemical indicators of stress tolerance in *in vitro* screening studies. We recommend that this statistical approach be applied to other species and stressors to validate its utility for screening varieties for stress tolerance *in vitro*.

Authors' Contributions

Conceptualization: OV, JCL; Data curation: DG, JCL; Formal analysis: JCL; Funding acquisition: OV, JCL; Investigation: DG, DE, JCL; Methodology: JCL; Project administration: JCL; Resources: OV, JCL; Supervision: EH, OV, S, JCL; Writing - original draft: DG, EH, OV, S, JCL; Writing - review and editing: OV. All authors read and approved the final manuscript.

Acknowledgements

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. This research was supported by the Bioplant Centre (University of Ciego de Ávila, Cuba), the Universidad de Concepción (Chile), the Agricultural Research Council-Tropical and Subtropical Crops (South Africa), the Universitat Politècnica de València (Spain), and the South African Technology Network (South Africa).

Conflict of Interests

The authors declare that there are no conflicts of interest related to this article

References

- AbdElgawad H, De Vos D, Zinta G, Domagalska MA, Beemster GT, Asard H (2015). Grassland species differentially regulate proline concentrations under future climate conditions: an integrated biochemical and modelling approach. *New Phytologist* 208:354-369.
- Anderson C, Kohorn B (2001). Inactivation of *Arabidopsis* SIP1 leads to reduced levels of sugars and drought tolerance. *Journal of Plant Physiology* 158:1215-1219.
- Avramova V, AbdElgawad H, Zhang Z, Fotschki B, Casadevall R, Vergauwen L, Knapen D, Taleisnik E, Guisez Y, Asard H, Beemster GTS (2015). Drought induces distinct growth response, protection and recovery mechanisms in the maize leaf growth zone. *Plant Physiology* 169:1382-1396.
- Avramova V, Nagel KA, AbdElgawad H, Bustos D, DuPlessis M, Fiorani F, Beemster GT (2016). Screening for drought tolerance of maize hybrids by multi-scale analysis of root and shoot traits at the seedling stage. *Journal of Experimental Botany* 67:2453-2466.
- Avramova V, AbdElgawad H, Vasileva I, Petrova AS, Holek A, Mariën J, Beemster GT (2017). High antioxidant activity facilitates maintenance of cell division in leaves of drought tolerant maize hybrids. *Frontiers in Plant Science* 8:84.
- Awasthi P, Gupta AP, Bedi YS, Vishwakarma RA, Gandhi SG (2016). Mannitol stress directs flavonoid metabolism toward synthesis of flavones via differential regulation of two cytochrome p450 monooxygenases in *Coleus forskohlii*. *Frontiers in Plant Science* 7:985.
- Barker AV (1999). Foliar ammonium accumulation as an index of stress in plants. *Communications in Soil Science and Plant Analysis* 30:167-174.
- Boestfleisch C, Wagenseil NB, Buhmann AK, Seal CE, Wade EM, Muscolo A, Papenbrock J (2014). Manipulating the antioxidant capacity of halophytes to increase their cultural and economic value through saline cultivation. *Annals of Botany* PLANTS 6:plu046. <https://doi.org/10.1093/aobpla/plu046>
- Boestfleisch C, Papenbrock J (2017). Changes in secondary metabolites in the halophytic putative crop species *Crithmum maritimum*, *Triglochin maritima* and *Halimione portulacoides* as reaction to mild salt stress. *PLoS ONE* 12:e0176303. <https://doi.org/10.1371/journal.pone.0176303>
- Chao SK, Kim JE, Jong AP, Eom TJ, Kim WT (2006). Constitutive expression of abiotic stress-inducible hot pepper CaXTH3, which encodes a xyloglucan endotransglycosylase/hydrolase homolog, improves drought and salt tolerance in transgenic *Arabidopsis* plants. *FEBS Letters* 580:3136-3144.
- Daquinta M, Benegas R (1997). Brief review of tissue culture of pineapple. *Pineapple News* 3:7-9.
- Díaz-López L, Gimeno V, Simón I, Martínez V, Rodríguez-Ortega W, García-Sánchez F (2012). *Jatropha curcas* seedlings show a water conservation strategy under drought conditions based on decreasing leaf growth and stomatal conductance. *Agricultural Water Management* 105:48-56.
- Escalona M, Lorenzo JC, González B, Daquinta M, Borroto C, González JL, Desjardines Y (1999). Pineapple micropropagation in temporary immersion systems. *Plant Cell Reports* 18:743-748.
- Fernández RJ, Wang M, Reynolds JF (2002). Do morphological changes mediate plant responses to water stress? A steady state experiment with two C4 grasses. *New Phytologist* 155:79-88.
- Galle A, Haldimann P, Feller U (2007). Photosynthetic performance and water relations in young pubescent oak (*Quercus pubescens*) trees during drought stress and recovery. *New Phytologist* 174:799-810.

- Gindaba J, Rozanov A, Negash L (2004). Response of seedlings of two *Eucalyptus* and three deciduous tree species from Ethiopia to severe water stress. *Forest Ecology and Management* 201:119-129.
- Gómez D, Hernández L, Valle B, Martínez J, Cid M, Escalona M, Hernández M, Beemster GTS, Tebbe CC, Yabor L, Lorenzo JC (2017). Temporary immersion bioreactors (TIB) provide a versatile, cost-effective and reproducible *in vitro* analysis of the response of pineapple shoots to salinity and drought. *Acta Physiologiae Plantarum* 39:277.
- Gupta P, Sharma S, Saxena S (2015). Biomass yield and steviol glycoside production in callus and suspension culture of *Stevia rebaudiana* treated with proline and polyethylene glycol. *Applied Biochemistry and Biotechnology* 176:863-874.
- Gurmani AR, Bano A, Saleem M (2007). Effect of ABA and BA on growth and ion accumulation of wheat under salinity stress. *Pakistan Journal of Botany* 39:141-149.
- Gurr S, McPherson J, Bowles D (1992). Lignin and associated phenolic acids in cell walls. In: Wilkinson DL (Ed). *Molecular Plant Pathology*. Oxford Press, Oxford, pp 51-56.
- Haghighi Z, Modarresi M, Mollayi S (2012). Enhancement of compatible solute and secondary metabolites production in *Plantago ovata* Forsk. by salinity stress. *Journal of Medical Plant Research* 6:3495-3500.
- Heath R, Packer J (1968). Photoperoxidation in isolated chloroplast: I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics* 125:189-198.
- Hernández L, Loyola-González O, Valle B, Martínez J, Díaz-López L, Aragón C, Vicente O, Papenbrock J, Trethowan R, Yabor L, Lorenzo JC (2015). Identification of discriminant factors after exposure of maize and common bean plantlets to abiotic stresses. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 43:589-598.
- Hoagland RE (1990). *Alternaria cassiae* alters phenylpropanoid metabolism in Sicklepod (*Casia obtusifolia*). *Journal of Phytopathology* 130:177-187.
- Hüve K, Bichele I, Kaldmäe H, Rasulov B, Valladares F, Niinemets Ü (2019). Responses of Aspen Leaves to Heatflecks: Both Damaging and Non-Damaging Rapid Temperature Excursions Reduce Photosynthesis. *Plants* 8:145.
- Ivanov Z (1989). *The Agricultural Experimentation Pueblo y Educación*, Havana, pp 332.
- Lichtenthaler HK (1987). Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods of Enzymology* 148:350-382.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiologiae Plantarum* 5:473-497.
- Naik PM, Al-Khayri JM (2016). Abiotic and biotic elicitors—Role in secondary metabolites production through *in vitro* culture of medicinal plants. In: Shanker AK, Shanker C (Eds). *Abiotic and Biotic Stress in Plants - Recent Advances and Future Perspectives*. InTech, Al-Hassa, pp 247-277. <http://dx.doi.org/10.5772/61442>
- Peet MM, Willits D, Gardner R (1997). Response of ovule development and post-pollen production processes in male-sterile tomatoes to chronic, sub-acute high temperature stress. *Journal of Experimental Botany* 48:101-111.
- Porra R (2002). The chequered history of the development and use of simultaneous equations for the accurate determination of chlorophylls a and b. *Photosynthesis Research* 73:149-156.
- Quan R, Shang M, Zhang H, Zhao Y, Zhang J (2004). Engineering of enhanced glycine betaine synthesis improves drought tolerance in maize. *Plant Biotechnology Journal* 2:477-486.
- Rivelli A, De Maria S, Pizza S, Gherbina P (2012). Growth and physiological response of hydroponically grown sunflower as affected by salinity and magnesium levels. *Journal of Plant Nutrition* 33:1307-1323.
- Runeckles V (1982). Relative death rate: a dynamic parameter describing plant response to stress. *Journal of Applied Ecology* 295-303.
- Schachtman D, Goodger J (2008). Chemical root to shoot signaling under drought. *Trends in Plant Science* 13:281-287.

- Selmar D, Kleinwächter M (2013). Influencing the product quality by deliberately applying drought stress during the cultivation of medicinal plants. *Indian Crops Production* 42:558- 566.
- Silva E, Ferreira-Silva S, Viegas R, Silveira J (2010). The role of organic and inorganic solutes in the osmotic adjustment of drought-stressed *Jatropha curcas* plants. *Environmental and Experimental Botany* 69:279-285.
- Vieira DA, Mesquita AC, Marinho LB, Souza Vd, Aida SdT, Carvalho MMP (2019). Gas exchanges of melon under water stress in the Submedium region of the São Francisco River Valley. *Acta Scientiarum Agronomy* 41. <https://doi.org/10.4025/actasciagron.v41i1.42686>
- Wijewardana C, Alsajri FA, Reddy KR (2019). Soybean seed germination response to *in vitro* osmotic stress. *Seed Technology* 39:143-154.
- Winkel-Shirley B (2002). Biosynthesis of flavonoids and effects of stress. *Current Opinion in Plant Biology* 5:218-223.
- Zaher-Ara T, Boroomand N, Sadat-Hosseini M (2016). Physiological and morphological response to drought stress in seedlings of ten citrus. *Trees* 30:985-993.
- Zaker A, Sykora C, Gössnitzer F, Abrishamchi P, Asili J, Mousavi SH, Wawrosch C (2015). Effects of some elicitors on tanshinone production in adventitious root cultures of *Perovskia abrotanoides* Karel. *Indian Crops Production* 67:97-102.



The journal offers free, immediate, and unrestricted access to peer-reviewed research and scholarly work. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.



License - Articles published in *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* are Open-Access, distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) License.

© Articles by the authors; UASVM, Cluj-Napoca, Romania. The journal allows the author(s) to hold the copyright/to retain publishing rights without restriction.