

## Orbitally discrete coarse spaces

IGOR PROTASOV

Taras Shevchenko National University of Kyiv, Department of Computer Science and Cybernetics,  
Academic Glushkov pr. 4d, 03680 Kyiv, Ukraine (i.v.protasov@gmail.com)

Communicated by F. Mynard

### ABSTRACT

---

Given a coarse space  $(X, \mathcal{E})$ , we endow  $X$  with the discrete topology and denote  $X^\sharp = \{p \in \beta G : \text{each member } P \in p \text{ is unbounded}\}$ . For  $p, q \in X^\sharp$ ,  $p||q$  means that there exists an entourage  $E \in \mathcal{E}$  such that  $E[P] \in q$  for each  $P \in p$ . We say that  $(X, \mathcal{E})$  is orbitally discrete if, for every  $p \in X^\sharp$ , the orbit  $\overline{p} = \{q \in X^\sharp : p||q\}$  is discrete in  $\beta G$ . We prove that every orbitally discrete space is almost finitary and scattered.

---

2010 MSC: 54D80; 20B35; 20F69.

KEYWORDS: coarse space; ultrafilter; orbitally discrete space; almost finitary space; scattered space.

### 1. INTRODUCTION AND PRELIMINARIES

Given a set  $X$ , a family  $\mathcal{E}$  of subsets of  $X \times X$  is called a *coarse structure* on  $X$  if

- each  $E \in \mathcal{E}$  contains the diagonal  $\Delta_X = \{(x, x) \in X : x \in X\}$ ;
- if  $E, E' \in \mathcal{E}$  then  $E \circ E' \in \mathcal{E}$  and  $E^{-1} \in \mathcal{E}$ , where  $E \circ E' = \{(x, y) : \exists z((x, z) \in E, (z, y) \in E')\}$ ,  $E^{-1} = \{(y, x) : (x, y) \in E\}$ ;
- if  $E \in \mathcal{E}$  and  $\Delta_X \subseteq E' \subseteq E$  then  $E' \in \mathcal{E}$ ;
- $\bigcup \mathcal{E} = X \times X$ .

A subfamily  $\mathcal{E}' \subseteq \mathcal{E}$  is called a *base* for  $\mathcal{E}$  if, for every  $E \in \mathcal{E}$ , there exists  $E' \in \mathcal{E}'$  such that  $E \subseteq E'$ . For  $x \in X$ ,  $A \subseteq X$  and  $E \in \mathcal{E}$ , we denote

$$E[x] = \{y \in X : (x, y) \in E\}, \quad E[A] = \bigcup_{a \in A} E[a], \quad E_A[x] = E[x] \cap A$$

and say that  $E[x]$  and  $E[A]$  are *balls of radius  $E$  around  $x$  and  $A$* .

The pair  $(X, \mathcal{E})$  is called a *coarse space* [19] or a *balleen* [12], [18].

For a coarse space  $(X, \mathcal{E})$ , a subset  $B \subseteq X$  is called *bounded* if  $B \subseteq E[x]$  for some  $E \in \mathcal{E}$  and  $x \in X$ . The family  $\mathcal{B}_{(X, \mathcal{E})}$  of all bounded subsets of  $(X, \mathcal{E})$  is called the *bornology* of  $(X, \mathcal{E})$ . We recall that a family  $\mathcal{B}$  of subsets of a set  $X$  is a *bornology* if  $\mathcal{B}$  is closed under taking subsets and finite unions, and  $\mathcal{B}$  contains all finite subsets of  $X$ .

A coarse space  $(X, \mathcal{E})$  is called *finitary*, if for each  $E \in \mathcal{E}$  there exists a natural number  $n$  such that  $|E[x]| < n$  for each  $x \in X$ .

Let  $G$  be a transitive group of permutations of a set  $X$ . We denote by  $X_G$  the set  $X$  endowed with the coarse structure with the base

$$\{(x, gx) : g \in F\} : F \in [G]^{<\omega}, \text{ id} \in F\}.$$

By [8, Theorem 1], for every finitary coarse structure  $(X, \mathcal{E})$ , there exists a transitive group  $G$  of permutations of  $X$  such that  $(X, \mathcal{E}) = X_G$ . For more general results, see [10].

Let  $X$  be a discrete space and let  $\beta X$  denote the *Stone-Ćech compactification* of  $X$ . We take the points of  $\beta X$  to be the ultrafilters on  $X$ , with the points of  $X$  identified with the principal ultrafilters, so  $X^* = \beta X \setminus X$  is the set of all free ultrafilters. The topology of  $\beta X$  is generated by the base consisting of the sets  $\bar{A} = \{p \in \beta X : A \in p\}$ , where  $A \subseteq X$ . The universal property of  $\beta X$  states that every mapping  $f : X \rightarrow Y$  to a compact Hausdorff space  $Y$  can be extended to a continuous mapping  $f^\beta : \beta X \rightarrow Y$ .

Given a coarse space  $(X, \mathcal{E})$ , we endow  $X$  with the discrete topology and denote by  $X^\sharp$  the set of all ultrafilters  $p$  on  $X$  such that each member  $P \in p$  is unbounded. Clearly,  $X^\sharp$  is a closed subset of  $X^*$  and  $X^\sharp = X^*$  if  $(X, \mathcal{E})$  is finitary.

Following [7], we say that two ultrafilters  $p, q \in X^\sharp$  are *parallel* (and write  $p||q$ ) if there exists  $E \in \mathcal{E}$  such that  $E[P] \in q$  for each  $P \in p$ . Then  $||$  is an equivalence on  $X^\sharp$ . We denote

$$\bar{p} = \{q \in X^\sharp : q||p\}$$

and say that  $\bar{p}$  is the *orbit* of  $p$ . If  $(X, \mathcal{E})$  is finitary and  $(X, \mathcal{E}) = X_G$  then  $\bar{p} = Gp$ .

A coarse space  $(X, \mathcal{E})$  is called *orbitally discrete* if, for every  $p \in X^\sharp$ , the orbit  $\bar{p}$  is discrete. Every discrete coarse space is orbitally discrete. We recall that  $(X, \mathcal{E})$  is *discrete* if, for each  $E \in \mathcal{E}$ , there exists a bounded subset  $B$  such that  $E[x] = \{x\}$  for each  $x \in X \setminus B$ . In this case,  $\bar{p} = \{p\}$  for each  $p \in X^\sharp$ .

Every bornology  $\mathcal{B}$  on a set  $X$  defines the discrete coarse structure on  $X$  with the base  $\{E_B : B \in \mathcal{B}\}$ ,  $E_B[x] = B$  if  $x \in B$ , and  $E_B[x] = \{x\}$  if  $x \in X \setminus B$ .

By [15, Theorem 5.4], for a finitary coarse space  $(X, \mathcal{E})$ , the following conditions are equivalent:  $X_G$  is orbitally discrete,  $X_G$  is scattered,  $X_G$  has no piecewise shifted FP-sets.

A coarse space  $(X, \mathcal{E})$  is called *scattered* if, for every unbounded subset  $A$  of  $X$ , there exists  $E \in \mathcal{E}$  such that  $A$  has asymptotically  $E$ -isolated balls: for each  $E' \in \mathcal{E}$ , there is  $a \in A$  such that  $E'_A[a] \setminus E_A[a] = \emptyset$ .

This notion arose in the characterization of the Cantor macrocube [3] and, in the case of finitary coarse groups, was explored in [2].

Let  $G$  be a group of permutations of a set  $X$ . Let  $(g_n)_{n \in \omega}$  be a sequence in  $G$  and let  $(x_n)_{n \in \omega}$  be a sequence in  $X$  such that

- (1)  $\{g_0^{\epsilon_0} \dots g_n^{\epsilon_n} x_n : (\epsilon_i)_{i=0}^n \in \{0, 1\}^{n+1}\} \cap \{g_0^{\epsilon_0} \dots g_m^{\epsilon_m} x_m : (\epsilon_i)_{i=0}^m \in \{0, 1\}^{m+1}\} = \emptyset$  for all distinct  $n, m \in \omega$ ;
- (2)  $|\{g_0^{\epsilon_0} \dots g_n^{\epsilon_n} x_n : (\epsilon_i)_{i=0}^n \in \{0, 1\}^{n+1}\}| = 2^{n+1}$  for every  $n \in \omega$ .

Following [15], we say that a subset  $Y$  of  $X$  is a *piecewise shifted FP-set* if there exist  $(g_n)_{n \in \omega}$ ,  $(x_n)_{n \in \omega}$  satisfying (1), (2) and such that

$$Y = \{g_0^{\epsilon_0} \dots g_n^{\epsilon_n} x_n : \epsilon_i \in \{0, 1\}\}, n \in \omega\}.$$

After exposition of results in Section 2, we survey some known classes of orbitally discrete spaces in Section 3.

## 2. RESULTS

A coarse space  $(X, \mathcal{E})$  is called *almost finitary* if, for every  $E \in \mathcal{E}$ , there exists a bounded subset  $B$  and a natural number  $n$  such that  $|E[x]| < n$  for each  $x \in X \setminus B$ . Every discrete space and every finitary space are almost finitary.

**Theorem 2.1.** *Every orbitally discrete coarse space is almost finitary.*

*Proof.* We suppose the contrary and choose  $E \in \mathcal{E}$ ,  $E = E^{-1}$  such that, for any bounded subset  $B$  and a natural number  $n$ , there exists  $x \in X \setminus B$  such that  $|E[x]| > n$ .

We claim that there exists  $p \in X^\sharp$  such that, for every  $P \in p$ ,  $\{x \in P : |E^2[x] \cap P| > 1\} \in p$ . Otherwise, for every  $p \in X^\sharp$ , there exists  $Q_p \in p$  such that  $\{x \in Q_p : |E^2[x] \cap Q_p| = 1\} \in p$ . We consider the open covering  $\{Q_p^\sharp : p \in X^\sharp\}$  of  $X^\sharp$  and choose its finite subcovering  $Q_{p_1}^\sharp, \dots, Q_{p_m}^\sharp$ . Then the set  $B = X \setminus (Q_{p_1} \cup \dots \cup Q_{p_m})$  is bounded and  $|E[x]| \leq m$  for each  $x \in X \setminus E[B]$ , but this contradicts the choice of  $E$ .

We show that the orbit  $\overline{p}$  is not discrete. Given any  $P \in p$ , we choose  $Q \in p$ ,  $Q \subseteq P$  such that  $|E^2[x] \cap P| > 1$  for each  $x \in Q$ . For every  $x \in Q$ , we take  $f(x) \in E^2[x] \cap P$  such that  $x \neq f(x)$ . Then we extend the mapping  $x \mapsto f(x)$

from  $Q$  to  $X$  by  $f(x) = x$  for each  $x \in X \setminus Q$ . Clearly,  $f^\beta(p) \neq p$ ,  $P \in f^\beta(p)$  and  $f^\beta(p) \not\ll p$  because  $(x, f(x)) \in E^2$  for each  $x \in X$ .  $\square$

To clarify the structure of an almost finitary coarse space, we use the following construction from [6]. A bornology  $\mathcal{B}$  on a coarse space  $(X, \mathcal{E})$  is called  $\mathcal{E}$ -compatible if  $E[B] \in \mathcal{B}$  for all  $B \in \mathcal{B}$ ,  $E \in \mathcal{E}$ . Every  $\mathcal{E}$ -compatible bornology  $\mathcal{B}$  defines the  $\mathcal{B}$ -strengthening  $(X, \mathcal{H})$  of  $(X, \mathcal{E})$ , where  $\mathcal{H}$  has the base

$$\{H_{B,E} : B \in \mathcal{B}, E \in \mathcal{E}\},$$

$$H_{B,E}[x] = \begin{cases} E[B], & \text{if } x \in B, \\ E[x], & \text{if } x \in X \setminus B. \end{cases}$$

For description of the upper bound  $\mathcal{E} \vee \mathcal{E}'$  of coarse structures, see [13].

**Theorem 2.2.** *For a coarse space  $(X, \mathcal{E})$ , the following statements are equivalent*

- (i)  $(X, \mathcal{E})$  is almost finitary;
- (ii)  $(X, \mathcal{E})$  is the  $\mathcal{B}$ -strengthening of some finitary coarse space  $(X, \mathcal{E}')$  by the bornology  $\mathcal{B}$  of bounded subspaces of  $(X, \mathcal{E})$ ;
- (iii)  $\mathcal{E}$  is the upper bound of a discrete and a finitary coarse structures on  $X$ .

*Proof.* (i)  $\implies$  (ii). For  $B \in \mathcal{B}$  and  $E \in \mathcal{E}$ , we pick  $B'_{B,E} \in \mathcal{B}$  and a natural number  $n$  such that  $B \subseteq B'_{B,E}$  and  $|E[x]| < n$  for each  $x \in X \setminus B'_{B,E}$ . We note that  $\{B'_{B,E} : B \in \mathcal{B}, E \in \mathcal{E}\}$  is a base for  $\mathcal{B}$ . For  $B \in \mathcal{B}$ ,  $E \in \mathcal{E}$  we put

$$E'_{B,E} = \begin{cases} x & \text{if } x \in B'_{B,E}, \\ E[x] & \text{if } x \in X \setminus B'_{B,E}, \end{cases}$$

denote by  $\mathcal{E}'$  the smallest coarse structure on  $X$  containing all entourages  $\{H_{B,E} : B \in \mathcal{B}, E \in \mathcal{E}\}$ , observe that  $\mathcal{E}'$  is finitary and  $(X, \mathcal{E})$  is the  $\mathcal{B}$ -strengthening of  $(X, \mathcal{E}')$ .

(ii)  $\implies$  (iii). If  $(X, \mathcal{E})$  is the  $\mathcal{B}$ -strengthening of  $(X, \mathcal{E}')$  then  $\mathcal{E}$  is the upper bounded of  $\mathcal{E}'$  and the discrete coarse structure on  $X$  defined by the bornology  $\mathcal{B}$ .

(iii)  $\implies$  (i). We assume that  $\mathcal{E}$  is the upper bound of finitary coarse structure  $\mathcal{E}'$  and discrete coarse structure on  $X$  defined by some bornology  $\mathcal{B}$ . We choose the smallest bornology  $\mathcal{B}'$  on  $X$  such that  $\mathcal{B} \subseteq \mathcal{B}'$  and  $E'(B') \in \mathcal{B}'$  for all  $E' \in \mathcal{E}'$ . Then  $\mathcal{B}'$  is the bornology of bounded subsets of  $(X, \mathcal{E})$  and  $(X, \mathcal{E})$  is the  $\mathcal{B}'$ -strengthening of  $(X, \mathcal{E}')$ , so  $(X, \mathcal{E})$  is almost finitary.  $\square$

**Remark.** Let  $(X, \mathcal{E})$  be the  $\mathcal{B}$ -strengthening of a finitary coarse space  $(X, \mathcal{E}')$ . If  $(X, \mathcal{E}')$  is orbitally discrete then  $(X, \mathcal{E})$  is orbitally discrete, but the converse statement needs not to be true. Let  $X$  be the disjoint union of

two infinite subsets  $Y, Z$ . We endow  $Y$  with the finitary coarse structure  $\mathcal{E}_Y$  such that  $(Y, \mathcal{E}_Y)$  is not orbitally discrete, and denote by  $\mathcal{E}_Z$  the discrete coarse structure on  $Z$  defined by the bornology of finite subset. We take the smallest coarse structure  $\mathcal{E}'$  on  $X$  such that  $\mathcal{E}'|_Y = \mathcal{E}_Y, \mathcal{E}'|_Z = \mathcal{E}_Z$ . Clearly,  $\mathcal{E}'$  is finitary but not orbitally discrete. We denote by  $\mathcal{B}$  the smallest bornology on  $X$  such that  $Y \in \mathcal{B}$ . Then the  $\mathcal{B}$ -strengthening of  $(X, \mathcal{E}')$  is discrete.

**Theorem 2.3.** *For almost finitary coarse space  $(X, \mathcal{E})$  and  $p, q \in X^\sharp$ , we have  $p||q$  if and only if there exist  $E \in \mathcal{E}$  and a permutation  $g$  of  $X$  such that  $gp = q, gp = \{gP : P \in p\}$  and  $(x, gx) \in E$  for each  $x \in X$ .*

*Proof.* Let  $p||q$ . We take  $E \in \mathcal{E}$  such that  $E = E^{-1}$  and  $E[P] \in q$  for each  $P \in p$ . Since  $(X, \mathcal{E})$  is almost finitary, there exist a bounded subset  $B$  of  $X$  and a natural number  $n$  such that  $|E[x]| < n$  for each  $x \in X \setminus B$ . We put  $Y = X \setminus E[B]$ , note that  $Y \in p$  and define a set-valued mapping  $\mathcal{F} : X \rightarrow [x]^{<\omega}$ .  $\mathcal{F}(x) = E[x]$  if  $x \in Y$  and  $\mathcal{F}(x) = \{x\}$  if  $x \in X \setminus Y$ . By Theorem 1 from [10], there exists bijection  $f_1, \dots, f_m$  of  $X$  such that  $f_i(x) \in \mathcal{F}(x)$  and  $f_1(x) \cup \dots \cup f_m(x) = \mathcal{F}(x)$ . We take  $i \in \{1, \dots, m\}$  such that  $f_i(P) \in q$  for each  $P \in p$  and put  $g = f_i$ .

The converse statement follows directly from the definition of the parallelity relation  $||$ . □

**Corollary 2.4.** *If  $(X, \mathcal{E})$  is almost finitary,  $p \in X^\sharp$  and  $p$  is an isolated point of  $\overline{p}$  then  $\overline{p}$  is discrete.*

*Proof.* We assume that some point  $q \in \overline{p}$  is not isolated in  $\overline{p}$ , use Theorem 2.3 to choose a permutation  $g$  of  $X$  such that  $gq = p$  and note that  $p$  is not isolated in  $\overline{p}$ . □

For a subset  $A$  of  $(X, \mathcal{E})$  and  $p \in X^\sharp$ , we denote  $\Delta_p(A) = \overline{p} \cap A^\sharp$ .

**Theorem 2.5.** *An almost finitary coarse space  $(X, \mathcal{E})$  is scattered if and only if, for every unbounded subset  $A$  of  $X$ , there exists  $p \in A$  such that  $\Delta_p(A)$  is finite.*

*Proof.* We suppose that  $X$  is scattered and choose  $E \in \mathcal{E}$  such that  $A$  has an asymptotically isolated  $E$ -balls. For each  $H \in \mathcal{E}$ , we denote  $P_H = \{x \in A : H_A[x] \setminus E_A[x] = \emptyset\}$  and take  $p \in A^\sharp$  such that  $P_H \in p$  for each  $H \in \mathcal{E}$ . If  $q \in A^\sharp$  and  $q||p$  then  $E[P] \in q$  for each  $P \in p$ . We take the bijections  $f_1, \dots, f_m$  from the proof of Theorem 2.3. Since  $q = gp$  for some  $g \in \{f_1, \dots, f_m\}$ , we have  $\Delta_p(A) \leq m$ .

Let  $\Delta_p(A) = \{p_1, \dots, p_m\}$ . For each  $i \in \{1, \dots, m\}$ , we pick  $E_i \in \mathcal{E}$  such that  $E_i[p] \in p_i$  for each  $P \in p$ . Then we take  $E \in \mathcal{E}$  such that  $E_i \subseteq E$  for each  $i \in \{1, \dots, m\}$ , and observe that  $A$  has an asymptotically isolated  $E$ -balls. □

**Theorem 2.6.** *Every orbitally discrete space is scattered.*

*Proof.* To apply Theorem 2.5, we take an arbitrary unbounded subset  $A$  of  $X$  and find  $p \in A^\sharp$  such that  $\Delta_p(A)$  is finite.

We use the Zorn lemma to choose a minimal (by inclusion) closed subset  $S$  of  $A^\sharp$  such that  $\Delta_q(A) \subseteq S$  for each  $q \in S$ . Let  $p \in S$  but  $\Delta_p(A)$  is infinite. We take the limit point  $q$  of  $\Delta_p(A)$ . By the minimality of  $S$ , we have  $p \in cl\Delta_q(A)$ . Applying Theorem 2.3, we conclude that  $p$  is not isolated in  $\overline{p}$ .  $\square$

**Question.** *Let  $X$  be an almost finitary scattered space. Is  $X$  orbitally discrete?*

### 3. COMMENTS

1. For a natural number  $n$ , a coarse space  $(X, \mathcal{E})$  is called *n-thin* if, for every  $E \in \mathcal{E}$ , there exists a bounded subset  $B$  of  $X$  such that  $|E[x]| \leq n$ , for every  $x \in X \setminus B$ . A space  $(X, \mathcal{E})$  is *n-thin* if and only if  $|\overline{p}| \leq n$  for each  $p \in X^\sharp$ .

For finite partitions of an *n-thin* space into discrete subspaces, see [5], [14], [17], [1, Section 6].

2. A coarse space  $(X, \mathcal{E})$  is called *sparse* if each orbit  $\overline{p}$ ,  $p \in X^\sharp$  is finite. Sparse subsets of groups are studied in [4], [16]. For sparse metric spaces, see [9].
3. A coarse space  $(X, \mathcal{E})$  is called *indiscrete* if each discrete subspace of  $X$  is bounded. By Theorem 3.15 from [11], a finitary indiscrete space has no unbounded orbitally discrete subspaces. We do not know whether this statement holds for any almost finitary indiscrete spaces.

### REFERENCES

- [1] T. Banach and I. Protasov, Set-theoretical problems in Asymptology, arXiv: 2004.01979.
- [2] T. Banach, I. Protasov and S. Slobodianiuk, Scattered subsets of groups, Ukr. Math. J. 67 (2015), 347–356.
- [3] T. Banach and I. Zarichnyi, Characterizing the Cantor bi-cube in asymptotic categories, Groups Geom. Dyn. 5, no. 4 (2011), 691–728.
- [4] Ie. Lutsenko and I. Protasov, Space, thin and other subsets of groups, Intern. J. Algebra Comput. 19 (2009), 491–510.
- [5] Ie. Lutsenko and I. V. Protasov, Thin subset of balleans, Appl. Gen. Topology 11 (2010), 89–93.
- [6] O. Petrenko and I. V. Protasov, Balleans and filters, Mat. Stud. 38 (2012), 3–11.
- [7] I. V. Protasov, Normal ball structures, Mat. Stud. 20 (2003), 3–16.
- [8] I. V. Protasov, Balleans of bounded geometry and  $G$ -spaces, Algebra Discrete Math. 7, no. 2 (2008), 101–108.
- [9] I. V. Protasov, Sparse and thin metric spaces, Mat. Stud. 41 (2014), 92–100.
- [10] I. Protasov, Decompositions of set-valued mappings, Algebra Discrete Math. 30, no. 2 (2020), 235–238.
- [11] I. Protasov, Coarse spaces, ultrafilters and dynamical systems, Topol. Proc. 57 (2021), 137–148.
- [12] I. Protasov and T. Banach, Ball Structures and Colorings of Groups and Graphs, Mat. Stud. Monogr. Ser. Vol. 11, VNTL, Lviv, 2003.
- [13] I. Protasov and K. Protasova, Lattices of coarse structures, Math. Stud. 48 (2017), 115–123.

- [14] I. V. Protasov and S. Slobodianiuk, Thin subsets of groups, *Ukrain. Math. J.* 65 (2013), 1245–1253.
- [15] I. Protasov and S. Slobodianiuk, On the subset combinatorics of  $G$ -spaces, *Algebra Discrete Math.* 17, no. 1 (2014), 98–109.
- [16] I. Protasov and S. Slobodianiuk, Ultracompanions of subsets of a group, *Comment. Math. Univ. Carolin.* 55, no. 1 (2014), 257–265.
- [17] I. Protasov and S. Slobodianiuk, The dynamical look at the subsets of groups, *Appl. Gen. Topology* 16 (2015), 217–224.
- [18] I. Protasov and M. Zarichnyi, *General Asymptology*, Math. Stud. Monogr. Ser., Vol. 12, VNTL, Lviv, 2007.
- [19] J. Roe, *Lectures on Coarse Geometry*, Univ. Lecture Ser., vol. 31, American Mathematical Society, Providence RI, 2003.