

Index

Abstract	1
Resumen	3
Resum	5
General Introduction	7
1. <i>Cannabis sativa</i> L. botany	9
1.1. Taxonomy	11
2. <i>Cannabis sativa</i> history, uses and economic potential	15
2.1. Origin, dispersion and domestication.....	15
2.2. <i>Cannabis</i> : a multipurpose crop	20
2.3. Legal context of its cultivation	23
2.4. <i>Cannabis</i> market and economic predictions	27
3. Biotechnological approaches for <i>C. sativa</i> genetic improvement	28
3.1. <i>In vitro</i> plant tissue culture	29
3.1.1. Plant <i>in vitro</i> regeneration	30
3.2. Polyploidization	31
3.2.1. Polysomaty	32
3.3. Double haploids	33
3.3.1. Androgenesis	34
3.3.2. Factors influencing androgenesis induction	35
3.4. Plant genetic transformation.....	38
Objectives	41
Chapter 1: “Development of a direct <i>in vitro</i> plant regeneration protocol from <i>Cannabis sativa</i> L. seedling explants: developmental morphology of shoot regeneration and ploidy level of regenerated plants”	45
1. Abstract	48
2. Introduction	50
3. Material and methods	52
3.1. Plant material and growth conditions	52
3.2. <i>In vitro</i> shoot organogenesis experiments	54
3.3. Developmental morphology of the <i>in vitro</i> regeneration process	55
3.4. Determination of ploidy level of explants and <i>in vitro</i> regenerants	55
3.5. Data analyses	56
4. Results	57
4.1. Effect of genotype, explant and medium on <i>in vitro</i> shoot organogenesis of <i>C. sativa</i>	57
4.2. Developmental morphology of <i>in vitro</i> shoot organogenesis in <i>C. sativa</i>	64
4.3. Rooting of explants and spontaneous rooting of hypocotyl derived plants	72
4.4. Acclimatization of <i>in vitro</i> regenerated plants	74
4.5. Ploidy evaluation of explants and <i>in vitro</i> regenerated plants of <i>C. sativa</i>	76
5. Discussion	79
5.1. Hypocotyl presents a high potential for <i>in vitro</i> direct regeneration of <i>C. sativa</i> plants without addition of plant growth regulators to the culture medium	79
5.2. Hypocotyl derived plants can root spontaneously in hormone-free medium, being completely acclimatized in only six weeks	82
5.3. Pericycle cells adjacent to xylem poles could be the origin of <i>in vitro</i> regenerated plants of <i>C. sativa</i>	83
5.4. Polysomaty is present in cotyledons and hypocotyls of <i>C. sativa</i> seedlings.....	85
5.5. Mixoploid plants can be regenerated after <i>in vitro</i> culture of hypocotyls and cotyledons coming from seedlings of <i>C. sativa</i>	86

6. Conclusion	87
7. Statements	88
8. References	89
Chapter 2: "Microgametophyte development in <i>Cannabis sativa</i> L. and first androgenesis induction through microspore embryogenesis"	101
1. Abstract	104
2. Introduction	105
3. Materials and methods	109
3.1. Plant material and growth conditions	109
3.2. Characterization of nuclear dynamics during meiosis, microsporogenesis and microgametogenesis through light and fluorescence microscopy	110
3.3. Analysis of exine evolution by means of scanning electron microscopy (SEM)	111
3.4. Study of anther wall formation using cryogenic scanning electron microscopy (cryo-SEM)	112
3.5. Histochemical detection of starch through light microscopy	113
3.6. Correlation of the bud length with the stage of development of microspores and pollen grains	114
3.7. <i>In vitro</i> microspore culture experiments	115
3.8. Data analyses	116
4. Results	117
4.1. Male reproductive anatomy of <i>C. sativa</i> , microspore density per flower bud and <i>in vivo</i> viability of microspores	117
4.2. Cellular characterization of the different developmental stages of meiosis, microsporogenesis and microgametogenesis	121
4.3. Evolution of the exine throughout microsporogenesis and microgametogenesis	127
4.4. Development of the different layers that compose the anther wall during stamen formation	130
4.5. Amyloplasts in anthers, microspores and pollen grains from <i>C. sativa</i>	133
4.6. Correlation of the bud length with the different <i>C. sativa</i> microgametophyte developmental stages	136
4.7. Effect of cold-shock bud pretreatment on microspore viability, amyloplast content, and development of multicellular structures of androgenic origin	138
5. Discussion	142
5.1. <i>Cannabis sativa</i> is a suitable candidate for androgenesis experiments mainly due to its fast production of viable and potentially inducible microspores and pollen grains	142
5.2. Anthers from <i>C. sativa</i> present a high degree of uniformity in the developmental stages of microspores and pollen grains contained	143
5.3. The exine ornamentation of <i>C. sativa</i> microspores and pollen grains presents a scabrate sculpture which is almost completely covered by sporopollenin at maturity	144
5.4. The anther wall of <i>C. sativa</i> is composed by an external epidermis, endothecium, two middle layers and a secretory type of tapetum	145
5.5. Starch content of microspores and pollen grains from <i>C. sativa</i> is coincident with the amyloplast pattern observed in species recalcitrant to androgenesis	147
5.6. The bud length can be used as a floral morphological marker to identify male floral buds containing specific microgametophyte developmental stages	148

5.7. Although with an extremely low frequency, cold-shock bud pretreatment can promote microspore embryogenesis in <i>C. sativa</i>	150
6. Conclusion	153
7. Statements	154
8. References	154
Chapter 3: "A novel and rapid method for <i>Agrobacterium</i>-mediated production of stably transformed <i>Cannabis sativa</i> L. plants"	169
1. Abstract	172
2. Introduction	174
3. Materials and methods	176
3.1. Plant material and growth conditions	176
3.2. Hypocotyl and cotyledon explant <i>in vitro</i> culture, and hypocotyl kanamycin dose-response experiments	179
3.3. Media used, bacterial strain and transformation vector	179
3.4. <i>Agrobacterium tumefaciens</i> co-cultivation of explants and subsequent culture in antibiotic-containing medium	180
3.5. Visual analysis of GUS expression in hypocotyls, cotyledons and already-developed meristems after regenerant-derived tissue incubation with X-Gluc	181
3.6. Primer design and polymerase chain reaction (PCR) detection of β -glucuronidase (<i>uidA</i>) and kanamycin resistance neomycin phosphotransferase II (<i>nptII</i>) genes	181
3.7. Data analyses	182
4. Results	183
4.1. Efficiency comparison of <i>C. sativa</i> hypocotyl and cotyledon <i>in vitro</i> plant regeneration protocols	183
4.2. Effect of kanamycin on <i>C. sativa</i> hypocotyl regeneration in <i>in vitro</i> culture	190
4.3. Effect of <i>A. tumefaciens</i> co-culture and subsequent explant culture in antibiotic-containing selective regeneration medium in <i>in vitro</i> <i>C. sativa</i> regeneration, spontaneous rooting and albinism rates	194
4.4. Effect of explant on the production of <i>C. sativa</i> transgenic plants after <i>A. tumefaciens</i> co-culture and subsequent explant culture in antibiotic-containing selective regeneration medium	196
4.5. <i>uidA</i> and <i>nptII</i> gene expression in <i>C. sativa</i> <i>in vitro</i> regenerating shoots	197
5. Discussion	207
6. Conclusions	209
7. Statements	210
8. References	211
General Discussion	221
Conclusions	229
References (Introduction and General Discussion)	233