

CONTENTS

Introduction	1
Achieving cell diversity	3
An overview of the eukaryotic gene expression.	3
The making of a neuron.	16
The cilium - a sensory organelle	21
Basic morphological features of the cilium.	23
The intraflagellar transport machinery.	26
Key stages of ciliogenesis.	29
Functional diversity of sensory cilia.	31
When cilia fail: the ciliopathies.	33
<i>Caenorhabditis elegans</i> and its ciliated system	37
<i>C. elegans</i> as a model organism.	37
General architecture of the ciliated system.	39
Overview of sensory perception mediated by ciliated neurons.	47
Phenotypes & behaviours of ciliary mutants.	53
Transcriptional regulatory logic of cilium formation	55
The RFX family of transcription factors.	55
FOX genes, FOXJ1 and the regulation of the motile ciliogenic program.	61
Interplays between ciliary transcriptional networks	68
The making of specialised cilia	71
Motivation & Aims	77
Materials & Methods	81
Experimental procedures	83
<i>C. elegans</i> strains and maintenance.	83
Worm lysis for genomic DNA preparations.	83
Generation of fluorescent reporters: fusion PCR.	83
Generation of reporter transgenic lines.	85
Strain crossing.	89
Genotyping.	90
Microscopy & Scoring.	91
Bioinformatic analysis of regulatory sequences.	92
Identification of candidate transcription factors.	93
<i>In-vivo</i> analysis of transcription factors expression.	93
Dye-filling assays.	94
<i>fkh-8</i> ChIP-seq analysis.	95

Enrichment analysis.	95
Population synchronization.	95
RNA extraction.	96
<i>fkh-8</i> RNA-seq analysis.	97
CRISPR-based generation of a <i>fkh-8</i> null allele.	99
Rescuing experiments.	100
Site directed mutagenesis of <i>xbx-1</i> cis-regulatory region.	102
Behavioural assays.	103
Statistical analysis.	106
Materials	110
Results	111
1. A regulatory-motif enrichment analysis of the <i>C. elegans</i> ciliome.	113
Subsets of ciliated neurons maintain the expression of ciliary genes in the absence of <i>daf-19</i> .	113
Generation of a <i>bona-fide</i> list of ciliary genes.	117
<i>De novo</i> motif enrichment analysis of ciliary gene regulatory sequences.	122
RFX/DAF-19 binding sites are enriched in the promoters of structural ciliary genes.	122
Additional TF binding motifs are enriched in structural ciliary genes with RFX binding motifs.	129
2. FKH-8 is expressed in all sensory ciliated neurons of <i>C. elegans</i>.	136
Ten transcription factors are highly and specifically enriched within the ciliated sensory neurons of <i>C. elegans</i> .	136
FKH-8 is broadly expressed across the whole ciliated system of <i>C. elegans</i>	139
3. FKH-8 preferentially binds regulatory regions of ciliome genes	149
FKH-8 preferentially binds to ciliome genes when compared to the binding of other TFs.	149
Genomic binding analysis of FKH-8.	153
4. RNA-seq profiling of <i>fkh-8</i> and <i>daf-19</i> single and double mutants.	164
Downregulated genes in young adult <i>fkh-8(tm292)</i> mutants do not enrich ciliary terms or functions.	164
Downregulated genes in double <i>daf-12</i> and <i>daf-19</i> young adult mutants relate to neuronal processes.	174
Deregulated genes in double <i>daf-12</i> and <i>daf-19</i> young adult mutants with FKH-8 binding enrich ciliary features.	177
Innate immune response might be impaired in young adult <i>fkh-8(tm292)</i> mutants.	182

5. Generation of a new null mutant allele of <i>fkh-8</i>.	190
<i>fkh-8(tm292)</i> is likely a hypomorphic allele.	190
<i>fkh-8(vlc43)</i> : an engineered full deletion of the <i>fkh-8</i> locus.	191
<i>fkh-8</i> mutants do not exhibit dye-filling defects.	192
6. <i>In vivo</i> reporter gene expression analysis uncovers the role of FKH-8 regulating structural ciliary genes.	193
Animals with whole deletion of the <i>fkh-8</i> locus show stronger defects in the expression of cilium-related reporters.	193
<i>fkh-8</i> mutation affects the expression of structural ciliary genes in a cell-type specific manner.	196
<i>fkh-8</i> acts cell-autonomously.	199
<i>cis</i> -mutation of putative FKH binding sites strongly affects expression of the panciliary gene <i>xbx-1</i> .	199
7. <i>fkh-8</i> mutants display defects in a wide range of sensory-mediated behaviours.	206
<i>fkh-8</i> mutants show normal response to body touch.	206
<i>fkh-8</i> mutants display nose touch defects.	207
<i>fkh-8(vlc43)</i> mutants are defective in chemotaxis to a volatile attractant.	209
<i>fkh-8(vlc43)</i> mutants show defects in the avoidance response to a volatile repellent.	210
<i>fkh-8</i> mutants do not show chemotaxis defects to the water-soluble attractant NaCl.	212
<i>fkh-8</i> mutants display defects in the avoidance response to water-soluble toxic substances.	213
8. <i>fkh-8</i> & <i>daf-19</i> exhibit cross-regulation and synergistic effects.	215
<i>fkh-8(tm292)</i> mutation has no major effect over the expression of <i>daf-19</i> .	219
Long <i>daf-19</i> isoforms repress <i>fkh-8</i> expression in non-ciliated neurons.	220
<i>daf-19</i> and <i>fkh-8</i> act synergistically in the regulation of some structural ciliary genes.	224
Discussion	229
<i>daf-19</i> does not act alone in the regulation of ciliary features in <i>C. elegans</i> .	231
<i>In-silico</i> approaches help decipher transcriptional regulatory signatures.	232
Expression analyses allow for the identification of candidate regulators of the ciliated fate.	235
A multi-angle approach reveals <i>fkh-8</i> is a terminal selector of the ciliated fate in <i>C. elegans</i> .	236
<i>cis</i> -mutation of FKH sites suggests compensatory effects upon <i>fkh-8</i> loss.	242
Interplay between RFX and FKH members is also seen in <i>C. elegans</i> .	244

RNA-seq uncovers a role for <i>fkh-8</i> in the regulation of the innate immune response in <i>C. elegans</i> .	245
Concluding remarks.	247
Conclusions	251
Annexes	255
References	285