

Article

Fixed Point of Interpolative Rus–Reich–Ćirić Contraction Mapping on Rectangular Quasi-Partial b-Metric Space

Pragati Gautam ^{1,†}, Luis Manuel Sánchez Ruiz ^{2,*}  and Swapnil Verma ^{1,†} 

¹ Department of Mathematics, Kamala Nehru College, University of Delhi, New Delhi 110049, India; pgautam@knc.du.ac.in (P.G.); swapnil@knc.du.ac.in (S.V.)

² ETSID-Departamento de Matematica Aplicada & CITG, Universitat Politecnica de Valencia, E-46022 Valencia, Spain

* Correspondence: LMSR@mat.upv.es

† These authors contributed equally to this work.

Abstract: The purpose of this study is to introduce a new type of extended metric space, i.e., the rectangular quasi-partial b-metric space, which means a relaxation of the symmetry requirement of metric spaces, by including a real number s in the definition of the rectangular metric space defined by Branciari. Here, we obtain a fixed point theorem for interpolative Rus–Reich–Ćirić contraction mappings in the realm of rectangular quasi-partial b-metric spaces. Furthermore, an example is also illustrated to present the applicability of our result.

Keywords: fixed point; interpolation; Rus–Reich–Ćirić contraction; rectangular quasi-partial b-metric space

1. Introduction and Preliminaries

In the year 1968, Kannan [1] extended the well-known Banach contraction:

$$d(G\vartheta, G\eta) \leq \rho[d(\vartheta, G\vartheta) + d(\eta, G\eta)] \quad \text{for all } \vartheta, \eta \in M,$$

where $\rho \in [0, \frac{1}{2})$. In 2018, Karapinar [2] established the generalized Kannan-type contraction by using the interpolative approach and proved that such an interpolative Kannan-type contraction mapping owns a fixed point in a complete metric space. Let us recall that given a metric space (M, d) , a self-map $G: M \rightarrow M$ is called an interpolative Kannan-type contraction map, if:

$$d(G\vartheta, G\eta) \leq \rho[d(\vartheta, G\vartheta)]^\alpha \cdot [d(\eta, G\eta)]^{1-\alpha} \quad \text{for all } \vartheta, \eta \in M \setminus \text{Fix}(G)$$

where $\text{Fix}(G) = \{z \in M: Gz = z\}$. Recently, Karapinar/Agarwal/Aydi [3] introduced the following notion of interpolative Rus–Reich–Ćirić contractions in the context of partial metric spaces [4], which keep symmetry as one of their intrinsic properties.

Theorem 1 ([3]). *In the setting of partial metric space (M, d) , if a self-map $G: M \rightarrow M$ is an interpolative Rus–Reich–Ćirić-type contraction, i.e., there are constants $\rho \in [0, 1)$ and $\alpha, \beta \in (0, 1)$ such that:*

$$d(G\vartheta, G\eta) \leq \rho[d(\vartheta, \eta)]^\beta [d(\vartheta, G\vartheta)]^\alpha \cdot [d(\eta, G\eta)]^{1-\alpha-\beta} \quad \text{for all } \vartheta, \eta \in M \setminus \text{Fix}(G),$$

then G owns a fixed point.

In the year 2000, Branciari [5] introduced the notion of the rectangular metric space by replacing the triangle inequality with the quadrilateral inequality in the definition of the metric space. It was noticed by Suzuki [6] that the topological structure of the standard



Citation: Gautam, P.; Sánchez Ruiz, L.M.; Verma, S. Fixed Point of Interpolative Rus–Reich–Ćirić Contraction Mapping on Rectangular Quasi-Partial b-Metric Space. *Symmetry* **2021**, *13*, 32. <https://dx.doi.org/10.3390/sym13010032>

Received: 5 December 2020
Accepted: 24 December 2020
Published: 28 December 2020

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

metric space and that the rectangular metric space are not comparable. In 2019, Karapinar [7] defined the interpolative Rus–Reich–Ćirić contraction map on the rectangular metric. Interesting work has been done by several authors [8–19] enriching this research field. Firstly, we recall some basic definitions and concepts on the rectangular metric space.

Definition 1 ([5]). Let $M \neq \emptyset$. Consider $r: M \times M \rightarrow \mathbb{R}^+$ such that for all $\vartheta, \eta \in M$ and $u, v \in M$:

- (r₁) $r(\vartheta, \eta) = 0$ iff $\vartheta = \eta$ (identification),
 - (r₂) $r(\vartheta, \eta) = r(\eta, \vartheta)$ (symmetry),
 - (r₃) $r(\vartheta, \eta) \leq r(\vartheta, u) + r(u, v) + r(v, \eta)$ (quadrilateral inequality).
- (M, r) is called a rectangular metric space.

Example 1. Let $M = [1, \infty)$. Define $r(\vartheta, \eta) = |\ln(\vartheta/\eta)|$.

Here, $r(\vartheta/\eta) = 0 \implies \ln \vartheta = \ln \eta$, which gives $\vartheta = \eta$.

Again, $r(\vartheta/\eta) = r(\eta/\vartheta)$.

Furthermore, $r(\vartheta/\eta) \leq r(\vartheta, u) + r(u, v) + r(v, \eta)$.

As $|\ln(\vartheta/\eta)| = |\ln \vartheta - \ln \eta| \leq |\ln \vartheta - \ln u + \ln u - \ln v + \ln v - \ln \eta|$.

It can be observed that $|\ln(\frac{\vartheta}{\eta})| \leq |\ln \vartheta - \ln u| + |\ln u - \ln v| + |\ln v - \ln \eta|$.

Therefore, (r₃) holds. Thus, (M, r) is a rectangular metric space.

Definition 2 ([7]). Let (M, r) be a rectangular metric. Then:

- (i) A sequence $\{\vartheta_n\} \subset M$ converges to $\vartheta \in M$ if $r(\vartheta, \vartheta) = \lim_{n \rightarrow \infty} r(\vartheta, \vartheta_n)$.
- (ii) A sequence $\{\vartheta_n\} \subset M$ is called a Cauchy sequence if for every $\varepsilon > 0$, there exists a positive integer $N = N(\varepsilon)$ such that $r(\vartheta_n, \vartheta_m) < \varepsilon$ for all $n, m > N$.
- (iii) (M, r) is said to be complete if each Cauchy sequence in M is convergent.

Definition 3 ([7]). Let (M, r) be a rectangular metric space. We say that a mapping $G: M \rightarrow M$ is continuous at $u \in M$, if we have $G\vartheta_n \rightarrow Gu$ (in other words $\lim_{n \rightarrow \infty} r(G\vartheta_n, Gu) = 0$), for any sequence $\{\vartheta_n\}$ in M that is convergent to $u \in M$, that is $\vartheta_n \rightarrow u$.

Proposition 1 ([7]). Suppose that $\{\vartheta_n\}$ is a Cauchy sequence in a rectangular metric space such that $\lim_{n \rightarrow \infty} r(\vartheta_n, u) = \lim_{n \rightarrow \infty} r(\vartheta_n, z) = 0$, where $u, z \in M$. Then, $u = z$.

The following definition gives room for the lack of symmetry in the spaces under study. Let us recall that quasi-metric spaces [20] satisfy the same axioms as metric spaces, but with the requirement of symmetry.

Definition 4 ([21]). A quasi-partial b -metric on a non-empty set M is a function $qp_b: M \times M \rightarrow \mathbb{R}^+$ such that for some real number $s \geq 1$ and all $\vartheta, \eta, \zeta \in M$:

$$(QPb_1) \quad qp_b(\vartheta, \vartheta) = qp_b(\vartheta, \eta) = qp_b(\eta, \eta) \text{ implies } \vartheta = \eta,$$

$$(QPb_2) \quad qp_b(\vartheta, \vartheta) \leq qp_b(\vartheta, \eta),$$

$$(QPb_3) \quad qp_b(\vartheta, \vartheta) \leq qp_b(\eta, \vartheta),$$

$$(QPb_4) \quad qp_b(\vartheta, \eta) \leq s[qp_b(\vartheta, \zeta) + qp_b(\zeta, \eta)] - qp_b(\zeta, \zeta).$$

(M, qp_b) is called a quasi-partial b -metric space. The number s is called the coefficient of (M, qp_b) .

Lemma 1 ([22]). Let (M, qp_b) be a quasi-partial b -metric space. Then, the following hold:

- (i) if $qp_b(\vartheta, \eta) = 0$, then $\vartheta = \eta$.
- (ii) if $\vartheta = \eta$, then $qp_b(\vartheta, \eta) > 0$ and $qp_b(\eta, \vartheta) > 0$.

Definition 5 ([23]). Let (M, qp_b) be a quasi-partial b-metric. Then:

- (i) a sequence $\{\vartheta_n\} \subset M$ converges to $\vartheta \in M$ if $qp_b(\vartheta, \vartheta) = \lim_{n \rightarrow \infty} qp_b(\vartheta, \vartheta_n)$.
- (ii) a sequence $\{\vartheta_n\} \subset M$ is called a Cauchy sequence if $\lim_{n, m \rightarrow \infty} qp_b(\vartheta_n, \vartheta_m)$ exists.
- (iii) a quasi-partial b-metric space (M, qp_b) is said to be complete if every Cauchy sequence $\{\vartheta_n\} \subset M$ converges with respect to τ_{qp_b} to a point $\vartheta \in M$ such that:

$$qp_b(\vartheta, \vartheta) = \lim_{n, m \rightarrow \infty} qp_b(\vartheta_n, \vartheta_m).$$

- (iv) a mapping $f: M \rightarrow M$ is said to be continuous at $\vartheta_0 \in M$ if, for every $\varepsilon > 0$, there exists:

$$\delta > 0 \text{ such that } f(B(\vartheta_0, \delta)) \subset B(f(\vartheta_0), \varepsilon).$$

Lemma 2 ([23]). Let (M, qp_b) be a quasi-partial b-metric space and (M, dqp_b) be the corresponding b-metric space. Then, (M, dqp_b) is complete if (M, qp_b) is complete.

Lemma 3 ([24]). Let (M, qp_b) be a quasi-partial b-metric space and $G: M \rightarrow M$ be a given mapping. G is said to be sequentially continuous at $z \in M$ if for each sequence $\{\vartheta_n\}$ in M converging to z , we have: $G\vartheta_n \rightarrow Gz$, that is, $qp_b(G\vartheta_n, Gz) = qp_b(Gz, Gz)$.

2. Main Results

We start this section by introducing the notion of interpolative Rus–Reich–Ćirić-type contractions in the setting of the rectangular quasi-partial b-metric space.

Definition 6. A rectangular quasi-partial b-metric on a non-empty set M is a function $rqp_b: M \times M \rightarrow \mathbb{R}^+$ such that for some real number $s \geq 1$ and all $\vartheta, \eta, u, v \in M$:

- (RQPb₁) $rqp_b(\vartheta, \vartheta) = rqp_b(\vartheta, \eta) = rqp_b(\eta, \eta) \Rightarrow \vartheta = \eta$,
- (RQPb₂) $rqp_b(\vartheta, \vartheta) \leq rqp_b(\vartheta, \eta)$,
- (RQPb₃) $rqp_b(\vartheta, \vartheta) \leq rqp_b(\eta, \vartheta)$,
- (RQPb₄) $rqp_b(\vartheta, \eta) \leq s[rqp_b(\vartheta, u) + rqp_b(u, v) + rqp_b(v, \eta)] - rqp_b(u, u) - rqp_b(v, v)$.

(M, rqp_b) is called a rectangular quasi-partial b-metric space. The number s is called the coefficient of (M, rqp_b) .

Example 2. Let $M = [0, 1]$. Define $rqp_b(\vartheta, \eta) = |\vartheta - \eta| + \vartheta$.

Here, $rqp_b(\vartheta, \vartheta) = rqp_b(\vartheta, \eta) = rqp_b(\eta, \eta) \Rightarrow \vartheta = \eta$ as $\vartheta = |\vartheta - \eta| + \vartheta = \eta$ gives $\vartheta = \eta$.

Again, $rqp_b(\vartheta, \vartheta) \leq rqp_b(\vartheta, \eta)$ as $\vartheta \leq |\vartheta - \eta| + \vartheta$, and similarly, $rqp_b(\vartheta, \vartheta) \leq rqp_b(\eta, \vartheta)$ as $\vartheta \leq |\eta - \vartheta| + \eta$ for $0 \leq \vartheta \leq \eta$.

Furthermore, $rqp_b(\vartheta, \eta) + rqp_b(u, u) + rqp_b(v, v) \leq s[rqp_b(\vartheta, u) + rqp_b(u, v) + rqp_b(v, \eta)]$ as $|\vartheta - \eta| + \vartheta + u + v \leq [|\vartheta - u| + \vartheta + |u - v| + u + |v - \eta| + v]$.

It can be observed that

$$|\vartheta - \eta| + \vartheta + u + v = |\vartheta - u + u - v + v - \eta| + \vartheta + u + v \leq |\vartheta - u| + |u - v| + |v - \eta| + \vartheta + u + v.$$

Therefore, (RQPb₄) holds. Thus, (M, rqp_b) is a rectangular quasi-partial b-metric space with $s = 1$.

Example 3. Let $M = \mathbb{R}$. Define $rqp_b(\vartheta, \eta) = |\vartheta - \eta| + |\vartheta| + |\vartheta - \eta|^2$ for any $(\vartheta, \eta) \in M \times M$ with $s \geq 2$. Here, we can show that (M, rqp_b) is a rectangular quasi-partial b-metric space.

If $rqp_b(\vartheta, \vartheta) = rqp_b(\vartheta, \eta) = rqp_b(\eta, \eta) \Rightarrow \vartheta = \eta$.

Furthermore, $rqp_b(\vartheta, \vartheta) \leq rqp_b(\vartheta, \eta)$, which satisfies (RQPb₂).

Next, $rqp_b(\vartheta, \vartheta) = |\vartheta| \leq |\vartheta - \eta| + |\vartheta| + |\vartheta - \eta|^2$.

Since,

$$\begin{aligned} |\vartheta| - |\eta| &\leq |(|\vartheta| - |\eta|)| \\ &\leq |\vartheta - \eta| \\ &\leq |\vartheta - \eta| + |\vartheta - \eta|^2 \end{aligned}$$

which proves (RQPb₃).

Now, (RQPb₄) follows from,

$$\begin{aligned} rqp_b(\vartheta, \eta) + rqp_b(u, u) + rqp_b(v, v) &= |\vartheta - \eta| + |\vartheta| + |\vartheta - \eta|^2 + |u| + |v| \\ &\leq 2(|\vartheta - u| + |u - v| + |v - \eta| + |v - u|^2 + |u - v|^2 + |v - \eta|^2) \\ &\leq s[rqp_b(\vartheta, u) + rqp_b(u, v) + rqp_b(v, \eta)]. \end{aligned}$$

Definition 7. Let (M, rqp_b) be a rectangular quasi-partial b -metric space. A self-mapping G on M is called an interpolative Rus–Reich–Ćirić-type contraction, if there are $\rho \in [0, 1)$ and positive reals α, β with $\alpha + \beta < 1$ such that:

$$rqp_b(G\vartheta, G\eta) \leq \rho[rqp_b(\vartheta, \eta)]^\beta \cdot [rqp_b(\vartheta, G\vartheta)]^\alpha \cdot \left[\frac{1}{s}rqp_b(\eta, G\eta)\right]^{1-\alpha-\beta} \quad (1)$$

for all $\vartheta, \eta \in M \setminus \text{Fix}(G)$.

Theorem 2. Let $G: M \rightarrow M$ be an interpolative Rus–Reich–Ćirić-type contraction on a complete rectangular quasi-partial b -metric space (M, rqp_b) , then G has a fixed point in M .

Proof. Let ϑ_0 be an arbitrary point in M . Consider ϑ_n by $\vartheta_n = G^n(\vartheta_0)$ for each positive integer n . If there exists n_0 such that $\vartheta_{n_0} = \vartheta_{n_0+1}$, then ϑ_{n_0} is a fixed point of G , and we are done. Throughout the proof, we assume that $\vartheta_n \neq \vartheta_{n+1}$ for each $n \geq 0$.

We shall prove that $\lim_{n \rightarrow \infty} rqp_b(\mu_n, \mu_{n+1}) = 0$.

By substituting the values $\vartheta = \vartheta_n$ and $\eta = \mu_{n-1}$, we find that:

$$\begin{aligned} rqp_b(\vartheta_{n+1}, \vartheta_n) &= rqp_b(G\vartheta_n, G\vartheta_{n-1}) \\ &\leq \rho[rqp_b(\vartheta_n, \vartheta_{n-1})]^\beta \cdot [rqp_b(\vartheta_n, G\vartheta_n)]^\alpha \left[\frac{1}{s}rqp_b(\vartheta_{n-1}, \vartheta_{n-1})\right]^{1-\alpha-\beta} \\ &\leq \rho[rqp_b(\vartheta_n, \vartheta_{n-1})]^\beta \cdot [rqp_b(\vartheta_n, \vartheta_{n+1})]^\alpha \left[\frac{1}{s}rqp_b(\vartheta_{n-1}, \vartheta_n)\right]^{1-\alpha-\beta} \\ &\leq [rqp_b(\vartheta_{n-1}, \vartheta_n)]^{1-\alpha} \cdot [rqp_b(\vartheta_n, \vartheta_{n+1})]^\alpha. \end{aligned} \quad (2)$$

We get,

$$[rqp_b(\vartheta_n, \vartheta_{n+1})]^{1-\alpha} \leq \rho[rqp_b(\vartheta_{n-1}, \vartheta_n)]^{1-\alpha}. \quad (3)$$

Therefore, we conclude that,

$$rqp_b(\vartheta_n, \vartheta_{n+1}) \leq \rho rqp_b(\vartheta_{n-1}, \vartheta_n) \text{ for all } n \geq 1. \quad (4)$$

That is, $\{rqp_b(\vartheta_{n+1}, \vartheta_n)\}$ is a non-increasing sequence with non-negative terms. Eventually, there is a non-negative constant L such that $\lim_{n \rightarrow \infty} rqp_b(\vartheta_{n+1}, \vartheta_n) = L$. Note that $L \geq 0$. Indeed, from (4), we deduce that:

$$rqp_b(\vartheta_n, \vartheta_{m+1}) \leq \rho rqp_b(\vartheta_{n+1}, \vartheta_n) \leq \rho^n rqp_b(\vartheta_0, \vartheta_1). \quad (5)$$

Since $\rho < 1$ and by taking $n \rightarrow \infty$ in the inequality (5), we deduce that $L = 0$.

Now, we shall show that

$$\lim_{n \rightarrow \infty} rqp_b(\vartheta_n, \vartheta_{n+2}) = 0$$

Using (4) and (5) and the quadrilateral inequality, we have

$$\begin{aligned} rqp_b(\vartheta_{n+2}, \vartheta_n) &= rqp_b(G\vartheta_{n+1}, G\vartheta_{n-1}) \\ &\leq \rho[rqp_b(\vartheta_{n+1}, \vartheta_{n-1})]^\beta [rqp_b(\vartheta_{n+1}, G\vartheta_{n+1})]^\alpha \left[\frac{1}{s} rqp_b(\vartheta_{n-1}, G\vartheta_{n-1})\right]^{1-\alpha-\beta} \\ &\leq \rho[rqp_b(\vartheta_{n+1}, \vartheta_{n-1})]^\beta [rqp_b(\vartheta_{n+1}, \vartheta_{n+2})]^\alpha [rqp_b(\vartheta_{n-1}, \vartheta_n)]^{1-\alpha-\beta} \\ &\leq \rho[rqp_b(\vartheta_{n+1}, \vartheta_{n-1})]^\beta [rqp_b(\vartheta_n, \vartheta_{n+1})]^\alpha [rqp_b(\vartheta_{n-1}, \vartheta_n)]^{1-\alpha-\beta} \\ &\leq \rho[rqp_b(\vartheta_{n+1}, \vartheta_{n-1})]^\beta [rqp_b(\vartheta_{n-1}, \vartheta_n)]^{1-\beta} \\ &\leq \rho[srqp_b(\vartheta_{n+1}, \vartheta_{n+2}) + srqp_b(\vartheta_{n+2}, \vartheta_n) + srqp_b(\vartheta_n, \vartheta_{n-1})]^\beta [rqp_b(\vartheta_{n+1}, \vartheta_n)]^{1-\beta} \\ &\leq \rho[srqp_b(\vartheta_{n+2}, \vartheta_n) + 2rsqp_b(\vartheta_n, \vartheta_{n-1})]^\beta [srqp_b(\vartheta_{n+2}, \vartheta_n) + 2rsqp_b(\vartheta_n, \vartheta_{n-1})]^{1-\beta} \\ &\leq \rho[s(rqp_b(\vartheta_{n+2}, \vartheta_n) + 2rqp_b(\vartheta_n, \vartheta_{n-1}))] \\ &\leq \rho srqp_b(\vartheta_{n+2}, \vartheta_n) + 2\rho srqp_b(\vartheta_n, \vartheta_{n-1}) \end{aligned} \quad (6)$$

Therefore,

$$rqp_b(\vartheta_{n+2}, \vartheta_n) \leq \frac{2\rho s}{1-\rho s} rqp_b(\vartheta_n, \vartheta_{n-1}), \text{ for all } n \geq 1.$$

We shall prove that $\{\vartheta_n\}$ is a Cauchy sequence, that is $\lim_{n \rightarrow \infty} rqp_b(\vartheta_n, \vartheta_{n+p}) = 0$ for all $p \in \mathbb{N}$.

Case 1. Let $p = 2m$. By the quadrilateral inequality, we find:

$$\begin{aligned} rqp_b(\vartheta_n, \vartheta_{n+2m}) &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2}) + rqp_b(\vartheta_{n+2}, \vartheta_{n+2m})] \\ &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2})] + s^2[rqp_b(\vartheta_{n+2}, \vartheta_{n+3}) + rqp_b(\vartheta_{n+3}, \vartheta_{n+4}) + \\ &\quad rqp_b(\vartheta_{n+4}, \vartheta_{n+2m})] \\ &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2})] + s^2[rqp_b(\vartheta_{n+2}, \vartheta_{n+3}) + rqp_b(\vartheta_{n+3}, \vartheta_{n+4})] \\ &\quad + s^3[rqp_b(\vartheta_{n+4}, \vartheta_{n+5}) + rqp_b(\vartheta_{n+5}, \vartheta_{n+6}) + rqp_b(\vartheta_{n+6}, \vartheta_{n+2m})] \\ &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2})] + \dots + s^{m-1}[rqp_b(\vartheta_{2m-4}, \vartheta_{2m-3}) \\ &\quad + rqp_b(\vartheta_{2m-3}, \vartheta_{2m-2})] + s^{m-1}[rqp_b(\vartheta_{n+2m-2}, \vartheta_{n+2m})] \\ &\leq s[\rho^n rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+1} rqp_b(\vartheta_0, \vartheta_1)] + s^2[\rho^{n+2} rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+3} rqp_b(\vartheta_0, \vartheta_1)] + \dots + \\ &\quad s^3[\rho^{n+4} rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+5} rqp_b(\vartheta_0, \vartheta_1)] + \dots + s^{m-1}[\rho^{2m-4} rqp_b(\vartheta_0, \vartheta_1) + \rho^{2m-3} rqp_b \\ &\quad (\vartheta_0, \vartheta_1)] + s^{m-1} \rho^{n+2m+2} rqp_b(\vartheta_0, \vartheta_1) \\ &\leq s\rho^n [1 + s\rho^2 + s^2\rho^4 + \dots] rqp_b(\vartheta_0, \vartheta_1) + s\rho^{n+1} [1 + s\rho^2 + s^2\rho^4 + \dots] rqp_b(\vartheta_0, \vartheta_1) \\ &\quad + s^{m-1} \rho^{n+2m-2} rqp_b(\vartheta_0, \vartheta_1) \\ rqp_b(\vartheta_n, \vartheta_{n+2m}) &\leq \frac{1+\rho}{1-s\rho^2} s\rho^n rqp_b(\vartheta_0, \vartheta_1) + (s\rho)^{2m} \rho^{n-2} rqp_b(\vartheta_0, \vartheta_1) \\ &\leq \frac{1+\rho}{1-s\rho^2} s\rho^n rqp_b(\vartheta_0, \vartheta_1) + \rho^{n-2} rqp_b(\vartheta_0, \vartheta_1) \end{aligned} \quad (7)$$

Case 2. Let $p = 2m + 1$.

$$\begin{aligned}
 rqp_b(\vartheta_n, \vartheta_{n+2m+1}) &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2}) + (rqp_b(\vartheta_{n+2}, \vartheta_{n+2m+1}))] \\
 &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2})] + s^2[rqp_b(\vartheta_{n+2}, \vartheta_{n+3}) + rqp_b(\vartheta_{n+3}, \vartheta_{n+4}) + \\
 &\quad rqp_b(\vartheta_{n+4}, \vartheta_{n+2m+1})] \\
 &\leq s[rqp_b(\vartheta_n, \vartheta_{n+1}) + rqp_b(\vartheta_{n+1}, \vartheta_{n+2})] + s^2[rqp_b(\vartheta_{n+2}, \vartheta_{n+3}) + rqp_b(\vartheta_{n+3}, \vartheta_{n+4})] + \dots + \\
 &\quad s^m rqp_b(\vartheta_{n+2m}, \vartheta_{n+2m+1}) \\
 &\leq s[\rho^n rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+1} rqp_b(\vartheta_0, \vartheta_1)] + s^2[\rho^{n+2} rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+3} rqp_b(\vartheta_0, \vartheta_1)] + \dots \\
 &\quad + s^3[\rho^{n+4} rqp_b(\vartheta_0, \vartheta_1) + \rho^{n+5} rqp_b(\vartheta_0, \vartheta_1)] + \dots + s^m \rho^{n+2m} rqp_b(\vartheta_0, \vartheta_1) \\
 &\leq s\rho^n [1 + s\rho + s^2\rho^4 + \dots] rqp_b(\vartheta_0, \vartheta_1) + s\rho^{n+1} [1 + s\rho^2 + s^2\rho^4 + \dots] rqp_b(\vartheta_0, \vartheta_1) \\
 &= \frac{1 + \rho}{1 - s\rho^2} s\rho^n rqp_b(\vartheta_0, \vartheta_1)
 \end{aligned}$$

Therefore,

$$rqp_b(\vartheta_n, \vartheta_{n+2m+1}) \leq \frac{1 + \rho}{1 - s\rho^2} s\rho^n rqp_b(\vartheta_0, \vartheta_1). \tag{8}$$

By (7) and (8), $\lim_{n \rightarrow \infty} rqp_b(\vartheta_n, \vartheta_{n+p}) = 0$.

Thus, $\{\vartheta_n\}$ is a Cauchy sequence. Since (M, rqp_b) is complete, there exists $z \in M$ such that $\lim_{n \rightarrow \infty} \vartheta_n = z$. Next, we shall prove that z is a fixed point of G .

Let $\vartheta = \vartheta_n$ and $\eta = z$,

$$\begin{aligned}
 rqp_b(\vartheta_{n+1}, Gz) &= rqp_b(G\vartheta_n, Gz) \\
 &\leq [rqp_b(\vartheta_n, z)]^\beta \cdot [rqp_b(\vartheta_n, G\vartheta_n)]^\alpha \cdot \left[\frac{1}{s} rqp_b(z, Gz)\right]^{1-\alpha-\beta}. \tag{9}
 \end{aligned}$$

Letting $n \rightarrow \infty$ in (9), we conclude $\lim_{n \rightarrow \infty} rqp_b(\vartheta_n, Gz) = 0$.

By Proposition 1, we get $Gz = z$. \square

Example 4. Let $M = \{0, 1, 2, 3\}$. Consider the complete rectangular quasi-partial b -metric as $rqp_b(\vartheta, \eta) = |\vartheta - \eta| + \vartheta$, that is:

$rqp_b(\vartheta, \eta)$	0	1	2	3
0	0	1	2	3
1	2	1	2	3
2	4	3	2	3
3	6	5	4	3

We define a self-map G on M as $G: \begin{pmatrix} 0 & 1 & 2 & 3 \\ 0 & 0 & 1 & 0 \end{pmatrix}$ as shown in Figure 1.

Choose $\alpha = \frac{1}{2}$, $\beta = \frac{1}{3}$, and $\rho = \frac{6}{10}$.

Case 1: Let $(\vartheta, \eta) = (1, 1)$. We have:

$$\begin{aligned}
 rqp_b(G\vartheta, G\eta) &\leq \rho[rqp_b(\mu, \eta)]^\beta [rqp_b(\vartheta, G\vartheta)]^\alpha \left[\frac{1}{s} rqp_b(\eta, G\eta)\right]^{1-\alpha-\beta} \\
 rqp_b(G1, G1) &= 0 \leq \rho[rqp_b(1, 1)]^{1/3} [rqp_b(1, G1)]^{1/2} \left[\frac{1}{s} rqp_b(1, G1)\right]^{1/6}
 \end{aligned}$$

Case 2: Let $(\vartheta, \eta) = (3, 3)$:

$$rqp_b(G3, G3) = 0 \leq \rho[rqp_b(3, 3)]^{1/3} [rqp_b(3, G3)]^{1/2} \left[\frac{1}{s} rqp_b(3, G3)\right]^{1/6}.$$

Thus, zero is the fixed point of G in the setting of the interpolative Rus–Reich–Ćirić-type contraction.




Figure 1. Zero is the fixed point of G .

Problem 1. Let (M, rqp_b) be a complete rectangular quasi-partial b -metric space. Consider a family of self-mappings $G_n: M \rightarrow M$, $n \geq 1$, and $s \geq 1$ such that

$$rqp_b(G_i\vartheta, G_j\eta) \leq \rho_{i,j} [rqp_b(\vartheta, \eta)]^{\beta_j} \cdot [rqp_b(\vartheta, G_i\vartheta)]^{\alpha_i} \cdot \left[\frac{1}{s} rqp_b(\eta, G_j\eta) \right]^{1-\alpha_i-\beta_j}.$$

What are the conditions on $\rho_{i,j}, \alpha_i, \beta_j$ for G_n to have a fixed point?

Definition 8. Let (M, rqp_b) be a rectangular quasi-partial b -metric space. A self-mapping G on M is called an interpolative Kannan contraction, if there are $\rho \in [0, 1)$ and positive reals $\alpha \in [0, 1)$ such that

$$rqp_b(G\vartheta, G\eta) \leq \rho [rqp_b(\vartheta, G\vartheta)]^\alpha \cdot \left[\frac{1}{s} rqp_b(\eta, G\eta) \right]^{1-\alpha}$$

for all $\vartheta, \eta \in M \setminus \text{Fix}(G)$.

Theorem 3. Let $G: M \rightarrow M$ be an interpolative Kannan contraction on a complete rectangular quasi-partial b -metric space (M, rqp_b) , then G has a fixed point in M .

We skip this proof as it is similar to Theorem 2.

3. Conclusions

In the present study, the authors investigated the interpolative Rus–Reich–Ćirić contraction mapping and its variants to attain the fixed point on a new metric space known as the rectangular quasi-partial b -metric space. Interpolation in fixed point theory is an advanced and widespread technique, which is acknowledged in several research areas such as metallurgy, earth sciences, and surface physics, etc., due to its application potential in the approximation of signal sensation analysis. The present research will find its place in these applications. Determining the fixed point for a non-self mapping and fractal interpolants will be an interesting work for future study.

Author Contributions: Conceptualization, P.G., L.M.S.R. and S.V.; investigation, P.G., L.M.S.R. and S.V.; methodology, P.G., L.M.S.R. and S.V.; writing—review editing, P.G., L.M.S.R. and S.V. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not Applicable.

Acknowledgments: The authors would like to thank the anonymous referees for their careful reading, valuable comments, and suggestions, which helped to improve the manuscript.

Conflicts of Interest: The authors declare that they have no competing interests.

References

1. Kannan, R. Some results on fixed points. *Bull. Calcutta Math. Soc.* **1968**, *60*, 71–76.
2. Karapınar, E. Revisiting the Kannan type contractions via interpolation. *Adv. Theory Nonlinear Anal. Appl.* **2018**, *2*, 85–87. [[CrossRef](#)]
3. Karapınar, E.; Agarwal, R.P.; Aydi, H. Interpolative Reich–Rus–Ćirić type contractions on partial-metric spaces. *Mathematics* **2018**, *6*, 256. [[CrossRef](#)]
4. Matthews, S.G. Partial metric topology. *Ann. N. Y. Acad. Sci.* **1994**, *728*, 183–197. [[CrossRef](#)]
5. Branciari, A. A fixed point theorem of Banach-Caccioppoli type on a class of generalized metric spaces. *Publ. Math. Debrecen* **2000**, *57*, 31–37.
6. Suzuki, T. Generalized metric space do not have the compatible topology. *Abstr. Appl. Anal.* **2014**, *2014*, 5. [[CrossRef](#)]
7. Aydi, H.; Chen, C.M.; Karapınar, E. Interpolative Reich–Rus–Ćirić Type Contractions via the Branciari Distance. *Mathematics* **2019**, *7*, 84. [[CrossRef](#)]
8. Karapınar, E.; Alqahtani, O.; Aydi, H. On interpolative Hardy–Rogers type contractions. *Symmetry* **2018**, *11*, 8. [[CrossRef](#)]
9. Ćirić, L.B. Generalized contractions and fixed-point theorems. *Publ. Inst. Math. (Beograd)* **1971**, *12*, 19–26.
10. Reich, S. Fixed point of contractive functions. *Boll. Un. Mat. Ital.* **1972**, *4*, 26–42.
11. Reich, S. Kannan’s fixed point theorem. *Boll. Un. Mat. Ital.* **1971**, *4*, 1–11.
12. Reich, S. Some remarks concerning contraction mappings. *Can. Math. Bull.* **1971**, *14*, 121–124. [[CrossRef](#)]
13. Hardy, G.E.; Rogers, T.D. A generalization of a fixed point theorem of Reich. *Can. Math. Bull.* **1973**, *16*, 201–206. [[CrossRef](#)]
14. Karapınar, E.; Chi, K.P.; Thanh, T.D. A generalization of Ćirić quasi-contractions. *Abstr. Appl. Anal.* **2012**. [[CrossRef](#)]
15. Mlaiki, N.; Abodayeh, K.; Aydi, H.; Abdeljawad, T.; Abuloha, M. Rectangular metric-like type spaces and related fixed points. *J. Math.* **2018**. [[CrossRef](#)]
16. Krein, S.G.; Petunin, J.I.; Semenov, E.M. *Interpolation of Linear Operators*; American Mathematical Society: Providence, RI, USA, 1978.
17. Aydi, H.; Karapınar, E.; Hierro, A.F.R. ω -Interpolative Reich–Rus–Ćirić type contractions. *Mathematics* **2019**, *7*, 57. [[CrossRef](#)]
18. Debnath, P.; de La Sen, M. Set-valued interpolative Hardy–Rogers and set-valued Reich–Rus–Ćirić-type contractions in b-metric spaces. *Mathematics* **2019**, *7*, 849. [[CrossRef](#)]
19. Karapınar, E.; Agarwal, R.P. Interpolative Reich–Rus–Ćirić type contractions via simulation functions. *Mathematics* **2018**, *6*, 256. [[CrossRef](#)]
20. Wilson, W.A. On quasi-metric spaces. *Am. J. Math.* **1931**, *53*, 675–684. [[CrossRef](#)]
21. Gautam, P.; Mishra, V.N.; Ali, R.; Verma, S. Interpolative Chatterjea and cyclic Chatterjea contraction on quasi-partial b-metric space. *AIMS Math.* **2021**, *6*, 1727–1742. [[CrossRef](#)]
22. Gupta, A.; Gautam, P. Quasi partial b-metric spaces and some related fixed point theorems. *Fixed Point Theory Appl.* **2015**, *18*, 1–12. [[CrossRef](#)]
23. Gupta, A.; Gautam, P. Some coupled fixed point theorems on quasi-partial b-metric spaces. *Int. J. Math. Anal.* **2015**, *9*, 293–306. [[CrossRef](#)]
24. Mishra, V.N.; Sánchez Ruiz, L.M.; Gautam, P.; Verma, S. Interpolative Reich–Rus–Ćirić and Hardy–Rogers Contraction on Quasi-Partial b-Metric Space and Related Fixed Point Results. *Mathematics* **2020**, *8*, 1598. [[CrossRef](#)]