

REVIEW PAPER

Genetic insights into the modification of the pre-fertilization mechanisms during plant domestication

Silvia Manrique¹, James Friel^{2,3}, Pietro Gramazio^{4,5}, Tomas Hasing³, Ignacio Ezquer¹ and Aureliano Bombarely^{1,3,*}

¹ Department of Biosciences, Università degli Studi di Milano, Milan, Italy

² Genetics and Biotechnology Laboratory, Plant and AgriBioscience Research Center (PABC), Ryan Institute, National University of Ireland Galway, Galway, Ireland

³ School of Plant and Environmental Sciences (SPES), Virginia Tech, Blacksburg, VA, USA

⁴ Instituto de Conservación y Mejora de la Agrodiversidad Valenciana (COMAV), Universitat Politècnica de València, Valencia, Spain

⁵ Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

* Correspondence: aurebg@vt.edu

Received 25 December 2018; Editorial decision 2 May 2019; Accepted 2 May 2019

Editor: Sílvia Coimbra, University of Porto, Portugal

Abstract

Plant domestication is the process of adapting plants to human use by selecting specific traits. The selection process often involves the modification of some components of the plant reproductive mechanisms. Allelic variants of genes associated with flowering time, vernalization, and the circadian clock are responsible for the adaptation of crops, such as rice, maize, barley, wheat, and tomato, to non-native latitudes. Modifications in the plant architecture and branching have been selected for higher yields and easier harvests. These phenotypes are often produced by alterations in the regulation of the transition of shoot apical meristems to inflorescences, and then to floral meristems. Floral homeotic mutants are responsible for popular double-flower phenotypes in Japanese cherries, roses, camellias, and lilies. The rise of peloric flowers in ornamentals such as snapdragon and florists' gloxinia is associated with non-functional alleles that control the relative expansion of lateral and ventral petals. Mechanisms to force outcrossing such as self-incompatibility have been removed in some tree crops cultivars such as almonds and peaches. In this review, we revisit some of these important concepts from the plant domestication perspective, focusing on four topics related to the pre-fertilization mechanisms: flowering time, inflorescence architecture, flower development, and pre-fertilization self-incompatibility mechanisms.

Keywords: Clonal propagation, domestication, flower development, flowering time, inflorescence architecture, self-incompatibility, sexual reproduction.

Introduction

Plant domestication is the process of adapting plants to human use by selecting specific traits. Domestication can also be understood as the process of selection of crucial traits by early farmers (domestication syndrome), being different from crop improvement, a later process where secondary traits are selected.

For this review, we have preferred to use domestication as a synonym of human-driven active selection of useful (or desirable) traits. Under this definition, domestication can be understood as a continuous process ranging from the active growing of a wild plant with a specific goal (pre-domestication) to their

genetic modification by modern techniques such as CRISPR (clustered regularly interspaced short palindromic repeats; molecular breeding).

Many traits selected during domestication are related to plant reproduction, since most plant-derived food is the product of plant reproduction (seeds and fruits). Plant reproduction is frequently altered in the case of ornamental plants too, as often the targets of the selection are the flowers.

Charles Darwin was one of the first scientists to study the phenotypic changes related to the domestication process. The first chapter of his book '*On the origin of species*' introduced several ideas about domestication such as the increase in phenotypic diversity and the pushing of the reproductive barriers during domestication (Darwin, 1859). Then, 9 years later in his book '*The variation of animals and plants under domestication*', Darwin developed his ideas and observations about domestication in greater detail. Chapters X and XI summarize Darwin's observations on the variation of flowers, buds, and reproduction modes (Darwin, 1868). He described floral homeotic mutations in which stamens and pistils are converted to petals for species such as *Aquilegia vulgaris* (columbine) and *Primula vulgaris* (hose-in-hose primroses), respectively. He also mentioned a poppy variety in which stamens have turned into pistils. DeVries (1904) also shared this observation in his book '*Species and varieties, their origin by mutation*' where he pointed to this phenotype in the species *Papaver commutatum*. Another change in flower morphology described by Charles Darwin was the transition from zygomorphic to actinomorphic (peloric) flowers in *Sinningia speciosa* (gloxinia) and *Antirrhinum majus* (snapdragon). Darwin's observations exemplify some of the changes in the reproductive mechanisms which occurred during the plant domestication process, but they are not the only ones. Changes in the transition from the vegetative to the reproductive phase, the fertilization process, fruit development, ripening, and abscission are also commonly associated with plant domestication. In this review, we present important examples of the alteration of these mechanisms. Due to space constraints, we will focus on the events occurring prior to the fertilization of the ovule: transition of the vegetative to the reproductive phase, inflorescence architecture and flower development, and self-incompatibility (SI). We summarize the main genes described in this article in Table 1.

Changes in flowering time associated with plant domestication

The transition from the vegetative to the reproductive stage is controlled by a complex genetic mechanism that translates changes in photoperiod, temperature, and plant hormones into the signal that induces the production of flowers. The plant domestication process involved the adaptation of human-selected populations to environments with different photoperiods and temperatures, which involves the selection of changes in the flowering time of these species. The genetic mechanisms that control flowering time have been extensively studied in several plant species, but most of the work has been done in the model species *Arabidopsis thaliana*, a long-day plant native to Africa

and Eurasia. Hundreds of genes have been described in the flowering time pathway, but, due to space constraints, we will focus on listing genes that have been selected during domestication without giving extensive detail about the pathways, as this has already been reviewed in many excellent articles.

The central player controlling flowering time is the *FLOWERING LOCUS T* gene (*FT*), which encodes a small phosphatidylethanolamide-binding protein (PEBP) that binds to phospholipids (Kobayashi *et al.*, 1999). *FT* is expressed in leaves and is induced by long-day treatment. *FT* is translocated to the shoot apex where it induces its own expression and activates the expression of floral determination genes to trigger flowering (Wigge *et al.*, 2005; Corbesier *et al.*, 2007).

Photoperiod is the most important environmental signal determining flowering time. In Arabidopsis, photoperiod information is connected to *FT* through the *CONSTANS* gene (*CO*), a zinc finger transcription activator expressed in leaves that activates the expression of *FT* (Putterill *et al.*, 1995; Kardailsky *et al.*, 1999). The *CONSTANS* (*CO*) gene is post-transcriptionally regulated by *GIGANTEA* (*GI*), a circadian clock gene (Park *et al.*, 1999; Huq *et al.*, 2000). The *FT* protein moves to the shoot apical meristem (SAM) and activates the expression of floral determination genes to trigger flowering (Wigge *et al.*, 2005; Corbesier *et al.*, 2007). However, *FT* is tightly regulated in order to integrate other inputs, such as light quality or temperature, to ensure that flowering aligns with seasonal cues for successful reproduction (Valverde *et al.*, 2004; Song *et al.*, 2015). Additionally, vernalization and temperature are also important factors in regulating the expression of *FT* in plants that require a period of cold before flowering such as the winter-annual ecotypes of Arabidopsis (Michaels and Amasino, 1999).

The flowering time pathways in other species share some similarities with Arabidopsis, but also have distinctive features (Zhu *et al.*, 2017). In the short-day monocot *Oryza sativa*, *Heading date 3a* (*Hd3a*) and *FLOWERING LOCUS T 1* (*RFT1*) represent the orthologs of the Arabidopsis *FT* gene (Komiya *et al.*, 2008). *Heading date 1* (*Hd1*), the homolog of *CO* in rice, is regulated by *OsGI* (the *GI* homolog in rice) (Yano *et al.*, 2000; Hayama *et al.*, 2003). *Hd1* promotes flowering by activating *Hd3a* expression under short-day conditions and delays flowering by repressing *Hd3a* expression under long-day conditions (Yano *et al.*, 2000; Kojima *et al.*, 2002; Komiya *et al.*, 2008). The *Early heading date 1* (*Ehd1*) pathway, which is unique to grasses and independent of *Hd1* (Doi *et al.*, 2004), also promotes the expression of *FLOWERING LOCUS T 1* (*RFT1*) and *Hd3a* (Doi *et al.*, 2004; Xue *et al.*, 2008; Itoh *et al.*, 2010).

A very important domestication trait related to flowering time is the ability to flower at latitudes different from the plant's native region. Therefore, specific alleles of several important players in the flowering time pathway have been positively selected during domestication to achieve this.

Cultivated rice, *O. sativa* L., is a good example of a plant originally from a tropical region that has been adapted to a wide range of latitudes, from 53°N to 40°S. The adaptation of rice to high latitudes was driven by the selection of natural variants of several genes from both *Hd1* and *Ehd1* pathways. Loss-of-function alleles of *Hd1* itself (Goretti *et al.*, 2017) or

Table 1. Summary of the genes described in this article

Gene ID	Arabidopsis homolog	Species	Molecular function	Pathway
HEADING DATE 3A (<i>hd3A</i> , Os06t0157700)	FLOWERING LOCUS (<i>FT</i> , AT1G65480)	<i>Oryza sativa</i>	Phosphatidylethanolamine binding (GO:0008429)	Flowering time
RICE FLOWERING-LOCUS T 1 (<i>RFT1</i> , Os06t0157500)	FLOWERING LOCUS (<i>FT</i> , AT1G65480)	<i>Oryza sativa</i>	Phosphatidylethanolamine binding (GO:0008429)	Flowering time
HEADING DATE 7 (<i>Ghd7</i> , Os07g0261200)	NA	<i>Oryza sativa</i>	DNA binding (GO:0003677), protein binding (GO:000551)	Flowering time
Heading date (<i>QTL</i>)-5(<i>t</i>) (<i>Hd5</i> , Os08g0174500)	NA	<i>Oryza sativa</i>	DNA binding (GO:0003677), protein binding (GO:0005515)	Flowering time
<i>Oryza sativa</i> Pseudo-Response Regulator37 (OsPRR37, Os07g0695100)	Two-component response regulator-like (<i>APRR7</i> , AT5G02810)	<i>Oryza sativa</i>	Protein binding (GO:0005515)	Flowering Time
HEADING DATE 1 (<i>Hd1</i> , Os06t0275000)	Pseudo-response regulator 3 (<i>PRR3</i> , AT5G60100)	<i>Oryza sativa</i>	Protein binding (GO:0005515), zinc ion binding (GO:0008270)	Flowering time
<i>Zea mays</i> CCT transcription factor 9 (<i>ZmCCT9</i>)	CONSTANS (<i>CO</i> , AT5G15840)	<i>Zea mays</i> ssp. <i>mays</i>	DNA binding (GO:0003677), protein binding (GO:000551)	Photoperiod
<i>Glycine max</i> Cryptochrome 1 (<i>GmCRY1a</i> , Glyma04G101500)	CRYPTOCROME 2 (<i>CRY2</i> , AT1G04400)	<i>Glycine max</i>	Protein binding (GO:000551), nucleotide binding (GO:0000166)	Photoperiod
<i>Glycine max</i> CONSTANS-like 7a (<i>GMCOL7a</i>)	CONSTANS (<i>CO</i> , AT5G15840)	<i>Glycine max</i>	Protein binding (GO:0005515), zinc ion binding (GO:0008270)	Flowering time
<i>GmGla</i> (Glyma10g36600)	Protein GIGANTEA (AT1G22770)	<i>Glycine max</i>	Protein binding (GO:0005515)	Flowering time
<i>Glycine max</i> phosphatidylethanolamine-binding protein <i>FT2a</i> (<i>GmFT2a</i> , Glyma16G150700)	FLOWERING LOCUS T (<i>FT</i> , AT1G65480)	<i>Glycine max</i>	Phosphatidylethanolamine binding (GO:0008429)	Flowering time
<i>Helianthus annuus</i> flowering locus T4 (<i>HaFT4</i> , 110873663)	FLOWERING LOCUS T (<i>FT</i> , AT1G65480)	<i>Helianthus annuus</i>	Phosphatidylethanolamine binding (GO:0008429)	Flowering time
<i>Solyc09g075080</i>	Phytochrome A-associated F-box protein (<i>EID1</i> , AT4G02440)	<i>Solanum lycopersicum</i>	Protein binding (GO:0005515), ubiquitin-protein ligase activity (GO:0004842)	Flowering time
<i>Solyc01g068560</i>	NIGHT LIGHT-INDUCIBLE AND CLOCK-REGULATED 2 (<i>LNK2</i> , AT3G54500)	<i>Solanum lycopersicum</i>		Photoperiod
Teosinte branched 1 (<i>tb1</i> , Zm00001d033673)	TEOSINTE BRANCHED 1 (<i>TCP24</i> , AT1G30210)	<i>Zea mays</i> ssp. <i>mays</i>	DNA binding (GO:0003677), DNA-binding transcription factor activity (GO:0003700)	Inflorescence architecture
ABERRANT PANICLE ORGANIZATION (Os06g0665400)	Floral meristem identity control protein <i>LEAFY</i> (<i>LFY</i> , AT5G61850)	<i>Oryza sativa</i>	Ubiquitin-protein transferase activity (GO:0004842), protein binding (GO:0005515)	Flower development
Putative phosphatidylethanolamine-binding protein <i>TFL1a</i> (<i>GmTFL1</i> , Glyma03G194700)	TERMINAL FLOWER 1 (<i>TFL1</i> , At5g03840)	<i>Glycine max</i>	Transcription co-regulator activity (GO:0003712)	Flower development
Self-pruning (<i>sp</i> , Solyc06g074350)	CENTRORADIALIS (<i>ATC</i> , AT2G27550)	<i>Solanum lycopersicum</i>	Phosphatidylethanolamine binding (GO: 0008429)	Flower development
<i>Brassica oleracea</i> Transcription factor <i>CAULIFLOWER</i> (<i>BoCAL</i> , 106320120)	Transcription factor <i>CAULIFLOWER</i> (<i>CAL</i> , At1g26310)	<i>Brassica oleracea</i>	Protein binding (GO:0005515), DNA-binding transcription factor activity (GO:0003700)	Flower development
<i>Falsiflora</i> (<i>fa</i> , Solyc03g118160)	Floral meristem identity control protein <i>LEAFY</i> (<i>LFY</i> , AT5G61850)	<i>Solanum lycopersicum</i>	Transcription factor binding (GO:0008134)	Flowering time, flower development
COMPOUND INFLORESCENCE (<i>S</i> , Solyc02g077390)	WUSCHEL HOMEODOMAIN 9 (<i>WOX9</i> , AT2G33880)	<i>Solanum lycopersicum</i>	DNA binding (GO:0003677)	Inflorescence architecture
LIGULELESS 1 (OsLG1, Os04g0656500)	Squamosa promoter-binding-like protein 8 (<i>SPL8</i> , AT1G02065)	<i>Oryza sativa</i>	DNA binding (GO:0003677), metal ion binding (GO:0046872)	Flower development
CYCLOIDEA (<i>CYC</i> , O49250)	TEOSINTE BRANCHED 1 (<i>TCP24</i> , AT1G30210)	<i>Antirrhinum majus</i>	DNA binding (GO:0003677), DNA-binding transcription factor activity (GO:0003700)	Flower development

Table 1. Continued

Gene ID	Arabidopsis homolog	Species	Molecular function	Pathway
DICHOTOMA (DICH, Q0GFJ4)	NA	<i>Antirrhinum majus</i>	DNA binding (GO:0003677)	Flower development
DIVARICATA (DIV, Q8S9H7)	NA	<i>Antirrhinum majus</i>	DNA binding (GO:0003677)	Flower development
RADIALIS (RAD, Q58FS3)	NA	<i>Antirrhinum majus</i>	DNA binding (GO:0003677)	Flower development

the repressors of *Ehd1*, *Grain number, plant height and heading date 7 (Ghd7)* (Xue *et al.*, 2008) and *Days To Heading on chromosome 8 (DTH8/Ghd8/OsHAP3H/Hd5)* (Xue *et al.*, 2008; Wei *et al.*, 2010; Fujino *et al.*, 2013) were selected to obtain plants with low photoperiod sensitivity. Selection of natural variation of *OsPRR37*, a pseudo-response regulator (PRR) gene that makes up part of the circadian clock, also contributed to the adaptation of rice to cultivation at higher latitudes (Koo *et al.*, 2013).

In maize, the adaptation to different geographical regions follows a similar pattern to that of rice. Maize (*Zea mays* ssp. *mays*) and its ancestor teosinte are native to tropical South-western Mexico. The adaptation of maize to higher latitudes is linked with the down-regulation of the gene *ZmCCT9* (homolog of the rice gene *Ghd7*) by the insertion of a Harbinger-like transposon in a distant regulative region, which results in photoperiod insensitivity, allowing flowering in long-day conditions (Huang *et al.*, 2018).

Conversely, the domestication of wheat and barley followed different trajectories. While some varieties were selected as winter crops, with day-neutral behaviour and a strong vernalization requirement, other varieties were selected as short-seasoned spring varieties not requiring vernalization (Blümel *et al.*, 2015). However, in both cases, most known variations are associated with changes in *VERNALIZATION 1 (VRN1)*, a MADS-box transcription factor gene involved in the vernalization process that promotes inflorescence initiation (Fjellheim *et al.*, 2014).

In the case of soybean (*Glycine max*), adaptation to other latitudes is associated with a strong selection of the genes *GmCRY1a* (homolog of the Arabidopsis blue light receptor *CRYPTOCROME 2*) and *GMCOL7a* (homolog of *CO*) (Li *et al.*, 2013). Soybean quantitative trait locus (QTL) analysis also linked four alleles (E1, E2, E3, and E4) to flowering time (Zhai *et al.*, 2014). The major contributor, E2, is an allele of the *GmGla* gene, the soybean homeolog of *GI* of Arabidopsis. The dominant E2 allele carries an early stop codon mutation that leads to induction of the expression of the *GmFT2a* gene, producing early flowering (Langewisch *et al.*, 2014). Another case in which FT is affected is sunflower (*Helianthus annuus* L.), where five members of the *FT* gene family were selected during domestication, including a non-functionalized copy (*HaFT4*) (Blackman *et al.*, 2011).

Other interesting cases are the mutations selected in the tomato genes *Solyc09g075080*, the homolog of the Arabidopsis F-box *EID1* that functions as a negative regulator in phytochrome A-specific light signaling, and *Solyc01g068560*, a homolog of the Arabidopsis *NIGHT LIGHT-INDUCIBLE*

AND CLOCK-REGULATED 2 (LNK2) that functions in the integration of light signaling and the circadian clock. They are responsible for the adaptation of this crop to the long summers in temperate regions. A single amino acid deletion in the *Solyc09g075080/EID1* protein delays the circadian phase (Müller *et al.*, 2016). For the *Solyc01g068560/LNK2* gene, an almost complete deletion lengthens the circadian period (Müller *et al.*, 2018).

Modifications in inflorescence architecture during plant domestication

The architecture of the inflorescence conditions the number of flowers and, as a consequence, fruits and seeds that are produced and their position on the plant (Wyatt, 1982; Evers *et al.*, 2011; Iwata *et al.*, 2012; Teo *et al.*, 2014). Therefore, genes controlling inflorescence development are instrumental for domestication as they have a profound impact on key agronomical aspects such as yield and crop management (Benlloch *et al.*, 2015).

Inflorescence architecture is determined by two main factors: the growth habit of the plant and the level of indeterminacy of the inflorescence meristem (IM). Regarding growth habit, in monopodial plants, such as Arabidopsis or rice, due to apical dominance, vertical growth results only from the SAM. In these plants, after floral transition, the main SAM develops into the leader inflorescence shoot with subordinate branches. Instead, in sympodial plants such as tomato, the role of the SAM is sequentially adopted by the uppermost axillary meristem, which, after a period of growth, will either terminate in reproductive structures or abort. Then, the growth will continue from a new axillary meristem that will repeat this pattern so, instead of a leader inflorescence, several inflorescences of similar size are formed along the shoot.

One of the best-known examples of genes modifying inflorescence architecture related to domestication is *TEOSINTE-BRANCHED 1 (TB1)* of maize/teosinte. *teosinte branched1 (tb1)* was identified as a major QTL contributing to the shift towards monopodial growth habit with a concomitant increase in ear size during teosinte domestication (Doebley *et al.*, 1997). *tb1* encodes a member of the TCP family of transcriptional regulators expressed in the axillary meristems (Doebley *et al.*, 1997; Hubbard *et al.*, 2002; Kebrom and Brutnell, 2015). The TB1 protein acts as a repressor of organ growth and contributes to apical dominance by repressing branch outgrowth. The maize allele of *TB1* is more highly expressed than that of teosinte, causing greater repression of branching compared with

teosinte (Doebley *et al.*, 1997; Studer *et al.*, 2011). Increased expression of *TB1* in maize is due to an insertion of a transposable element 65 kb upstream of the *TB1* coding region (Studer *et al.*, 2011; Kebrom and Brutnell, 2015).

The balance between the maintenance of indeterminacy or commitment to flower is the main determinant of the architecture of the inflorescence. Upon perception of inducing environmental and/or internal cues, the SAM transitions from a vegetative to a reproductive identity, becoming the primary IM (I_1) can either produce flowers or remain indeterminate to produce branch meristems (I_2), which iterate the pattern of I_1 (Prusinkiewicz *et al.*, 2007; Teo *et al.*, 2014). Based on the activity of the I_1 and the I_2 , inflorescences can be classified into four groups: when the I_1 terminates as a flower, they are determinate inflorescences; in contrast, when the I_1 grows indefinitely until senescence to produce a floral meristem (FM) or I_2 , they are indeterminate inflorescences. Additionally, when the I_2 forms the FM, we are in the presence of simple inflorescences. Alternatively, if the I_2 forms further IMs, increasing the complexity of the architecture, we refer to them as compound inflorescences (Weberling, 1989; Benlloch *et al.*, 2007, 2015; Cheng *et al.*, 2018). A basic genetic model to explain how the balance between IM and FM identity is determined has been developed using studies in the model plant *A. thaliana*. Although the applicability of this model to other species is variable, it is useful to set a frame for comparison.

The development of the Arabidopsis inflorescence can be mostly explained by the function of three genes: *TERMINAL FLOWER 1* (*TFL1*), *LEAFY* (*LFY*), and *APETALA 1* (*AP1*) (Shannon and Meeks-Wagner, 1993; Liljegren *et al.*, 1999; Blázquez *et al.*, 2006; Benlloch *et al.*, 2015). These genes coordinate to maintain the balance between IM and FM identity at the inflorescence apex. Broadly speaking, *TFL1* promotes IM identity, while *LFY* and *AP1* promote FM identity. Therefore, it has been proposed that differences in their expression patterns or function can explain much of the diversity of inflorescence architectures observed among angiosperms (Ratcliffe *et al.*, 1999; Blázquez *et al.*, 2006; Benlloch *et al.*, 2007; Serrano-Mislata *et al.*, 2017). Briefly, *TFL1*, a PEBP, is specifically expressed in the center of the I_1 and I_2 , and promotes IM identity by repressing *LFY* and its direct target *AP1* (and its paralog *CAULIFLOWER*, *CAL*) to prevent early inflorescence termination (Mandel *et al.*, 1992; Weigel *et al.*, 1992; Weigel and Nilsson, 1995; Parcy *et al.*, 1998; Teo *et al.*, 2014). Conversely, *LFY*, a plant-specific transcription factor gene, and *AP1* and *CAL*, two paralog MADS-box transcription factor genes, are expressed in the lateral FM primordia produced by the IM. The joint action of *LFY* and *AP1/CAL* in the newly formed FM leads to the repression of *TFL1*, allowing the up-regulation of floral organ identity genes and leading to the formation of flowers (Parcy *et al.*, 1998; Liljegren *et al.*, 1999; Wagner *et al.*, 1999; Kaufmann *et al.*, 2009). However, the mechanism leading to *TFL1* repression in the FM is not linear. Recent works pointed out that, actually, *LFY* activates *TFL1* in the FM while *AP1/CAL* represses it (Goslin *et al.*, 2017; Serrano-Mislata *et al.*, 2017). This indicates that *LFY* and *AP1* might be part of a feed-forward loop that could serve to ensure that

flower development starts only when *AP1/CAL* levels are high enough to over-ride *LFY* inhibitory action, ensuring that the conditions for stable development of flowers are already established (Goslin *et al.*, 2017). These results imply that *LFY* might also be involved in maintaining the indeterminate growth of the IM. Indeed, the activity of the *TFL1* promoter is reduced in *lfy* mutants (Serrano-Mislata *et al.*, 2017). Although *LFY* is not expressed in the SAM, *LFY* protein is mobile (Sessions *et al.*, 2000) so it can travel to the IM and bind to the *TFL1* promoter. Therefore, the relationship between *TFL1* and *LFY* is not entirely antagonistic as previously thought.

Regarding the situation in other plants, of the three genes forming the model, *TFL1* is the most conserved. *TFL1* orthologs exist in most land plants, and investigations in various plant species have shown that their role is mostly conserved (Chardon and Damerval, 2005; Carmona *et al.*, 2007; Danilevskaya *et al.*, 2010; C. Liu *et al.*, 2013; Teo *et al.*, 2014; Mahrez *et al.*, 2016). Mutations of *TFL1* produce a switch to more determinate inflorescences. Examples of crops where this has been selected during domestication are soybean and tomato (Wang *et al.*, 2018). *Glycine soja*, the wild progenitor of soybean (*G. max*), is indeterminate. Instead, many cultivated soya varieties have a determinate growth habit. This trait was found to be controlled by the *Dt1* locus encoded by *GmTFL1* (Liu *et al.*, 2010). As soybean contains several *TFL1* paralogs, complementation of the Arabidopsis *tfl1* mutant with *GmTFL1* demonstrated that it was the functional *TFL1* ortholog (Tian *et al.*, 2010).

Instead, the presence and role of *LFY* and *AP1* vary among species. For example, the rice ortholog of *LFY*, *ABERRANT PANICLE ORGANIZATION 2/RICE FLORICAULA*, is not expressed in FMs and, in contrast to Arabidopsis, it has a role in suppressing the transition from IM to FM (Kyojuka *et al.*, 1998; Ikeda-Kawakatsu *et al.*, 2012).

Another example is the tomato *self-pruning* (*sp*) mutant. *sp* was discovered 90 years ago and has facilitated the transformation of indeterminate tomato plants into new determinate forms (Yeager, 1927; Pnueli *et al.*, 1998; Wang *et al.*, 2018), leading to a more compact phenotype and synchronized growth which is adequate for mechanical harvesting (McGarry *et al.*, 2016). For this reason, the *SP* mutation was rapidly bred into all industrial tomatoes. However, it must be noted that although the product of *SP* is a PEBP and the functional equivalent of that of *TFL1* in tomato, the real ortholog of *SP* in Arabidopsis is another PEBP-encoding gene, *Arabidopsis thaliana* *CENTRORADIALIS* (*ATC*) (Mimida *et al.*, 2001; McGarry *et al.*, 2016).

Obvious examples of inflorescence architecture re-shaping associated with domestication are the mutants of *Brassica oleracea*, cauliflower (*B. oleracea* ssp. *botrytis*), and broccoli (*B. oleracea* ssp. *italica*). The cauliflower head is composed of a hypertrophied mass of IMs and FMs. In broccoli, developmental arrest happens at a later stage, so although the inflorescence also develops into a large hypertrophied structure, flower buds are eventually formed (Carr and Irish, 1997; Schilling *et al.*, 2018). Upon the characterization of the *ap1-1/cal-1* mutant of Arabidopsis, the similarity of broccoli and cauliflower with it led to speculation that the *AP1* and *CAL* orthologs

from *B. oleracea* might be responsible for this phenotype (Smith and King, 2000). Surprisingly, the link is not as clear as initially thought and, at present, the basis of these phenotypes is still not completely clear. Molecular and population genetic studies indicate that the function of the *B. oleracea* *CAL* (*BoCAL*) is compromised in both varieties (Kempin *et al.*, 1995; Lowman and Purugganan, 1999; Purugganan *et al.*, 2000; Smith and King, 2000). The situation for *AP1*-like genes is less clear, since several copies of *AP1*-like genes exist in *B. oleracea* (Lowman and Purugganan, 1999) and, although they are associated with the phenotype, they do not explain it entirely, indicating that additional genes might be involved (Labate *et al.*, 2006; Duclos and Björkman, 2008; Schilling *et al.*, 2018).

Besides *TFL1*, *LFY*, and *AP1*, many other genes involved in the network that regulates inflorescence architecture have been targeted during domestication. For example, although mutations in *FALSIFLORA* (*FA*), the ortholog of *LFY* in tomato, cause an increase in inflorescence branching (Moliner-Rosales *et al.*, 1999; Zheng *et al.*, 2017), most commercial highly branched tomato varieties carry mutations on *COMPOUND INFLORESCENCE* (*S*). *S* encodes a homolog of *WUSCHEL HOMEODOMAIN 9* (*WOX9*) of Arabidopsis. *WOX9* in Arabidopsis is involved in SAM and root apical meristem (RAM) maintenance (Wu *et al.*, 2005, 2007) and embryo patterning (Haecker *et al.*, 2004; Wu *et al.*, 2007; Ueda *et al.*, 2011), and has no effect on inflorescence branching in Arabidopsis, a difference probably originating from the different growth habits (Lippman *et al.*, 2008). Another example is the mutation on the regulatory regions of *OsLG1* or *OsSPL8*, encoding a squamosa promoter-binding-like protein which is responsible for the switch from a spread panicle in wild rice to the compact panicle of domesticated rice (Zhu *et al.*, 2013).

Floral crop selection as an example of alterations in flower development pathways

Perfect flowers contain four types of organs arranged in concentric rings known as whorls. From the outermost to innermost whorl, the organ types are: sepals, petals, stamens, and carpels. A combinatorial model that explains how these four organ types are specified within the FM was proposed in the early 1990s, based on the observation of a series of homeotic mutants in Arabidopsis and *Antirrhinum* (Causier *et al.*, 2010; Moyroud and Glover, 2017). Presently known as the ABCDE model, the model was originally proposed as the ABC model and extended later on (Coen and Meyerowitz, 1991; Weigel and Meyerowitz, 1994; Causier *et al.*, 2010).

The model proposes that five functions named A, B, C, D, and E specify which organs form in each whorl of the flower. A+E genes specify sepals, A+B+E specify petals, B+C+E specify stamens, C+E specify carpels, and C+D+E specify ovules, and the ABCDE genes are sufficient to superimpose floral organ identity in vegetative organs of angiosperms (Parcy *et al.*, 1998; Honma and Goto, 2001; Pelaz *et al.*, 2001). Additionally, it was observed that C-function expands into the outer whorls in A-function mutants and vice versa, so mutual repression between

the A- and C-functions was integrated into the model to explain it (Causier *et al.*, 2010). In Arabidopsis, the genes responsible for A-function are *APETALA1* (*AP1*) and *APETALA2* (*AP2*), B-function is encoded by *APETALA3* (*AP3*) and *PISTILLATA* (*PI*), and C-function is encoded by *AGAMOUS* (*AG*) (Yanofsky *et al.*, 1990; Jack *et al.*, 1992; Mandel *et al.*, 1992; Goto and Meyerowitz, 1994; Jofuku *et al.*, 1994; Theißen *et al.*, 2016). Three paralog MADS-box genes are responsible for the D-function: *SEEDSTICK* (*STK*), *SHATTERPROOF1* (*SHP1*), and *SHATTERPROOF2* (*SHP2*) (Favaro *et al.*, 2003; Pinyopich *et al.*, 2003), and E-function is redundantly encoded by *SEPALLATA* genes (*SEP1–SEP4*) (Pelaz *et al.*, 2000, 2001; Ditta *et al.*, 2004). Except for *AP2*, which is an AP2/EREBP (ethylene-responsive element-binding protein), all the other ABCDE genes encode MADS-box transcription factors. MADS-domain proteins can interact with each other, forming tetramers, explaining the combinatorial nature of the model: each organ is determined by a specific tetrameric combination of floral identity MADS-box proteins (Theißen and Saedler, 2001; Theißen *et al.*, 2016). This is known as the floral quartet model (Stewart *et al.*, 2016).

Although the model is 30 years old, the conceptual framework continues to be broadly valid. Floral identity genes are also present in gymnosperms, where a 'BC' model has been proposed (Baum and Hileman, 2006; Theissen and Melzer, 2007; Chanderbali *et al.*, 2016), in which C-function is expressed in male and female cones while B-function is restricted to male cones (Irish, 2017). In this context, the ABCDE model can be seen as an evolutionary extension of the 'BC' model. Instead, A-function has always been controversial as it seems to be much less conserved than the others and, in the last years, evidence has been accumulating pointing to the fact that outside Arabidopsis and close relatives, a classical A-function is rare (Litt, 2007; Ye *et al.*, 2016; Morel *et al.*, 2017; Wils and Kaufmann, 2017; Schilling *et al.*, 2018). However, a recent study has shown that, in rice, A-function exists and it is performed by *OsMADS14* and *OsMADS15* genes which belong to the AP1/FUL clade (Wu *et al.*, 2017), so the debate is still open.

The other functions are quite well conserved across angiosperms, particularly eudicots and monocots, although, for example, in petunia C- and D-function cannot be strictly distinguished from each other (Heijmans *et al.*, 2012; Theißen *et al.*, 2016; Schilling *et al.*, 2018). Finally, some differences that do not fit in a whorl-based ABCDE model are observed in some basal angiosperms. However, those differences can be explained as modifications of the ABCDE model like the 'fading borders model' (Causier *et al.*, 2010; Chanderbali *et al.*, 2016; Wils and Kaufmann, 2017).

Regarding modification of flower morphology due to domestication, probably the most frequent and notorious examples are the alterations in *AG* that result in double flowers where stamens and carpels are replaced by petals (Bowman *et al.*, 1989; Schilling *et al.*, 2018). Since the determinacy of the meristem is also disturbed, this pattern is iterated multiple times, leading to flowers with very high numbers of petals (Bowman *et al.*, 1989). In some cases, double flowers are associated with loss-of-function mutations in *AG*-like genes, such as, for example, in rue-anemone (*Thalictrum thalictroides*) or

Japanese cherry (*Prunus lannesiana*) (Galimba *et al.*, 2012; Z. Liu *et al.*, 2013; Schilling *et al.*, 2018). However, the formation of double flowers in rose, *Camellia*, or lily is instead associated with a restriction of the expression domain of AG-like genes toward the center of the meristem (Dubois *et al.*, 2010; Akita *et al.*, 2011; Sun *et al.*, 2014). The underlying molecular cause of the shrinkage of the AG expression domain was unknown, but recent data from rose and peach indicate that it might be caused by mutations in euAP2 genes, which are known to repress AG in many species (François *et al.*, 2018; Gattolin *et al.*, 2018). It seems that in both cases the mutations responsible for the change in the AG expression pattern were caused by the loss of the *miR172*-binding site in the euAP2 gene (François *et al.*, 2018; Gattolin *et al.*, 2018).

Besides floral organs, the symmetry of flowers is a characteristic often modified during domestication, especially in ornamental crops. Flowers can have two types of symmetry: radial or actinomorphy and bilateral or zygomorphy (Endress, 1999; Krizek and Fletcher, 2005; Smyth, 2018). Zygomorphy is thought to have evolved many times from an ancestral actinomorphic condition as a strategy apparently associated with the attraction of pollinating insects with bilateral vision (Krizek and Fletcher, 2005; Smyth, 2018). Although the classification is binary, the outcome is not, as differential development of each whorl can give rise to several intermediate situations. For example, while orchids are the classical example of zygomorphy and show a strong bilateral symmetry spanning all whorls, some species of *Solanum* present a restricted zygomorphy affecting only the stamen whorl, related to their interaction with pollen-collecting bees (Glover *et al.*, 2004).

Snapdragon (*A. majus*) is the classic genetic model in which zygomorphy has been studied. Its symmetry is based on the expansion of the dorsal petals relative to the lateral and ventral ones and abortion of the dorsal stamen (Krizek and Fletcher, 2005). Dorsal identity is specified by two paralogous TCP-domain family transcription factor genes with overlapping functions, *CYCLOIDEA* (*CYC*) and *DICHOTOMA* (*DICH*). Ventral identity is specified by the MYB-domain transcription factor gene *DIVARICATA* (*DIV*), expressed all across the flower. Expression of *CYC* and *DICH* in the dorsal domain activates *RADIALIS* (*RAD*), which encodes a protein with a single MYB domain. *RAD* antagonizes *DIV* function in dorsal cells and limits its activity to the lateral and ventral domains by competing for *DIV* interaction partners, the *DIV*-AND-*RAD*-INTERACTING-FACTORS (DRIFs), which are required for *DIV* activity in specifying ventral symmetry (Luo *et al.*, 1996, 1999; Galego and Almeida, 2002; Corley *et al.*, 2005; Perez-Rodriguez *et al.*, 2005; Gao *et al.*, 2017). Thus, *rad* and *cyc dich* double mutants are ventralized and have a radially symmetrical appearance. Many regressions to actinomorphic symmetry, both in natural populations (Cubas *et al.*, 1999; Reardon *et al.*, 2009) and in domesticated crops, are caused by mutations affecting *CYC* expression. A recent example of zygomorphic to actinomorphic reversion in a domesticated plant caused by a mutation in a *CYC* ortholog are gloxinias (*Sinningia speciosa*), where the loss of *CYC* function is caused by a 10 bp deletion in the coding sequence of the gene (Dong *et al.*, 2018).

CYC-like genes have also been associated with asymmetric pigmentation in zygomorphic flowers. For example, *TfCYC2* in wishbone flower (*Torenia fournieri*) evolved regulatory loops to bind to the promoter region of an R2R3-MYB factor gene repressing its transcription, which under normal circumstances promotes anthocyanin-related pigmentation in the epidermal cells of petals (Su *et al.*, 2017).

Limitations in plant breeding driven by pre-fertilization self-incompatibility

Plant fitness/yield has been fundamentally changed during domestication by hybridization, genetic bottlenecks, alteration of reproductive strategies, and polyploidization. Such changes have greatly modulated current plant traits in agriculture. Angiosperms exhibit a wide array of reproductive strategies, both asexual and sexual, with sexual reproduction including self-fertilization and cross-fertilization strategies. In unstable or unpredictable environments, reproductive strategies promoting cross-fertilization are fundamental to evolutionary success as they contribute to the creation of genetically diverse populations which increase the probability that at least one individual in a population will survive under changing conditions. However, in situations of low presence of sexual partners, or in stable and predictable environments, an asexual strategy and self-fertilization are effective means of reproduction that can be favored. Numerous wild plants display efficient mechanisms that ensure outcrossing promoting high levels of plant heterozygosity. SI is reported in >100 families and distributed among an estimated 39% of species (Ilg and Kohn, 2006). During angiosperm evolution, different molecular mechanisms for promoting SI have evolved at least 35 times (Iwano and Takayama, 2012). Self-compatibility (SC), on the other hand, might have evolved to adapt to conditions such as the loss of pollinators (Gervasi and Schiestl, 2017).

Perennial species are generally outcrossers, while annuals are more tolerant to SC. Since annuals only have one chance to reproduce during their lifespan, for them it might be better to self-pollinate, rather than not reproducing at all. Conversely, perennial species can wait longer and avoid the detrimental effects of inbreeding over time (Pekkala *et al.*, 2014).

Outcrossing can be achieved either through the spatiotemporal separation of the sexes via (hetero) dichogamy or dioecy, or by SI (Miller and Gross, 2011). SI is a genetically controlled mechanism that induces a higher sexual selection by preventing self-fertilization in wild-type plants. In some cases, SI arises from floral morphology (heteromorphology) thanks to genetically controlled physical or temporal barriers that prevent self-pollination. This is the case for primrose (*Primula*) which exhibits two floral forms (morphs) that differ in morphology, primarily in the relative placement of stigmas and anthers, and pollinations succeed only between different morphs (De Nettancourt, 2001). Multiple varieties of heteromorphic SI systems are present in the plant kingdom (seen in *Passifloraceae*, *Lythraceae*, *Polygonaceae*, and *Primulaceae*) which are considered to have evolved independently (Fujii *et al.*, 2016).

Nevertheless, most SI systems in plants belong to homomorphic systems where incompatibility is achieved by one-to-one interaction of two or a few genes. SI was originally classified into two types: (i) gametophytic SI (GSI) and (ii) sporophytic SI (SSI), based on the genetic control of the SI phenotype by pollen. A late-acting SI system (LSI) has been also described in several species recently (see review by Gibbs, 2014).

GSI has been described in *Rosaceae*, *Plantaginaceae*, *Papaveraceae*, and *Solanaceae*. According to the phylogeny and the conserved structure of the female S-RNase gene, it seems that this gene evolved only once, before the separation of the *Asterideae* and *Rosidaeae*, ~120 million years ago (Vieira *et al.*, 2009). GSI has a common molecular basis across many plant families and is probably the ancestral condition for flowering plants (McClure, 2006). In contrast, SSI is present in at least 10 plant families and derives from at least 17 distinct evolutionary origins (Ilgic *et al.*, 2008; Koseva *et al.*, 2017). In SSI, S-specificity is determined by the genotype of the sporophyte that produced the pollen grain (Sehgal and Singh, 2018). SSI has been most deeply characterized in *Brassicaceae* (Hiscock and Tabah, 2003; Kitashiba and Nasrallah, 2014; Lao *et al.*, 2014; Iwano *et al.*, 2015; Baldwin and Schoen, 2017). More recently, substantial advances have been made in describing SSI in *Asteraceae* (Gonthier *et al.*, 2013; Koseva *et al.*, 2017).

Finally, in LSI, both compatible and incompatible pollen grains can reach the ovary with a similar rate of pollen tube growth. However, whilst the double fertilization is completed after 24 h in compatible reactions, in incompatible reactions the male nuclei are released into the embryo sac, but they fail to achieve gamete fusion, resulting in floral abscission. LSI cases range from *Malvaceae*, *Apocynaceae*, and *Bignoniaceae*, to the monocot families *Amaryllidaceae* and *Xanthorrhoeaceae* (Gibbs, 2014). A well-characterized LSI case is *Theobroma cacao* (da Silva *et al.*, 2016; Lanaud *et al.*, 2017). Most of the *T. cacao* accessions are self incompatible; however, some anciently domesticated varieties, such as 'Criollo' varieties from Central America originally cultivated by the Mayas, or 'Comun' from Brazil and 'Nacional' from Ecuador, are self-compatible (Loor Solorzano *et al.*, 2012; Santos *et al.*, 2015).

During domestication, SI has been introduced, or removed, according to agronomic parameters of interest (McClure, 2012). For example, in some fruit trees of the *Rosaceae* family, such as apple, Japanese pear, sweet cherry, or apricot, trees of different cross-compatible varieties should be planted to ensure fruit production due to SI (Sassa, 2016). Similar situations have been reported for other crops such as cabbage, chicory, or sugar beet (Ockendon, 1974; Broothaerts, 2003; Wünsch and Hormaza, 2004; Hunt *et al.*, 2010; Gonthier *et al.*, 2013; Sassa, 2016; Saumitou-Laprade *et al.*, 2017; Farinelli *et al.*, 2018). An interesting example is the genus *Prunus*, a large genus in the *Rosaceae* family, that includes multiple domesticated crops such as almond, apricot, cherry, peach, and plum. Most of the *Prunus* species exhibit S-RNase-based gametophytic SI. In some species such as almond, domestication goals were exclusively focused on improving organoleptic aspects such as reduced toxicity, thinner endocarp, and increased seed size. In

contrast, in other cases, such as peach, domestication was focused not only on improving aspects affecting fruit morphology but also on the introduction of SC (Miller and Gross, 2011). Other crops of the genus *Prunus* such as cherries and plums also have had SC introduced during their domestication (Spiegel-Roy, 1986). Modern breeding programs oriented towards disrupting SI in almond only began recently (Martínez-Gómez *et al.*, 2006).

Other perennial species have evolved SC under domestication. For instance, wild grapevine is dioecious while the domesticated relative is hermaphrodite and self-compatible (De Mattia *et al.*, 2008). In contrast to perennials where few crops derive from selfing wild populations, many annual crops have been domesticated from SC wild ancestors, such as barley, chickpea, eggplant, lentils, pea, chile, tomatoes, and wheat (Miller and Gross, 2011).

However, SI can also be a desirable trait for breeders. SI systems prevent self-fertilization, which forces outcrossing and increases genetic diversity, which is useful for the breeding of hybrid varieties of economically important plant families. Accordingly, breeding programs towards introducing functional SI have been activated in many crops (Kaothien-Nakayama *et al.*, 2010; Havlíčková *et al.*, 2014; Cheng *et al.*, 2018; Xiao *et al.*, 2019). In order to control mating, many advances have been made toward understanding the SI mechanisms. However, transferring these mechanisms across wide phylogenetic distances is often difficult, or even impossible, for breeders. Recently, de Graaf and co-workers introduced an SI system (from *Papaver*) in a species with no SI system that diverged ~140 million years ago (de Graaf *et al.*, 2012), demonstrating that this transfer may be easier than previously thought.

Asexually (clonally) propagated crops where sexual reproduction is reduced have also been promoted by breeders. Vegetative propagation might be preferred to sexual reproduction either to avoid the segregation of traits in SI species (McKey *et al.*, 2010) or to speed up breeding and growing cycles in perennials with long juvenile phases. For this reason, many tree crops such as avocado or olive trees are propagated clonally (Diez *et al.*, 2015; Kuhn *et al.*, 2019).

An interesting example of a herbaceous plant that is clonally propagated is banana. The banana domestication involved hybridizations between diverse species and subspecies that generated diploid and triploid sterile hybrids. Nonetheless, the hybrids are able to produce parthenocarpic fruits that have been thereafter dispersed by vegetative propagation (D'Hont *et al.*, 2012).

Clonally propagated crops can potentially produce a wider range of adaptations with respect to sexual reproductive families that are easily maintained, but genetic homogeneity is an important drawback for survival if adverse conditions arise. For example, half of the 2018 banana world production relies on somaclones derived from a single triploid genotype (Cavendish) (Lescot, 2010). Since pests and diseases have gradually become adapted, at present this genetic homogeneity represents an imminent danger for global banana production (de Bellaire *et al.*, 2010; Dita *et al.*, 2010).

Final remarks

Many of the changes selected during domestication are related to reproductive traits, either because the outcome of the reproductive process (seeds or fruits) is the desired result, or because the alteration of the reproductive process is necessary to achieve it (i.e. to be able to cross varieties/species or to avoid undesirable phenotype variation). In this review, we have summarized some of the changes associated with molecular mechanisms that human action have introduced into the reproductive structures and strategies of domesticated plants. Darwin already noticed many of the changes in reproductive structures and strategies introduced by domestication, but he lacked the tools to understand their basis. At present, many domestication-driven changes in plant reproduction are still not well understood because crop molecular biology research has developed more slowly than that based on model plants. Recently developed tools such as genome editing and next-generation sequencing are changing this scenario and quickly increasing our knowledge of the molecular basis of domestication in crop species. This will also allow for better planning of breeding strategies in the future.

Acknowledgements

We would like to acknowledge the FruitLook FP7-PEOPLE-2013-IRSES-Marie Curie Action (Project ID: 612640), the SexSeed H2020-EU.1.3.3. MSCA-RISE-Marie Skłodowska-Curie Research, Innovation Staff Exchange project (Project ID: 690946), H2020-MSCA RISE Project (ExpoSeed GA-691109), and the Università degli Studi di Milano (UNIMI-RTD and Linea 2-DBS) for bringing together the team that developed this article. PG is grateful to Universitat Politècnica de València (PAID-10-18) and to the Japan Society for the Promotion of Science [FY2019 JSPS Postdoctoral Fellowship for Research in Japan (Standard)] for their respective postdoctoral grants. We would also like to thank the Society of Experimental Biology for the opportunity of summarizing some of our ideas in this article.

References

- Akita Y, Nakada M, Kanno A.** 2011. Effect of the expression level of an *AGAMOUS*-like gene on the petaloidy of stamens in the double-flowered lily, 'Elodie'. *Scientia Horticulturae* **128**, 48–53.
- Baldwin SJ, Schoen DJ.** 2017. Genetic variation for pseudo-self-compatibility in self-incompatible populations of *Leavenworthia alabamica* (Brassicaceae). *New Phytologist* **213**, 430–439.
- Baum DA, Hileman LC.** 2006. A developmental genetic model for the origin of the flower. In: Ainsworth C, ed. *Flowering and its manipulation*. Oxford: Blackwell Publishing Ltd, 1–27.
- Benlloch R, Berbel A, Ali L, Gohari G, Millán T, Madueño F.** 2015. Genetic control of inflorescence architecture in legumes. *Frontiers in Plant Science* **6**, 543.
- Benlloch R, Berbel A, Serrano-Mislata A, Madueño F.** 2007. Floral initiation and inflorescence architecture: a comparative view. *Annals of Botany* **100**, 659–676.
- Blackman BK, Rasmussen DA, Strasburg JL, Raduski AR, Burke JM, Knapp SJ, Michaels SD, Rieseberg LH.** 2011. Contributions of flowering time genes to sunflower domestication and improvement. *Genetics* **187**, 271–287.
- Blázquez MA, Ferrándiz C, Madueño F, Parcy F.** 2006. How floral meristems are built. *Plant Molecular Biology* **60**, 855–870.
- Blümel M, Dally N, Jung C.** 2015. Flowering time regulation in crops—what did we learn from Arabidopsis? *Current Opinion in Biotechnology* **32**, 121–129.
- Bowman JL, Smyth DR, Meyerowitz EM.** 1989. Genes directing flower development in *Arabidopsis*. *The Plant Cell* **1**, 37–52.
- Brothhaerts W.** 2003. New findings in apple S-genotype analysis resolve previous confusion and request the re-numbering of some S-alleles. *Theoretical and Applied Genetics* **106**, 703–714.
- Carmona MJ, Calonje M, Martínez-Zapater JM.** 2007. The FT/TFL1 gene family in grapevine. *Plant Molecular Biology* **63**, 637–650.
- Carr SM, Irish VF.** 1997. Floral homeotic gene expression defines developmental arrest stages in *Brassica oleracea* L. vars. *botrytis* and *italica*. *Planta* **201**, 179–188.
- Causier B, Schwarz-Sommer Z, Davies B.** 2010. Floral organ identity: 20 years of ABCs. *Seminars in Cell & Developmental Biology* **21**, 73–79.
- Chanderbali AS, Berger BA, Howarth DG, Soltis PS, Soltis DE.** 2016. Evolving ideas on the origin and evolution of flowers: new perspectives in the genomic era. *Genetics* **202**, 1255–1265.
- Chardon F, Damerval C.** 2005. Phylogenomic analysis of the PEBP gene family in cereals. *Journal of Molecular Evolution* **61**, 579–590.
- Cheng X, Li G, Tang Y, Wen J.** 2018. Dissection of genetic regulation of compound inflorescence development in *Medicago truncatula*. *Development* **145**, dev158766.
- Coen ES, Meyerowitz EM.** 1991. The war of the whorls: genetic interactions controlling flower development. *Nature* **353**, 31–37.
- Corbesier L, Vincent C, Jang S, et al.** 2007. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science* **316**, 1030–1033.
- Corley SB, Carpenter R, Copsey L, Coen E.** 2005. Floral asymmetry involves an interplay between TCP and MYB transcription factors in *Antirrhinum*. *Proceedings of the National Academy of Sciences, USA* **102**, 5068–5073.
- Cubas P, Vincent C, Coen E.** 1999. An epigenetic mutation responsible for natural variation in floral symmetry. *Nature* **401**, 157–161.
- Danilevskaya ON, Meng X, Ananiev EV.** 2010. Concerted modification of flowering time and inflorescence architecture by ectopic expression of *TFL1*-like genes in maize. *Plant Physiology* **153**, 238–251.
- Darwin C.** 1859. *On the origin of species*. London: Routledge.
- Darwin C.** 1868. *The variation of animals and plants under domestication*. London: John Murray.
- da Silva MR, Clément D, Gramacho KP, Monteiro WR, Argout X, Lanaud C, Lopes U.** 2016. Genome-wide association mapping of sexual incompatibility genes in cacao (*Theobroma cacao* L.). *Tree Genetics & Genomes* **12**, 62.
- de Bellaire LDL, Fouré E, Abadie C, Carlier J.** 2010. Black leaf streak disease is challenging the banana industry. *Fruits* **65**, 327–342.
- de Graaf BH, Vátovec S, Juárez-Díaz JA, Chai L, Kooblall K, Wilkins KA, Zou H, Forbes T, Franklin FC, Franklin-Tong VE.** 2012. The Papaver self-incompatibility pollen S-determinant, PrpS, functions in *Arabidopsis thaliana*. *Current Biology* **22**, 154–159.
- De Mattia F, Imazio S, Grassi F, Baneh HD, Scienza A, Labra M.** 2008. Study of genetic relationships between wild and domesticated grapevine distributed from Middle East Regions to European countries. *Rendiconti Lincei Scienze Fisiche e Naturali* **19**, 223–240.
- De Nettancourt D.** 2001. *Incompatibility and incongruity in wild and cultivated plants*. Berlin: Springer Science & Business Media.
- DeVries H.** 1904. *Species and varieties: their origin by mutation*. Chicago, IL: Open Court Publishing Company.
- D'Hont A, Denoeud F, Aury JM, et al.** 2012. The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature* **488**, 213–217.
- Diez CM, Trujillo I, Martínez-Urdiroz N, Barranco D, Rallo L, Marfil P, Gaut BS.** 2015. Olive domestication and diversification in the Mediterranean Basin. *New Phytologist* **206**, 436–447.

- Dita MA, Waalwijk C, Buddenhagen IW, Souza Jr MT, Kema GHJ.** 2010. A molecular diagnostic for tropical race 4 of the banana fusarium wilt pathogen. *Plant Pathology* **59**, 348–357.
- Ditta G, Pinyopich A, Robles P, Pelaz S, Yanofsky MF.** 2004. The SEP4 gene of *Arabidopsis thaliana* functions in floral organ and meristem identity. *Current Biology* **14**, 1935–1940.
- Doebley J, Stec A, Hubbard L.** 1997. The evolution of apical dominance in maize. *Nature* **386**, 485–488.
- Doi K, Izawa T, Fuse T, Yamanouchi U, Kubo T, Shimatani Z, Yano M, Yoshimura A.** 2004. *Ehd1*, a B-type response regulator in rice, confers short-day promotion of flowering and controls *FT*-like gene expression independently of *Hd1*. *Genes & Development* **18**, 926–936.
- Dong Y, Liu J, Li PW, Li CQ, Lü TF, Yang X, Wang YZ.** 2018. Evolution of Darwin's peloric glloxinia (*Sinningia speciosa*) is caused by a null mutation in a pleiotropic TCP gene. *Molecular Biology and Evolution* **35**, 1901–1915.
- Dubois A, Raymond O, Maene M, Baudino S, Langlade NB, Boltz V, Vergne P, Bendahmane M.** 2010. Tinkering with the C-function: a molecular frame for the selection of double flowers in cultivated roses. *PLoS One* **5**, e9288.
- Duclos DV, Björkman T.** 2008. Meristem identity gene expression during curd proliferation and flower initiation in *Brassica oleracea*. *Journal of Experimental Botany* **59**, 421–433.
- Endress PK.** 1999. Symmetry in flowers: diversity and evolution. *International Journal of Plant Sciences* **160**, S3–S23.
- Evers JB, van der Krol AR, Vos J, Struik PC.** 2011. Understanding shoot branching by modelling form and function. *Trends in Plant Science* **16**, 464–467.
- Farinelli D, Breton C, Koubouris G, Famiani F, Villemur P, Bervillé A.** 2018. Reply to Saumitou-Laprade *et al.* (2017). Controlling for genetic identity of varieties, pollen contamination and stigma receptivity is essential to characterize the self-incompatibility system of *Olea europaea* L. *Evolutionary Applications* **11**, 1465–1470.
- Favaro R, Pinyopich A, Battaglia R, Kooiker M, Borghi L, Ditta G, Yanofsky MF, Kater MM, Colombo L.** 2003. MADS-box protein complexes control carpel and ovule development in *Arabidopsis*. *The Plant Cell* **15**, 2603–2611.
- Fjellheim S, Boden S, Trevaskis B.** 2014. The role of seasonal flowering responses in adaptation of grasses to temperate climates. *Frontiers in Plant Science* **5**, 431.
- François L, Verdenaud M, Fu X, Ruleman D, Dubois A, Vandenbussche M, Bendahmane A, Raymond O, Just J, Bendahmane M.** 2018. A miR172 target-deficient AP2-like gene correlates with the double flower phenotype in roses. *Scientific Reports* **8**, 12912.
- Fujii S, Kubo K, Takayama S.** 2016. Non-self- and self-recognition models in plant self-incompatibility. *Nature Plants* **2**, 16130.
- Fujino K, Yamanouchi U, Yano M.** 2013. Roles of the Hd5 gene controlling heading date for adaptation to the northern limits of rice cultivation. *Theoretical and Applied Genetics* **126**, 611–618.
- Galego L, Almeida J.** 2002. Role of DIVARICATA in the control of dorsoventral asymmetry in Antirrhinum flowers. *Genes & Development* **16**, 880–891.
- Galimba KD, Tolkin TR, Sullivan AM, Melzer R, Theißen G, Di Stilio VS.** 2012. Loss of deeply conserved C-class floral homeotic gene function and C- and E-class protein interaction in a double-flowered ranunculid mutant. *Proceedings of the National Academy of Sciences, USA* **109**, E2267–E2275.
- Gao A, Zhang J, Zhang W.** 2017. Evolution of *RAD*- and *DIV*-like genes in plants. *International Journal of Molecular Sciences* **18**, 1961.
- Gattolin S, Cirilli M, Pacheco I, et al.** 2018. Deletion of the miR172 target site in a TOE-type gene is a strong candidate variant for dominant double-flower trait in Rosaceae. *The Plant Journal* **96**, 358–371.
- Gervasi DD, Schiestl FP.** 2017. Real-time divergent evolution in plants driven by pollinators. *Nature Communications* **8**, 14691.
- Gibbs PE.** 2014. Late-acting self-incompatibility—the pariah breeding system in flowering plants. *New Phytologist* **203**, 717–734.
- Glover BJ, Bunnewell S, Martin C.** 2004. Convergent evolution within the genus *Solanum*: the specialised anther cone develops through alternative pathways. *Gene* **331**, 1–7.
- Gonthier L, Blassiau C, Mörchen M, Cadalen T, Poirat M, Hendriks T, Quillet MC.** 2013. High-density genetic maps for loci involved in nuclear male sterility (NMS1) and sporophytic self-incompatibility (S-locus) in chicory (*Cichorium intybus* L., Asteraceae). *Theoretical and Applied Genetics* **126**, 2103–2121.
- Goretti D, Martignago D, Landini M, et al.** 2017. Transcriptional and post-transcriptional mechanisms limit heading date 1 (*Hd1*) function to adapt rice to high latitudes. *PLOS Genetics* **13**, e1006530.
- Goslin K, Zheng B, Serrano-Mislata A, et al.** 2017. Transcription factor interplay between *LEAFY* and *APETALA1/CAULIFLOWER* during floral initiation. *Plant Physiology* **174**, 1097–1109.
- Goto K, Meyerowitz EM.** 1994. Function and regulation of the *Arabidopsis* floral homeotic gene *PISTILLATA*. *Genes & Development* **8**, 1548–1560.
- Haecker A, Gross-Hardt R, Geiges B, Sarkar A, Breuninger H, Herrmann M, Laux T.** 2004. Expression dynamics of *WOX* genes mark cell fate decisions during early embryonic patterning in *Arabidopsis thaliana*. *Development* **131**, 657–668.
- Havlíčková L, Jozová E, Klíma M, Kucera V, Curn V.** 2014. Detection of self-incompatible oilseed rape plants (*Brassica napus* L.) based on molecular markers for identification of the class I S haplotype. *Genetics and Molecular Biology* **37**, 556–559.
- Hayama R, Yokoi S, Tamaki S, Yano M, Shimamoto K.** 2003. Adaptation of photoperiodic control pathways produces short-day flowering in rice. *Nature* **422**, 719–722.
- Heijmans K, Ament K, Rijpkema AS, Zethof J, Wolters-Arts M, Gerats T, Vandenbussche M.** 2012. Redefining C and D in the petunia ABC. *The Plant Cell* **24**, 2305–2317.
- Hiscock SJ, Tabah DA.** 2003. The different mechanisms of sporophytic self-incompatibility. *Philosophical Transactions of the Royal Society B: Biological Sciences* **358**, 1037–1045.
- Honma T, Goto K.** 2001. Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* **409**, 525–529.
- Huang C, Sun H, Xu D, et al.** 2018. *ZmCCT9* regulates maize flowering time. *Proceedings of the National Academy of Sciences, USA* **115**, E334–E341.
- Hubbard L, McSteen P, Doebley J, Hake S.** 2002. Expression patterns and mutant phenotype of teosinte branched1 correlate with growth suppression in maize and teosinte. *Genetics* **162**, 1927–1935.
- Hunt HV, Denyer K, Packman LC, Jones MK, Howe CJ.** 2010. Molecular basis of the waxy endosperm starch phenotype in broomcorn millet (*Panicum miliaceum* L.). *Molecular Biology and Evolution* **27**, 1478–1494.
- Huq E, Tepperman JM, Quail PH.** 2000. GIGANTEA is a nuclear protein involved in phytochrome signaling in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **97**, 9789–9794.
- Igic B, Kohn JR.** 2006. The distribution of plant mating systems: study bias against obligately outcrossing species. *Evolution* **60**, 1098–1103.
- Igic B, Lande R, Kohn JR.** 2008. Loss of self-incompatibility and its evolutionary consequences. *International Journal of Plant Sciences* **169**, 93–104.
- Ikeda-Kawakatsu K, Maekawa M, Izawa T, Itoh J, Nagato Y.** 2012. *ABERRANT PANICLE ORGANIZATION 2/RFL*, the rice ortholog of *Arabidopsis* *LEAFY*, suppresses the transition from inflorescence meristem to floral meristem through interaction with *APO1*. *The Plant Journal* **69**, 168–180.
- Irish V.** 2017. The ABC model of floral development. *Current Biology* **27**, R887–R890.
- Itoh H, Nonoue Y, Yano M, Izawa T.** 2010. A pair of floral regulators sets critical day length for *Hd3a* florigen expression in rice. *Nature Genetics* **42**, 635–638.
- Iwano M, Ito K, Fujii S, et al.** 2015. Calcium signalling mediates self-incompatibility response in the Brassicaceae. *Nature Plants* **1**, 15128.
- Iwano M, Takayama S.** 2012. Self/non-self discrimination in angiosperm self-incompatibility. *Current Opinion in Plant Biology* **15**, 78–83.
- Iwata T, Nagasaki O, Ishii HS, Ushimaru A.** 2012. Inflorescence architecture affects pollinator behaviour and mating success in *Spiranthes sinensis* (Orchidaceae). *New Phytologist* **193**, 196–203.
- Jack T, Brockman LL, Meyerowitz EM.** 1992. The homeotic gene *APETALA3* of *Arabidopsis thaliana* encodes a MADS box and is expressed in petals and stamens. *Cell* **68**, 683–697.

- Jofuku KD, den Boer BG, Van Montagu M, Okamoto JK. 1994. Control of *Arabidopsis* flower and seed development by the homeotic gene *APETALA2*. *The Plant Cell* **6**, 1211–1225.
- Kaothien-Nakayama P, Isogai A, Takayama S. 2010. Self-incompatibility systems in flowering plants. In: Pua E, Davey M, eds. *Plant developmental biology—biotechnological perspectives*. Berlin: Springer Berlin Heidelberg, 459–485.
- Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harrison MJ, Weigel D. 1999. Activation tagging of the floral inducer *FT*. *Science* **286**, 1962–1965.
- Kaufmann K, Muiño JM, Jauregui R, Airoidi CA, Smaczniak C, Krajewski P, Angenent GC. 2009. Target genes of the MADS transcription factor *SEPALLATA3*: integration of developmental and hormonal pathways in the *Arabidopsis* flower. *PLoS Biology* **7**, e1000090.
- Kebrom TH, Brutnell TP. 2015. Tillering in the *sugary1* sweet corn is maintained by overriding the teosinte branched1 repressive signal. *Plant Signaling & Behavior* **10**, e1078954.
- Kempin SA, Savidge B, Yanofsky MF. 1995. Molecular basis of the cauliflower phenotype in *Arabidopsis*. *Science* **267**, 522–525.
- Kitashiba H, Nasrallah JB. 2014. Self-incompatibility in Brassicaceae crops: lessons for interspecific incompatibility. *Breeding Science* **64**, 23–37.
- Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T. 1999. A pair of related genes with antagonistic roles in mediating flowering signals. *Science* **286**, 1960–1962.
- Kojima S, Takahashi Y, Kobayashi Y, Monna L, Sasaki T, Araki T, Yano M. 2002. *Hd3a*, a rice ortholog of the *Arabidopsis FT* gene, promotes transition to flowering downstream of *Hd1* under short-day conditions. *Plant & Cell Physiology* **43**, 1096–1105.
- Komiya R, Ikegami A, Tamaki S, Yokoi S, Shimamoto K. 2008. *Hd3a* and *RFT1* are essential for flowering in rice. *Development* **135**, 767–774.
- Koo BH, Yoo SC, Park JW, Kwon CT, Lee BD, An G, Zhang Z, Li J, Li Z, Paek NC. 2013. Natural variation in *OsPRR37* regulates heading date and contributes to rice cultivation at a wide range of latitudes. *Molecular Plant* **6**, 1877–1888.
- Koseva B, Crawford DJ, Brown KE, Mort ME, Kelly JK. 2017. The genetic breakdown of sporophytic self-incompatibility in *Tolpis coronopifolia* (Asteraceae). *New Phytologist* **216**, 1256–1267.
- Krizek BA, Fletcher JC. 2005. Molecular mechanisms of flower development: an armchair guide. *Nature Reviews. Genetics* **6**, 688–698.
- Kuhn DN, Livingstone DS III, Richards JH, Manosalva P, Van den Berg N, Chambers AH. 2019. Application of genomic tools to avocado (*Persea americana*) breeding: SNP discovery for genotyping and germplasm characterization. *Scientia Horticulturae* **246**, 1–11.
- Kyozuka J, Konishi S, Nemoto K, Izawa T, Shimamoto K. 1998. Down-regulation of *RFL*, the *FLO/LFY* homolog of rice, accompanied with panicle branch initiation. *Proceedings of the National Academy of Sciences, USA* **95**, 1979–82.
- Labate JA, Robertson LD, Baldo AM, Björkman T. 2006. Inflorescence identity gene alleles are poor predictors of inflorescence type in broccoli and cauliflower. *Journal of the American Society for Horticultural Science* **131**, 667–673.
- Lanaud C, Fouet O, Legavre T, et al. 2017. Deciphering the *Theobroma cacao* self-incompatibility system: from genomics to diagnostic markers for self-compatibility. *Journal of Experimental Botany* **68**, 4775–4790.
- Langewisch T, Zhang H, Vincent R, Joshi T, Xu D, Bilyeu K. 2014. Major soybean maturity gene haplotypes revealed by SNPviz analysis of 72 sequenced soybean genomes. *PLoS One* **9**, e94150.
- Lao X, Suwabe K, Niikura S, Kakita M, Iwano M, Takayama S. 2014. Physiological and genetic analysis of CO₂-induced breakdown of self-incompatibility in *Brassica rapa*. *Journal of Experimental Botany* **65**, 939–951.
- Lescot T. 2010. The genetic diversity of banana in figures. *FruiTrop* **177**, 48–52.
- Li YH, Zhao SC, Ma JX, et al. 2013. Molecular footprints of domestication and improvement in soybean revealed by whole genome re-sequencing. *BMC Genomics* **14**, 579.
- Liljgren SJ, Gustafson-Brown C, Pinyopich A, Ditta GS, Yanofsky MF. 1999. Interactions among *APETALA1*, *LEAFY*, and *TERMINAL FLOWER1* specify meristem fate. *The Plant Cell* **11**, 1007–1018.
- Lippman ZB, Cohen O, Alvarez JP, Abu-Abied M, Pekker I, Paran I, Eshed Y, Zamir D. 2008. The making of a compound inflorescence in tomato and related nightshades. *PLoS Biology* **6**, e288.
- Litt A. 2007. An evaluation of A-function: evidence from the *APETALA1* and *APETALA2* gene lineages. *International Journal of Plant Sciences* **168**, 73–91.
- Liu B, Watanabe S, Uchiyama T, et al. 2010. The soybean stem growth habit gene *Dt1* is an ortholog of *Arabidopsis TERMINAL FLOWER1*. *Plant Physiology* **153**, 198–210.
- Liu C, Teo ZW, Bi Y, Song S, Xi W, Yang X, Yin Z, Yu H. 2013. A conserved genetic pathway determines inflorescence architecture in *Arabidopsis* and rice. *Developmental Cell* **24**, 612–622.
- Liu Z, Zhang D, Liu D, Li F, Lu H. 2013. Exon skipping of *AGAMOUS* homolog *PrseAG* in developing double flowers of *Prunus lannesiana* (Rosaceae). *Plant Cell Reports* **32**, 227–237.
- Loor Solorzano RG, Fouet O, Lemainque A, Pavék S, Boccara M, Argout X, Amores F, Courtois B, Risterucci AM, Lanaud C. 2012. Insight into the wild origin, migration and domestication history of the fine flavour Nacional *Theobroma cacao* L. variety from Ecuador. *PLoS One* **7**, e48438.
- Lowman AC, Purugganan MD. 1999. Duplication of the *Brassica oleracea APETALA1* floral homeotic gene and the evolution of domesticated cauliflower. *Journal of Heredity* **90**, 514–520.
- Luo D, Carpenter R, Copley L, Vincent C, Clark J, Coen E. 1999. Control of organ asymmetry in flowers of *Antirrhinum*. *Cell* **99**, 367–376.
- Luo D, Carpenter R, Vincent C, Copley L, Coen E. 1996. Origin of floral asymmetry in *Antirrhinum*. *Nature* **383**, 794–799.
- Mahrez W, Arellano MS, Moreno-Romero J, Nakamura M, Shu H, Nanni P, Köhler C, Grisse W, Hennig L. 2016. H3K36ac is an evolutionary conserved plant histone modification that marks active genes. *Plant Physiology* **170**, 1566–1577.
- Mandel MA, Gustafson-Brown C, Savidge B, Yanofsky MF. 1992. Molecular characterization of the *Arabidopsis* floral homeotic gene *APETALA1*. *Nature* **360**, 273–277.
- Martínez-Gómez P, Sánchez-Pérez R, Dicenta F. 2006. Fruit development in almond. *Acta Horticulturae* **726**, 241–246.
- McClure B. 2006. New views of S-RNase-based self-incompatibility. *Current Opinion in Plant Biology* **9**, 639–646.
- McClure B. 2012. Plant self-incompatibility: ancient system becomes a new tool. *Current Biology* **22**, R86–R87.
- McGarry RC, Prewitt SF, Culpepper S, Eshed Y, Lifschitz E, Ayre BG. 2016. Monopodial and sympodial branching architecture in cotton is differentially regulated by the *Gossypium hirsutum* *SINGLE FLOWER TRUSS* and *SELF-PRUNING* orthologs. *New Phytologist* **212**, 244–258.
- McKey D, Elias M, Pujol B, Duputié A. 2010. The evolutionary ecology of clonally propagated domesticated plants. *New Phytologist* **186**, 318–332.
- Michaels SD, Amasino RM. 1999. *FLOWERING LOCUS C* encodes a novel MADS domain protein that acts as a repressor of flowering. *The Plant Cell* **11**, 949–956.
- Mimida N, Goto K, Kobayashi Y, Araki T, Ahn JH, Weigel D, Murata M, Motoyoshi F, Sakamoto W. 2001. Functional divergence of the *TFL1*-like gene family in *Arabidopsis* revealed by characterization of a novel homologue. *Genes to Cells* **6**, 327–336.
- Miller AJ, Gross BL. 2011. From forest to field: perennial fruit crop domestication. *American Journal of Botany* **98**, 1389–1414.
- Molinero-Rosales N, JAMILENA M, Zurita S, Gómez P, Capel J, Lozano R. 1999. *FALSIFLORA*, the tomato orthologue of *FLORICAULA* and *LEAFY*, controls flowering time and floral meristem identity. *The Plant Journal* **20**, 685–693.
- Morel P, Heijmans K, Rozier F, Zethof J, Chamot S, Bento SR, Vialette-Guiraud A, Chambrier P, Trehin C, Vandenbussche M. 2017. Divergence of the floral A-function between an asterid and a rosid species. *The Plant Cell* **29**, 1605–1621.
- Moyroud E, Glover BJ. 2017. The evolution of diverse floral morphologies. *Current Biology* **27**, R941–R951.
- Müller NA, Wijnen CL, Srinivasan A, et al. 2016. Domestication selected for deceleration of the circadian clock in cultivated tomato. *Nature Genetics* **48**, 89–93.

- Müller NA, Zhang L, Koornneef M, Jiménez-Gómez JM. 2018. Mutations in *EID1* and *LNK2* caused light-conditional clock deceleration during tomato domestication. *Proceedings of the National Academy of Sciences, USA* **115**, 7135–7140.
- Ockendon DJ. 1974. Distribution of self-incompatibility alleles and breeding structure of open-pollinated cultivars of Brussels sprouts. *Heredity* **33**, 159–171.
- Parcy F, Nilsson O, Busch MA, Lee I, Weigel D. 1998. A genetic framework for floral patterning. *Nature* **395**, 561–566.
- Park DH, Somers DE, Kim YS, Choy YH, Lim HK, Soh MS, Kim HJ, Kay SA, Nam HG. 1999. Control of circadian rhythms and photoperiodic flowering by the Arabidopsis *GIGANTEA* gene. *Science* **285**, 1579–1582.
- Pelaz S, Ditta GS, Baumann E, Wisman E, Yanofsky MF. 2000. B and C floral organ identity functions require *SEPALLATA* MADS-box genes. *Nature* **405**, 200–203.
- Pelaz S, Tapia-López R, Alvarez-Buylla ER, Yanofsky MF. 2001. Conversion of leaves into petals in Arabidopsis. *Current Biology* **11**, 182–184.
- Perez-Rodriguez M, Jaffe FW, Butelli E, Glover BJ, Martin C. 2005. Development of three different cell types is associated with the activity of a specific MYB transcription factor in the ventral petal of *Antirrhinum majus* flowers. *Development* **132**, 359–370.
- Pinyopich A, Ditta GS, Savidge B, Liljegren SJ, Baumann E, Wisman E, Yanofsky MF. 2003. Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature* **424**, 85–88.
- Pekkala N, Knott KE, Kotiaho JS, Nissinen K, Puurtinen M. 2014. The effect of inbreeding rate on fitness, inbreeding depression and heterosis over a range of inbreeding coefficients. *Evolutionary Applications* **7**, 1107–1119.
- Pnueli L, Carmel-Goren L, Hareven D, Gutfinger T, Alvarez J, Ganai M, Zamir D, Lifschitz E. 1998. The SELF-PRUNING gene of tomato regulates vegetative to reproductive switching of sympodial meristems and is the ortholog of CEN and TFL1. *Development* **125**, 1979–1989.
- Prusinkiewicz P, Erasmus Y, Lane B, Harder LD, Coen E. 2007. Evolution and development of inflorescence architectures. *Science* **316**, 1452–1456.
- Purugganan MD, Boyles AL, Suddith JI. 2000. Variation and selection at the *CAULIFLOWER* floral homeotic gene accompanying the evolution of domesticated *Brassica oleracea*. *Genetics* **155**, 855–862.
- Putterill J, Robson F, Lee K, Simon R, Coupland G. 1995. The *CONSTANS* gene of *Arabidopsis* promotes flowering and encodes a protein showing similarities to zinc finger transcription factors. *Cell* **80**, 847–857.
- Ratcliffe OJ, Bradley DJ, Coen ES. 1999. Separation of shoot and floral identity in Arabidopsis. *Development* **126**, 1109–1120.
- Reardon W, Fitzpatrick DA, Fares MA, Nugent JM. 2009. Evolution of flower shape in *Plantago lanceolata*. *Plant Molecular Biology* **71**, 241–250.
- Santos ES, Cerqueira-Silva CB, Mori GM, Ahnert D, Mello DL, Pires JL, Corrêa RX, de Souza AP. 2015. Genetic structure and molecular diversity of cacao plants established as local varieties for more than two centuries: the genetic history of Cacao plantations in Bahia, Brazil. *PLoS One* **10**, e0145276.
- Sassa H. 2016. Molecular mechanism of the S-RNase-based gametophytic self-incompatibility in fruit trees of Rosaceae. *Breeding Science* **66**, 116–121.
- Saumitou-Laprade P, Vernet P, Vekemans X, *et al.* 2017. Elucidation of the genetic architecture of self-incompatibility in olive: evolutionary consequences and perspectives for orchard management. *Evolutionary Applications* **10**, 867–880.
- Schilling S, Pan S, Kennedy A, Melzer R. 2018. MADS-box genes and crop domestication: the jack of all traits. *Journal of Experimental Botany* **69**, 1447–1469.
- Sehgal N, Singh S. 2018. Progress on deciphering the molecular aspects of cell-to-cell communication in Brassica self-incompatibility response. *Biotech* **8**, 347.
- Serrano-Mislata A, Goslin K, Zheng B, Rae L, Wellmer F, Graciet E, Madueño F. 2017. Regulatory interplay between *LEAFY*, *APETALA1/CAULIFLOWER* and *TERMINAL FLOWER1*: new insights into an old relationship. *Plant Signaling & Behavior* **12**, e1370164.
- Sessions A, Yanofsky MF, Weigel D. 2000. Cell-cell signaling and movement by the floral transcription factors *LEAFY* and *APETALA1*. *Science* **289**, 779–782.
- Shannon S, Meeks-Wagner DR. 1993. Genetic interactions that regulate inflorescence development in Arabidopsis. *The Plant Cell* **5**, 639–655.
- Smith LB, King GJ. 2000. The distribution of *BoCAL*-a alleles in *Brassica oleracea* is consistent with a genetic model for curd development and domestication of the cauliflower. *Molecular Breeding* **6**, 603–613.
- Smyth DR. 2018. Evolution and genetic control of the floral ground plan. *New Phytologist* **220**, 70–86.
- Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T. 2015. Photoperiodic flowering: time measurement mechanisms in leaves. *Annual Review of Plant Biology* **66**, 441–464.
- Spiegel-Roy P. 1986. Domestication of fruit trees. *Developments in Agricultural and Managed Forest Ecology* **16**, 201–211.
- Stewart D, Graciet E, Wellmer F. 2016. Molecular and regulatory mechanisms controlling floral organ development. *FEBS Journal* **283**, 1823–1830.
- Studer A, Zhao Q, Ross-Ibarra J, Doebley J. 2011. Identification of a functional transposon insertion in the maize domestication gene *tb1*. *Nature Genetics* **43**, 1160–1163.
- Su S, Xiao W, Guo W, *et al.* 2017. The *CYCLOIDEA-RADIALIS* module regulates petal shape and pigmentation, leading to bilateral corolla symmetry in *Torenia fournieri* (Linderniaceae). *New Phytologist* **215**, 1582–1593.
- Sun Y, Fan Z, Li X, Li J, Yin H. 2014. The *APETALA1* and *FRUITFUL* homologs in *Camellia japonica* and their roles in double flower domestication. *Molecular Breeding* **33**, 821–834.
- Teo ZW, Song S, Wang YQ, Liu J, Yu H. 2014. New insights into the regulation of inflorescence architecture. *Trends in Plant Science* **19**, 158–165.
- Theissen G, Melzer R. 2007. Molecular mechanisms underlying origin and diversification of the angiosperm flower. *Annals of Botany* **100**, 603–619.
- Theißen G, Melzer R, Rümpler F. 2016. MADS-domain transcription factors and the floral quartet model of flower development: linking plant development and evolution. *Development* **143**, 3259–3271.
- Theißen G, Saedler H. 2001. Floral quartets. *Nature* **409**, 469–471.
- Tian Z, Wang X, Lee R, Li Y, Specht JE, Nelson RL, McClean PL, Qiu L, Ma J. 2010. Artificial selection for determinate growth habit in soybean. *Proceedings of the National Academy of Sciences, USA* **107**, 8563–8568.
- Ueda M, Zhang Z, Laux T. 2011. Transcriptional activation of Arabidopsis axis patterning genes *WOX8/9* links zygote polarity to embryo development. *Developmental Cell* **20**, 264–270.
- Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G. 2004. Photoreceptor regulation of *CONSTANS* protein in photoperiodic flowering. *Science* **303**, 1003–1006.
- Vieira J, Fonseca NA, Vieira CP. 2009. RNase-based gametophytic self-incompatibility evolution: questioning the hypothesis of multiple independent recruitments of the S-pollen gene. *Journal of Molecular Evolution* **69**, 32–41.
- Wagner D, Sablowski RW, Meyerowitz EM. 1999. Transcriptional activation of *APETALA1* by *LEAFY*. *Science* **285**, 582–584.
- Wang B, Smith SM, Li J. 2018. Genetic regulation of shoot architecture. *Annual Review of Plant Biology* **69**, 437–468.
- Weberling F. 1989. Structure and evolutionary tendencies of inflorescences in the Leguminosae. *Monographs in Systematic Botany from the Missouri Botanical Garden* **29**, 35–58.
- Wei X, Xu J, Guo H, Jiang L, Chen S, Yu C, Zhou Z, Hu P, Zhai H, Wan J. 2010. *DTH8* suppresses flowering in rice, influencing plant height and yield potential simultaneously. *Plant Physiology* **153**, 1747–1758.
- Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM. 1992. *LEAFY* controls floral meristem identity in *Arabidopsis*. *Cell* **69**, 843–859.
- Weigel D, Meyerowitz EM. 1994. The ABCs of floral homeotic genes. *Cell* **78**, 203–209.
- Weigel D, Nilsson O. 1995. A developmental switch sufficient for flower initiation in diverse plants. *Nature* **377**, 495–500.
- Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D. 2005. Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science* **309**, 1056–1059.

- Wils CR, Kaufmann K.** 2017. Gene-regulatory networks controlling inflorescence and flower development in *Arabidopsis thaliana*. *Biochimica et Biophysica Acta* **1860**, 95–105.
- Wu F, Shi X, Lin X, Liu Y, Chong K, Theißen G, Meng Z.** 2017. The ABCs of flower development: mutational analysis of *AP1/FUL*-like genes in rice provides evidence for a homeotic (A)-function in grasses. *The Plant Journal* **89**, 310–324.
- Wu X, Chory J, Weigel D.** 2007. Combinations of *WOX* activities regulate tissue proliferation during *Arabidopsis* embryonic development. *Developmental Biology* **309**, 306–316.
- Wu X, Dabi T, Weigel D.** 2005. Requirement of homeobox gene *STIMPY/WOX9* for *Arabidopsis* meristem growth and maintenance. *Current Biology* **15**, 436–440.
- Wünsch A, Hormaza JI.** 2004. Molecular evaluation of genetic diversity and S-allele composition of local Spanish sweet cherry (*Prunus avium* L.) cultivars. *Genetic Resources and Crop Evolution* **51**, 635–641.
- Wyatt R.** 1982. Inflorescence architecture: how flower number, arrangement, and phenology affect pollination and fruit-set. *American Journal of Botany* **69**, 585–594.
- Xiao Z, Han F, Hu Y, et al.** 2019. Overcoming cabbage crossing incompatibility by the development and application of self-compatibility-QTL-specific markers and genome-wide background analysis. *Frontiers in Plant Science* **10**, 189.
- Xue W, Xing Y, Weng X, et al.** 2008. Natural variation in *Ghd7* is an important regulator of heading date and yield potential in rice. *Nature Genetics* **40**, 761–767.
- Yano M, Katayose Y, Ashikari M, et al.** 2000. *Hd1*, a major photoperiod sensitivity quantitative trait locus in rice, is closely related to the *Arabidopsis* flowering time gene *CONSTANS*. *The Plant Cell* **12**, 2473–2484.
- Yanofsky MF, Ma H, Bowman JL, Drews GN, Feldmann KA, Meyerowitz EM.** 1990. The protein encoded by the *Arabidopsis* homeotic gene *agamous* resembles transcription factors. *Nature* **346**, 35–39.
- Ye L, Wang B, Zhang W, Shan H, Kong H.** 2016. Gains and losses of cis-regulatory elements led to divergence of the *Arabidopsis* *APETALA1* and *CAULIFLOWER* duplicate genes in the time, space, and level of expression and regulation of one paralog by the other. *Plant Physiology* **171**, 1055–1069.
- Yeager AF.** 1927. Determinate growth in the tomato. *Journal of Heredity* **18**, 263–265.
- Zhai H, Lü S, Wang Y, et al.** 2014. Allelic variations at four major maturity *E* genes and transcriptional abundance of the *E1* gene are associated with flowering time and maturity of soybean cultivars. *PLoS One* **9**, e97636.
- Zheng H, Kawabata S, Zheng H, Kawabata S.** 2017. Identification and validation of new alleles of *FALSIFLORA* and *COMPOUND INFLORESCENCE* genes controlling the number of branches in tomato inflorescence. *International Journal of Molecular Sciences* **18**, 1572.
- Zhu S, Wang J, Cai M, et al.** 2017. The *OsHAPL1-DTH8-Hd1* complex functions as the transcription regulator to repress heading date in rice. *Journal of Experimental Botany* **68**, 553–568.
- Zhu Z, Tan L, Fu Y, Liu F, Cai H, Xie D, Wu F, Wu J, Matsumoto T, Sun C.** 2013. Genetic control of inflorescence architecture during rice domestication. *Nature Communications* **4**, 2200.