



UNIVERSITAT  
POLITÈCNICA  
DE VALÈNCIA

**Embryo cryopreservation and transfer to rederive  
a paternal rabbit line after 18 generations.  
Evaluation of growth and reproductive traits.**

***Ph.D. Thesis by***

Jorge Daniel Juárez Moreno

***Supervisor***

José Salvador Vicente Antón

**València, May 2022**



**Embryo cryopreservation and transfer to rederive a paternal rabbit line after 18 generations. Evaluation of growth and reproductive traits.**

## **INDEX OF CONTENTS**

## INDEX

<b>1. GENERAL INTRODUCTION .....</b>	<b>2</b>
1.1. GENERALITIES .....	2
1.2. GENETIC IMPROVEMENT OF MEAT RABBIT .....	3
1.2.1. Effects of growth rate selection on weight and meat quality .....	4
1.2.2. Effects of growth rate selection on reproductive performance .....	6
1.2.2.1. <i>Semen quality traits</i> .....	6
1.2.2.2. <i>Effects on prenatal development and reproductive performance</i> .....	8
1.2.2.3. <i>Effect on health and robustness</i> .....	9
1.3. CONTROL POPULATIONS TO EVALUATE SELECTION EFFECTS .....	10
1.4. REFERENCES .....	11
<b>2. OBJECTIVES .....</b>	<b>22</b>
<b>3. CHAPTER I. EFFECTS OF REDERIVATION BY EMBRYO VITRIFICATION ON PERFORMANCE IN A RABBIT PATERNAL LINE .....</b>	<b>25</b>
3.1. INTRODUCTION .....	26
3.2. MATERIAL AND METHODS .....	27
3.2.1. Animals and experimental design .....	28
3.2.2. Growing traits analysis .....	29
3.2.3. Female reproductive management .....	30
3.2.4. Evaluation of litter size components .....	30
3.2.5. Foetal growth. Ultrasound measurement .....	31
3.2.6. Statistical analysis .....	31
3.3. RESULTS .....	33
3.3.1. Growing traits. Rederived effect .....	34
3.3.2. Evolution of growth traits by filial generation .....	34
3.3.2. Reproductive Traits .....	35
3.3.2.1. <i>Reproductive performance of F2 (R38 and R38V) females.</i> .....	35
3.3.2.2. <i>Litter size components of F2 (R38 and R38V) females</i> .....	36
3.3.2.3. <i>Foetal growth during gestation in F2 (R38 and R38V) females</i> .....	36
3.4. DISCUSSION .....	40
3.4. CONCLUSIONS .....	43
3.5. REFERENCES .....	45
<b>4. CHAPTER II. EVALUATION BY RE-DERIVATION OF A PATERNAL LINE AFTER 18 GENERATIONS ON SEMINAL TRAITS, PROTEOME AND FERTILITY .....</b>	<b>53</b>

4.1. INTRODUCTION .....	54
4.2. MATERIAL AND METHODS .....	56
4.2.1. Animals .....	56
4.2.2. Experimental design .....	57
4.2.3. Semen collection .....	58
4.2.4. Evaluation of ejaculates .....	58
4.2.4.1. Ejaculate quantity and sperm production .....	58
4.2.4.2. Ejaculate quality .....	58
4.2.5. Plasma and sperm protein extraction samples .....	60
4.2.6. Proteomic relative quantification analysis: SWATH (DIA) MSMS analysis .....	60
4.2.6.1. Spectral libraries building .....	60
4.2.6.2. ProteinPilot v5.0. search engine (Sciex) .....	61
4.2.6.3. Swath analysis of samples .....	62
4.2.7. Fertility parameters. Artificial insemination .....	63
4.2.7.1. Semen collection and evaluation .....	63
4.2.7.2. Semen extension .....	63
4.2.7.3. Insemination procedure .....	64
4.2.8. Statistical analyses .....	64
4.3. RESULTS .....	65
4.3.1. Ejaculates and sperm traits .....	65
4.3.2. Plasma and sperm proteome .....	66
4.3.3. Fertility parameters .....	68
4.4. DISCUSSION AND CONCLUSION .....	75
4.5. REFERENCES .....	79

<b>5. CHAPTER III. REDERIVATION BY CRYOPRESERVATION OF A PATERNAL LINE OF RABBITS SUGGESTS EXHAUSTION OF SELECTION FOR POST-WEANING DAILY WEIGHT GAIN AFTER 37 GENERATIONS .....</b>	<b>91</b>
5.1. INTRODUCTION .....	92
5.2. MATERIAL AND METHODS .....	93
5.2.1. Animals .....	93
5.2.2. Embryo vitrification and transfer to population rederivation .....	94
5.2.3. Reproduction management in rederived and filial generations .....	95
5.2.4. Growing traits analysis .....	96
5.2.5. Statistical analysis .....	96
5.3. RESULTS .....	98
5.3.1. Descriptive growing traits .....	98
5.3.2. Growing traits by generation interval of rederived populations. Selection effect .....	100
5.3.3. Evolution of growth traits by filial generation .....	101

5.3.4. Estimated growth curves: F1 rederived populations (R19V and R37V) .....	102
5.3.5. Estimated growth curves: F3 rederived population (R21 and R39).....	102
5.4. DISCUSSION .....	106
5.5. CONCLUSIONS .....	110
5.6. REFERENCES .....	111
<b>6. CHAPTER IV. EVALUATION OF FOETAL GROWTH, LITTER SIZE AND REPRODUCTIVE PERFORMANCE IN RABBIT AFTER 18 GENERATIONS OF SELECTION FOR GROWTH RATE USING CRYOPRESERVED EMBRYOS .....</b>	<b>119</b>
6.1. INTRODUCTION .....	120
6.2. MATERIAL AND METHODS .....	122
6.2.1. Animals .....	122
6.2.2. Reproduction management .....	123
6.2.3. Laparoscopy and evaluated litter size components .....	124
6.2.4. Foetal growth. ultrasound measurement .....	124
6.2.5. Statistical analyses .....	125
6.3. RESULTS .....	126
6.3.1. Reproductive performance between rederived populations .....	126
6.3.2. Litter size components between rederived groups .....	127
6.3.3. Foetal growth during gestation .....	127
6.4. DISCUSSION AND CONCLUSIONS .....	130
6.5. REFERENCES .....	134
<b>7. GENERAL DISCUSSION .....</b>	<b>144</b>
REFERENCES .....	151
<b>8. CONCLUSIONS .....</b>	<b>160</b>

## INDEX OF FIGURES

<b>Figure 3.1.</b> Graphical abstract .....	29
<b>Figure 3.2.</b> Effect of rederivation by embryo cryopreservation on: A, weaning weight (WW, kg); B, weight at end of fattening (EFW, Kg) and C, average daily gain (ADG, g/day) from filial generations (F1-R37 and R37V, F2-R38 and R38V and F3-R39 and R39V). Number of rabbits for each population and filial generation is noted into columns .....	35
<b>Figure 3.3.</b> Effect of rederivation by embryo cryopreservation on fetal growth at 12, 19 and 26 days of gestation from a paternal rabbit line. Number of fetal traits measured is noted into columns. A, Fetal sac area (cm <sup>2</sup> ); B, crown-rump length of fetuses (mm); C, maternal placenta area (mm <sup>2</sup> ) and fetal placenta (mm <sup>2</sup> ) .....	37
<b>Figure 4.1.</b> Flowchart of the experiment performed to obtain the evaluated generations of a rabbit line selected by growth rate .....	57
<b>Figure 4.2.</b> a. Partial Least Squares Discriminant Analysis (PLS-DA) showing the classification of seminal plasma samples belonging to G21V and G39V. b. Partial Least Squares Discriminant Analysis (PLS-DA) showing the classification of sperm samples belonging to G21V and G39V .....	69
<b>Figure 4.3.</b> a. Heat map representing levels of differentially expressed seminal plasma proteins between male groups (G21V and G39V). b. Heat map representing levels of differentially expressed sperm proteins between male groups (G21V and G39V) .....	71
<b>Figure 4.4.</b> a. Distribution of molecular function, biological process and cell components of differentially expressed seminal plasma proteins between male groups (G21V and G39V). b. Distribution of molecular function, biological process and cell components of differentially expressed sperm proteins between male groups (G21V and G39V) .....	73
<b>Figure 5.1.</b> Experimental design: Two experimental progenies were developed from vitrified embryos stored in 2000 (19th generation of selection) and 2015 (37th generation of selection). All rabbits were identified and weighted at weaning and end of the fattening period to calculate the average daily gain. A sample of males and females were weighted weekly from birth to 20 weeks age in F1 y F3 generation to estimate Gompertz curve parameters .....	94

<b>Figure 5.2.</b> Growth curves from birth to 20-week-old between F1 rederived populations from R19V and R37V generations. V: Rederived from vitrified embryos. Growth curves were fitted using the Gompertz equation for (A) R19V and R37V generations and (B) sex .....	103
<b>Figure 5.3.</b> Growth curves from birth to 20-week-old between F3 filial generations from R21 and R39generations. V: Rederived from vitrified embryos. Growth curves were fitted using the Gompertz equation for (A) R21 and R39 generations and (B) sex .....	105
<b>Figure 5.4.</b> Gompertz growth curve from birth to 20-week-old for interaction between populations and sex of animals .....	106
<b>Figure 6.1.</b> Graphical abstract of the experimental study .....	123
<b>Figure 6.2.</b> Ultrasonography measurements of the foetal sac (FS), crown-rump length (CRL) of foetus and the placental measurements of the two-lobed foetal (L1FP and L2FP) and maternal (MP) at 12, 19 and 26 day of gestation .....	125



## INDEX OF TABLES

<b>Table 3.1.</b> Total number of parents used to generate the offspring analysed .....	28
<b>Table 3.2.</b> Effect of rederivation by embryo cryopreservation on weaning weight (WW, kg), weight at end of fattening (EFW, Kg) and average daily gain (ADG, g/day) from a paternal rabbit line .....	34
<b>Table 3.3.</b> Effect of rederivation by embryo cryopreservation on total litter size, liveborn, litter size at weaning, perinatal and lactation mortality rates from a paternal rabbit line .....	38
<b>Table 3.4.</b> Effect of rederivation by embryo cryopreservation on litter size components from a paternal rabbit line .....	39
<b>Table 3.5.</b> Number of young rabbits weighted from weaning to end of fattening period and average and standard deviation to birth alive (BA), weaning weight (WW), weight at end fattening (EFW) and average daily weight gain (ADG) .....	44
<b>Table 4.1.</b> Seminal traits in G21V and G39V .....	66
<b>Table 4.2.</b> Seminal parameters of G21V and G39V groups used in proteome analysis and fertility assay .....	67
<b>Table 4.3.</b> Reproductive performance of inseminated does (least square mean $\pm$ standard error least) .....	68
<b>Table 4.4a.</b> Highlighted differentially seminal plasma proteins between male groups (G21V and G39V) with a fold change (FC) $\geq$ 1.5 after $\log_2$ transformation.	74
<b>Table 4.4b.</b> Highlighted differentially sperm proteins between male groups (G21V and G39V) with a fold change (FC) $\geq$ 1.5 after $\log_2$ transformation .....	75
<b>Table 5.1.</b> The total number of parents used to generate the offspring analysed	98
<b>Table 5.2.</b> Number of young rabbits weighted from weaning to end fattening period and average and standard deviation to birth alive (BA), weaning weight (WW), weight at end fattening (EFW) and average daily weight gain (ADG) .....	99
<b>Table 5.3.</b> Effect of selection for growth rate on weaning weight (WW, kg), weight at end of fattening period (EFW, kg) and average daily weight gain (ADG, g/day) from two rederived population after 18 generations .....	100

<b>Table 5.4.</b> Differences of least-square means of weaning weight (WW, Kg), weight at end of fattening period (EFW, kg) and average daily weight gain (ADG, g/day) from filial generations. V: Rederived from vitrified embryos .....	101
<b>Table 5.5.</b> Gompertz curve parameters from R19V and R37V generations .....	102
<b>Table 5.6.</b> Gompertz curve parameters from R21 and R39 generations .....	104
<b>Table 6.1.</b> Effect of selection for growth rate on total litter size, liveborn, litter size at weaning, perinatal and lactation mortality rates and delivery rate from two rederived population separated by 18 generations .....	128
<b>Table 6.2.</b> Effect of selection and lactation status on litter size components from two rederived population separated by 18 generations .....	129
<b>Table 6.3.</b> Effect of selection and lactation status on foetal sac and placentae area and foetal size at 12, 19 and 26 days of gestation from two rederived population separated by 18 generations .....	131

