

Contents

Glossary	19
1. Introduction	25
1.1. Minimal hepatic encephalopathy	27
1.1.1. Factors contributing to minimal hepatic encephalopathy	28
1.1.1.1. Hyperammonemia	30
1.1.1.2. Alterations during peripheral inflammation	31
1.1.1.3. The interplay between neuroinflammation and systemic inflammation	33
1.1.1.4. Inflammation in cases of liver cirrhosis	35
1.1.1.5. Microbiota	37
1.1.2. Treatments that reduce peripheral inflammation	39
1.2. The human immune system and inflammatory responses	40
1.2.1. Immune system cells	41
1.2.1.1. CD4+ T cells	42
1.2.2. Communication between cells during immune responses	43
1.2.3. Inflammatory response	45
1.2.4. T-cell receptor	46
1.3. Omics and systems biology	49
1.3.1. Overview of multi-omics data	50
1.3.1.1. High-throughput RNA quantification	52
1.3.1.2. Mass spectrometry platforms	53
1.3.2. Analysis steps and integration of multi-omics data	55
1.3.2.1. Single-omics analysis	55
1.3.2.2. Multi-omics analysis	62
1.3.2.2.1. Multivariate approaches to omics data analysis	65
1.3.3. Computational immunology	67
2. Hypothesis, aims, and contributions	71
2.1. Hypothesis and objectives	73
2.2. Aims	73
2.3. Contributions	74
2.3.1. Journal papers	78
2.3.2. Conferences	79
3. Multi-omic analysis of changes in the peripheral immune system associated with the appearance of minimal hepatic encephalopathy in patients with cirrhosis	81
3.1. Introduction	83
3.2. Methods	85

3.2.1. Overview of the analysis strategy	85
3.2.2. Patients and sample collection	87
3.2.3. Transcriptomics profiling of plasma samples	88
3.2.4. Metabolomics profiling of serum samples	89
3.2.5. Analysis of cytokines in serum samples	91
3.2.6. Data pre-processing plots	92
3.2.7. Single-omic analysis	92
3.2.8. Omic power analysis	94
3.2.9. Obtaining modules of coordinated metabolites and cytokines	94
3.2.10. Integration of multi-omic datasets	95
3.2.11. Data and code availability	98
3.3. Results and discussion	98
3.3.1. Data pre-processing	98
3.3.2. Transcriptomics analysis confirms changes in the immunophenotype in patients with minimal hepatic encephalopathy and unveils new altered pathways	102
3.3.3. A coordinated metabolic and cytokinetic signature is present in patients with minimal hepatic encephalopathy	110
3.3.4. Multi-omic integration analysis highlights altered genes and associated pathways related with metabolites and cytokines in patients with minimal hepatic encephalopathy	115
3.4. Conclusion	126
4. Identification of altered signaling pathways in CD4+ lymphocytes isolated from patients with minimal hepatic encephalopathy	127
4.1. Introduction	129
4.2. Methods	131
4.2.1. Patients	131
4.2.2. Assessment of cognitive function using psychometric tests	132
4.2.3. Isolation of CD4+ T lymphocytes	132
4.2.4. RNA extraction and sequencing	132
4.2.5. RNA-seq analysis	133
4.2.6. miRNA-seq analysis	134
4.2.7. Omic power analysis	134
4.2.8. Biological integration analysis	135
4.2.9. Data availability	135
4.3. Results and discussion	136
4.3.1. Data pre-processing	136
4.3.2. CD4+ lymphocytes from patients with minimal hepatic encephalopathy showed alterations in 167 mRNA and 20 signaling	

pathways compared to patients without minimal hepatic encephalopathy	140
4.3.3. CD4+ lymphocytes showed altered levels for 13 miRNAs and 7 transcription factors comparing patients with versus without minimal hepatic encephalopathy	149
4.3.4. Identification of miRNAs and transcription factors that modulate the differential expression of key genes involved in regulating immune responses in minimal hepatic encephalopathy	150
4.4. Conclusion	159
5. A Nextflow pipeline for T-cell receptor repertoire reconstruction and analysis from RNA sequencing data	161
5.1. Introduction	163
5.2. Methods	165
5.2.1. Overview of the analysis strategy	165
5.2.2. Sample collection, RNA sequencing, and read pre-processing	167
5.2.3. Repertoire reconstruction	167
5.2.4. Hill-based evenness profiles	167
5.2.5. Shannon Evenness	168
5.2.6. Jaccard similarity	169
5.2.7. k-mer-based CDR3 β analysis	169
5.2.8. T-cell receptor sequence similarity architecture	169
5.2.9. Generation of the graphics	170
5.2.10. Antigen/Disease-specific T-cell receptor databases	170
5.2.11. Fisher's exact test analysis	170
5.2.12. Nextflow pipeline	171
5.2.13. Data and code availability	171
5.3. Results and discussion	172
5.3.1. Step-by-step analysis overview	172
5.3.2. T-cell receptor sequences can be recovered from RNA-seq data (steps 1–5)	175
5.3.3. T-cell receptor sequence profiling in patients with minimal hepatic encephalopathy	179
5.3.3.1. Low clonal convergence among patient samples (step 6a)	179
5.3.3.2. High clonal expansion in all the samples, independently of the immune status (step 6b)	180
5.3.3.3. Increased within-repertoire similarity based on repertoire architecture was unconstrained by immune status (steps 6c–d)	181
5.3.4. T-cell receptors with known disease associations were overrepresented in minimal hepatic encephalopathy CDR3 repertoires (step 6e)	183

5.4. Conclusion	187
6. General discussion and conclusions	191
6.1. General discussion	193
6.2. Conclusions	195
6.3. Future perspectives	197
References	201