
INDEX	1
ABBREVIATIONS.....	5
ABSTRACT	13
RESUMEN	19
RESUM	25
I. INTRODUCTION	31
1. CYCLOOXYGENASE AND PROSTAGLANDINS	33
1.1 Cyclooxygenase description, function, and localisation.....	33
1.1 COX isoforms, expression, and regulation.....	34
1.2 Prostanoids and prostaglandins.....	36
2. PROSTAGLANDINS AND COX-2 IN HEPATIC PHYSIOPATHOLOGY	38
2.1 The liver: function and structure.....	38
2.2 Hepatic COX-2 expression.....	40
2.3 COX-2 in liver pathology.....	41
3. LIVER ISCHEMIA-REPERFUSION INJURY	43
3.1 Liver transplant.....	43
3.2 Ischemia-reperfusion injury.....	44
3.3 Inflammatory immune response.....	46
3.3.1 <i>Liver macrophages activation and recruitment</i>	47
3.3.2 <i>Neutrophil activation and recruitment</i>	48
3.4 Oxidative stress and antioxidant response	48
3.5 Endoplasmic reticulum stress	49
3.6 Cellular death.....	51
3.6.1 <i>Apoptosis</i>	51
3.6.2 <i>Necrosis</i>	52
3.7 Autophagy	53
4. MITOCHONDRIA.....	54
4.1 Mitochondrial respiration	56
4.1.1 <i>Electron transport chain and oxidative phosphorylation system</i>	56
4.1.1 <i>Supercomplexes</i>	57
4.2 Mitochondrial dynamics	60
4.3 Role of the mitochondria in ischemia-reperfusion injury	62
II. OBJECTIVES	65
III. MATERIALS AND METHODS.....	69
1. REAGENTS AND EQUIPMENT.....	71
2. ANIMAL MODELS AND EXPERIMENTAL DESIGN.....	72
2.1 Generation and maintenance of the <i>h-COX-2 Tg</i> mice.....	72

2.2	Pure genetic background <i>h-COX-2 Tg</i> mice line generation (B6).....	74
2.3	F1 hybrid genetic background animals (B6D2 F1)	74
2.4	Mice genotyping	74
2.5	Experimental design.....	76
2.5.1	<i>Liver ischemia-reperfusion model</i>	76
2.5.2	<i>Ischemia preconditioning</i>	77
2.5.3	<i>Drug-inhibition of COX-2</i>	78
3.	SUBCELLULAR FRACTIONATION	78
3.1	Mitochondrial isolation	78
3.2	Nuclear fractionation.....	79
4.	HISTOLOGICAL ANALYSIS	80
4.1	Fixation, paraffin tissue inclusion, and sectioning	80
4.2	Deparaffination, rehydration, and dehydration of samples	80
4.3	Fixed paraffined tissue stains.....	80
4.3.1	<i>H&E staining</i>	80
4.3.2	<i>Ly6G⁺ staining</i>	81
4.4	Tissue cryopreservation and sectioning.....	82
4.5	Dihydroethidium staining.....	82
4.6	Transmission Electronic Microscopy (TEM)	82
5.	PROTEIN DETECTION	83
5.1	Protein extraction and quantification	83
5.2	Protein electrophoresis and immunoblotting	84
5.3	Protein native electrophoresis for supercomplex detection (BN-PAGE).....	85
5.4	Immunofluorescence	88
6.	GENE DETECTION AND EXPRESSION	88
6.1	Isolation of total DNA from frozen liver tissue	88
6.2	Determination of mitochondrial DNA content	89
6.3	RNA extraction, RT, and qPCR.....	90
6.4	Determination of <i>Xbp1</i> splicing	93
6.5	RNAseq and NGS analysis.....	94
7.	DETERMINATION OF METABOLITES AND CYTOKINES	94
7.1	Plasma collection	94
7.2	PGE ₂	94
7.3	Plasma transaminases.....	95
7.4	Plasma cytokines.....	95
7.5	Total NAD and NADH content determination	96
7.6	AMP and ATP determination	96
8.	ENZYMATIC ACTIVITIES	97
8.1	Caspase 3.....	97
8.2	Catalase	97
8.3	Superoxide dismutase.....	97
8.4	Glutathione reductase and Glutathione Peroxidase	98
8.5	Total glutathione and oxidised glutathione measurement.....	98
8.6	Myeloperoxidase activity	99
8.7	Indirect measurement of ROS production	99
9.	FUNCTIONAL ASSAYS.....	100
9.1	Measurement of mitochondrial membrane potential	100

9.2	High resolution respirometry	100
10.	PATIENTS	103
11.	STATISTICS	103
IV.	RESULTS.....	105
1.	CHARACTERIZATION OF THE ROLE OF COX-2 IN HEPATIC ISCHEMIA-REPERFUSION MODEL	109
1.1	Establishment of the ischemia-reperfusion injury pathology.....	109
1.2	Liver of <i>h-COX-2 Tg</i> mice have less tissue damage compared to the <i>Wt</i> animals	109
1.3	A mixed genetic background is required to reproduce the ischemia-reperfusion injury model in mice	112
1.4	Lower hepatic damage is due specifically to COX-2 expression	114
1.5	Hepatic preconditioning attenuates liver damage in <i>Wt</i> mice	115
1.6	The gene expression profile is altered after ischemia-reperfusion injury	117
2.	THE INFLAMMATORY RESPONSE IS ATTENUATED WHEN COX-2 IS EXPRESSED	119
2.1	Ischemia-reperfusion increases inflammatory markers that are lower in <i>h-COX-2 Tg</i> mice.....	119
2.2	Neutrophil infiltration is decreased with COX-2 expression	121
2.3	COX-2 expression attenuated NF- κ B pathway	123
3.	CELLULAR STRESS IS REDUCED WHEN COX-2 IS EXPRESSED IN HEPATOCYTES	124
3.1	Constitutive hepatocyte COX-2 expression leads to decreased apoptosis	124
3.2	Hepatocyte COX-2 expression activates autophagy after ischemia-reperfusion..	126
3.3	Unfolded protein response is activated in ischemia-reperfusion	127
3.4	The antioxidant response is enhanced when COX-2 is overexpressed.....	130
4.	STUDY OF THE MITOCHONDRIAL FUNCTION.....	134
4.1	Mitochondrial respiration is higher in <i>h-COX-2 Tg</i> livers after ischemia-reperfusion.....	134
4.2	Higher respiration in <i>h-COX-2 Tg</i> mitochondria is not due to an increase in ETC complexes expression	138
4.3	Ischemia-reperfusion causes loss in mitochondrial membrane potential and is prevented by COX-2 expression	140
4.4	Mitochondria number and phenotype are not altered because of the COX-2 expression.....	141
4.5	Mitochondrial dynamics is altered after ischemia-reperfusion	144
4.6	OPA1 processing is enhanced by a higher activity of OMA1 in <i>Wt</i> mitochondria after ischemia-reperfusion	146
5.	INCREASED PLASMATIC PGE ₂ ARE ASSOCIATED WITH AN IMPROVEMENT IN LIVER FUNCTION	148

INDEX

V.	DISCUSSION.....	151
VI.	CONCLUSIONS	171
VII.	BIBLIOGRAPHY	175
VIII.	ANNEXES	203
	ANNEX 1 – PATIENT INFORMATION TABLE	205
	ANNEX 2 – INDEX OF FIGURES	211
	ANNEX 3 – INDEX OF TABLES.....	215
	ANNEX 4 – PUBLICATIONS	217