

ANEXOS

Anexo 1. Cebadores directos y reversos empleados para analizar la expresión diferencial de genes mediante qRT-PCR (F: cebador directo, R: cebador reverso)

qRT <i>WRKY48</i> F	CAAAAACAGTCCTTATCCCAGAAGC
qRT <i>WRKY48</i> R	CGTGAGCATTCCGATGTGTC
qRT <i>MYB15</i> F	CCAAATACTTGGCAATAGATGGTCAG
qRT <i>MYB15</i> R	TGCTGGTCTTAGGTTTAGCTGG
qRT <i>MYB96</i> F	CCCTTTTAGGCAACAGATGGGC
qRT <i>MYB96</i> R	TGAACTAGTGTGTTGATGAAGAGACACC
qRT <i>MYB42</i> F	TCTTCTCGGCAACAGATGGTC
qRT <i>MYB42</i> R	TGTCGGCTTTCGATGAAGTTTCAG
qRT <i>MYB85</i> F	CATGCTAATCTCGGCAATAAGTGGTC
qRT <i>MYB85</i> R	CACTGAGACATCGTCTGGATTGG
qRT <i>MYB64</i> F	TGGCCGTTATTGGAGAAATCGTC
qRT <i>MYB64</i> R	CATCACCAACTTTATCAACTTCCTGTCT
qRT <i>CESA8</i> F	GCGGCAATCCTTACGATGAGAAT
qRT <i>CESA8</i> R	CAGCTCTCCACTCTGTTCTTCC
qRT <i>WRKY33</i> F	GAACAATGGAGCCAAACCGAG
qRT <i>WRKY33</i> R	GTTTCATTGTCCCCTTCCATCTC
qRT <i>CYP94B1</i> F	CCGTTCCAGAACTTGTTAAGGCT
qRT <i>CYP94B1</i> R	GGCCCGGATGATCTCAGAC

