

THESIS: Treating boar sperm with cholesterol-loaded cyclodextrins or cyclodextrins prior to cryopreservation: effects on post-thaw in vitro sperm quality of sperm cryopreserved in different freezing extenders

AUTHOR: Eva Blanch Torres

TABLE OF CONTENTS

ABSTRACT	1
RESUMEN	3
RESUM	5
CHAPTER 1 GENERAL INTRODUCTION	9
1. Use of frozen-thawed sperm in pig production	9
1.1. Stress conditions during the cryopreservation process: cold shock and osmotic stress	10
1.1.1. Cold shock	10
1.1.2. Osmotic stress	11
1.2. New strategies to improve frozen-thawed boar sperm quality	12
2. Cholesterol as strategy for improving the quality of frozen-thawed sperm	13
2.1. Effect of cholesterol	13
2.2. Pre-freezing treatment of sperm with cholesterol-loaded cyclodextrins (CLC)	15
2.3. Treatment of boar sperm with CLC	16
3. Cryoprotectant requirements for CLC-treated boar sperm	16

3.1. Cryoprotectants	16
3.2. Glycerol	18
3.3. Alternative cryoprotectants for the cryopreservation of CLC-treated boar sperm	19
CHAPTER 2 OBJECTIVES	31
CHAPTER 3 STUDY 1: Response of boar sperm to the treatment with cholesterol-loaded cyclodextrins added prior to cryopreservation	35
1. Introduction	37
2. Material and methods	40
2.1. Reagents and media	40
2.2. Preparation of cholesterol-loaded cyclodextrins (CLC)	40
2.3. Semen collection and dilution	41
2.4. Cryopreservation and thawing	41
2.5. Sperm evaluation	42
2.6. Statistical analyses	43
2.7. Experimental design	44
2.7.1. Experiment 1: Effect of CLC concentration on boar sperm cryosurvival	44
2.7.2. Experiment 2: Effect of CLC treatment on sperm cryosurvival from males classified as good or poor freezers	44
3. Results	45
3.1. Experiment 1: Effect of CLC concentration on boar sperm cryosurvival	45
3.2. Experiment 2: Effect of CLC treatment on sperm cryosurvival from males classified as good or poor freezers	47
4. Discussion	49
CHAPTER 4 STUDY 2: Egg yolk and glycerol requirements for freezing boar spermatozoa treated with methyl-β-cyclodextrin or cholesterol-loaded	57

cyclodextrin

1. Introduction	60
2. Materials and methods	61
2.1. Reagents and media	61
2.2 Preparation of cholesterol-loaded cyclodextrins (CLC)	62
2.3 Semen collection and dilution	62
2.4. Sperm cryopreservation	63
2.5 Sperm quality evaluation	63
2.6. Experimental design	65
2.6.1. Experiment 1: Requirement of egg yolk for freezing CLC- or methyl- β -cyclodextrin (MBCD)-treated boar spermatozoa	65
2.6.2. Experiment 2: Requirement of glycerol for freezing CLC- or MBCD-treated boar spermatozoa	65
2.6.3. Experiment 3: Effect of increasing glycerol concentration on cryosurvival of CLC-treated boar sperm	66
2.7. Statistical analyses	66
3. Results	67
3.1. Experiment 1: Requirement of egg yolk for freezing CLC- or MBCD-treated boar spermatozoa	67
3.2. Experiment 2: Requirement of glycerol for freezing CLC- or MBCD-treated boar spermatozoa	68
3.3. Experiment 3: Effect of increasing glycerol concentration on cryosurvival of CLC-treated boar sperm	70
4. Discussion	71

CHAPTER 5 STUDY 3: The contradictory effect of amides on	85
cryopreservation of boar semen treated with cholesterol-loaded cyclodextrin, methyl- β -cyclodextrin or non treated	
1. Introduction	88
2. Materials and methods	90
2.1. Reagents and extenders	90
2.2 Preparation of cholesterol-loaded cyclodextrins (CLC)	91
2.3 Semen collection and dilution	91
2.4. Sperm cryopreservation	91
2.5 Sperm quality evaluation	92
2.6. In vitro fertilizing ability evaluation	93
2.7. Experimental design	95
2.7.1. Experiment 1: The effect of different amides on fresh semen	95
2.7.2. Experiment 2: Determining the best concentration of dimethylacetamide (DMA) to cryopreserve boar semen	95
2.7.3. Experiment 3: Determining the effectiveness of different amides as cryoprotectants for boar semen	96
2.7.4. Experiment 4: Determining the in vitro fertilizing ability of boar sperm cryopreserved with dimethylformamide (DMF) or glycerol	96
2.8. Statistical analyses	97
3. Results	98
3.1. Experiment 1: The effect of different amides on fresh semen	98
3.2. Experiment 2: Determining the best concentration of DMA to cryopreserve boar semen	100
3.3. Experiment 3: Determining the effectiveness of different amides as cryoprotectants for boar semen	101

3.4. Experiment 4: Determining the in vitro fertilizing ability of boar sperm cryopreserved with DMF or glycerol	104
4. Discussion	106
CHAPTER 6 GENERAL DISCUSSION	119
1. Response of boar sperm to the treatment with cholesterol-loaded cyclodextrins (CLC) or cyclodextrins alone (MBCD) using a conventional freezing extender	121
2. Response of boar sperm to the treatment with cholesterol-loaded cyclodextrins (CLC) or cyclodextrins alone (MBCD) using alternative glycerol and egg yolk concentrations in the freezing extender	123
2.1. Egg yolk	123
2.2. Glycerol	124
3. Response of boar sperm to the treatment with cholesterol-loaded cyclodextrins (CLC) or cyclodextrins alone (MBCD) using alternative cryoprotectants (amides) in the freezing extender	126
CHAPTER 7 CONCLUSIONS	135

INDEX OF TABLES

- Chapter 3, Table 1:** Percentages of viable, total motile and progressively motile sperm (least square means \pm standard error) 10 min after thawing and after 1 h of incubation at 39°C when boar spermatozoa were treated with different concentrations (mg/120 x 10⁶ sperm) of methyl- β -cyclodextrin (MBCD), or methyl- β -cyclodextrin pre-loaded with cholesterol (CLC) and cryopreserved in Lactose-egg yolk diluent. n = 10. 46
- Chapter 3, Table 2:** Percentages of viable, total motile, progressively motile sperm (least square means \pm standard error) 10 min and 1 h after thawing, when spermatozoa from boars whose sperm normally cryopreserve well (good freezers) or poorly (poor freezers) were cryopreserved using normal methods (control) or were treated with 1 mg/120 x 10⁶ sperm of methyl- β -cyclodextrin pre-loaded with cholesterol (CLC) prior to cryopreservation. n = 4 categorized as good freezers and 7 categorized as poor freezers. 48
- Chapter 4, Table 1:** Post-thaw percentages of total motile sperm (TMS), rapid progressively motile sperm (RPM) and sperm with intact plasma membrane (sperm viability; SV) in boar semen samples treated with 1 mg/120 x 10⁶ sperm of methyl- β -cyclodextrin (MBCD) or methyl- β -cyclodextrin pre-loaded with cholesterol (CLC) or no treated (control) and subsequently frozen in a extender with 20 or 10% of egg yolk. 68
- Chapter 4, Table 2:** Percentages of total motile sperm (TMS), rapid progressively motile sperm (RPM) and plasma membrane intact sperm (sperm viability; SV) after thawing when boar spermatozoa were treated with 1 69

mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD), or methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) or no treated (control) and then cryopreserved with different percentages of glycerol. Data of interaction between semen treatments and glycerol concentrations.

Chapter 4, Table 3: Percentages of total motile sperm (TMS), rapid 71 progressively motile sperm (RPM), plasma membrane intact sperm (sperm viability; SV) and viable sperm with low mitochondrial membrane potential (MMP) after thawing when boar spermatozoa were treated with 1 mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD) or no treated (control) and then cryopreserved with 3% of glycerol or treated with methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) and then cyropreserved with different percentages of glycerol.

Chapter 5, Table 1: Molecular weights (MW) of the cryoprotectants and 99 percentages of sperm with intact plasma membrane (sperm viability; SV) in boar semen samples treated with 1 mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD) or methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) and subsequently diluted in a extender with 3% of glycerol or 5% of different amides. Evaluation was performed before (0h) and after incubation at 22 °C for 4h (n = 6 ejaculates).

Chapter 5, Table 2: Post-thaw percentages of total motile sperm (TMS), rapid 101 progressively motile sperm (RPM) and sperm with intact plasma membrane (sperm viability; SV) when boar spermatozoa (n = 10 ejaculates) were treated with 1 mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD), or methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) and then cyropreserved with 3% of glycerol or different percentages (5 or 7%) of dimethylacetamide (DMA).

Chapter 5, Table 3: Post-thaw percentages of total motile sperm (TMS), rapid 103

progressively motile sperm (RPM), sperm with intact plasma membrane (sperm viability; SV) and viable sperm with low mitochondrial membrane potential (VS-LMMP) when boar spermatozoa (n = 10 ejaculates) were treated with 1 mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD), or methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) and then cryopreserved with 3% of glycerol, 5% of methylformamide (MF), 5% of dimethylacetamide (DMA) or 5% of dimethylformamide (DMF).

Chapter 5, Table 4: Sperm penetration ability of sow's immature oocytes in 105

vitro (penetration rate (%)) and number of sperm per oocytes) when boar spermatozoa (n = 3 pools of semen) were treated with 1 mg/120 x 10⁶ sperm of methyl-β-cyclodextrin (MBCD), or methyl-β-cyclodextrin pre-loaded with cholesterol (CLC) and then cryopreserved with 3% of glycerol or 5% of dimethylformamide (DMF).