

Document downloaded from:

<http://hdl.handle.net/10251/65191>

This paper must be cited as:

Viudes De Castro, MP.; Saenz De Juano Ribes, MDLD.; Marco Jiménez, F.; Herreros Pomares, A.; Vicente Antón, JS. (2015). Effect of luteinizing hormone on rabbit ovarian superstimulation and embryo developmental potential. *Theriogenology*. 84(3):446-451. doi:10.1016/j.theriogenology.2015.04.001.



The final publication is available at

<https://dx.doi.org/10.1016/j.theriogenology.2015.04.001>

Copyright Elsevier

Additional Information

1 **Effect of luteinizing hormone on rabbit ovarian superstimulation and embryo**
2 **developmental potential**

3

4 M.P. Viudes-de-Castro¹, A. Pomares², M.D. Saenz de Juano i Ribes², F. Marco-
5 Jiménez², J.S. Vicente²

6 ¹Centro de Investigación y Tecnología Animal-Instituto Valenciano de
7 Investigaciones Agrarias (CITA-IVIA), Polígono La Esperanza nº 100, 12400,
8 Segorbe, Castellón, Spain

9 ²Instituto de Ciencia y Tecnología Animal, Universidad Politécnica de Valencia,
10 Departamento de Ciencia Animal: Laboratorio de Biotecnología de la Reproducción,
11 46071, Valencia, Spain

12

13

14

15

16 **Abstract**

17 Assisted reproduction technologies require ovarian stimulation to increase the
18 number of oocytes and embryos. Currently, superstimulation is achieved by
19 gonadotropin treatment, but the embryo yield and quality are highly variable.
20 Commonly, commercial preparations derived from pituitary and urinary origin are
21 used to superovulate. Hence, ovarian superstimulation protocols have usually
22 included both FSH and LH. The appearance of recombinant gonadotropins
23 manufactured by genetic engineering techniques has ensured high quality and batch-
24 to-batch consistency. Moreover, this enables us to assess the importance of LH in
25 the ovarian stimulation. The main aim of this study was to evaluate the effect of
26 recombinant human LH supplementation (10%) on embryonic development produced
27 by rabbit does superovulated with low or high concentration (18.75 or 37.50 IU) of
28 recombinant human FSH (rhFSH). Females treated with rhFSH increased the
29 ovulation rate, and it was significantly higher when the high FSH dose was
30 supplemented with LH. The superstimulation treatment used did not significantly
31 affect in vitro development rate until the expanded blastocyst stage. The results of
32 this study seem to suggest that, in terms of superovulatory response, when rabbit
33 does are treated with 37.5-IU rhFSH, the use of LH supplementation allows an
34 increase in the number of follicles recruited and the quality of embryos, in terms of
35 ability to develop in vitro until blastocyst, and the expression profile of OCT4,
36 NANOG, and SOX2 genes is not affected.

37

38 **Keywords:** rabbit, aminopeptidase activity; dilution rate, male genotype, reproductive
39 performance.

40

41 INTRODUCTION

42

43 Superstimulation of females from domestic animals is used to augment conventional
44 genetic improvement programs and facilitates advances in reproductive biotechnologies. Rabbit presents great interest as an experimental animal model for
45 biomedical research, genetic modeling, and fur and meat production [1,2]. Superstimulation is considered a fundamental method to establish an abundant source of
46 oocytes and embryos and requires the administration of exogenous gonadotropin.
47 Although it is well established that FSH alone can induce follicular growth, small
48 amounts of LH are necessary to support the follicular development [3,4]. The action
49 of LH on follicular development is not limited to providing androgen substrate for
50 aromatization but also exerts a direct effect on the stimulation and modulation of
51 folliculogenesis [5,6], and the influence and the amount of LH necessary for optimal
52 follicular stimulation is still being investigated. Until recently, superovulation
53 treatments in farm mammals were based on the use of eCG and FSH preparations
54 derived from pituitary extracts (porcine, ovine, or equine), in which the variable LH
55 concentration present may interfere the FSH activity [7,8], and it is one of the most
56 important factors for the variability in the superstimulation response, affecting the
57 ability of the ovary to respond to exogenous gonadotropin treatments. The use of
58 pituitary-derived products presents other problems including contamination from
59 other hormones, inconsistencies within and among batches, and the possibility of the
60 spread of disease-transmitting agents [8]. Several studies have attributed detrimental
61 effects of an excessive LH content on follicular development, but these studies have
62 not established an optimal dose of LH or an optimal ratio FSH:LH to maximize the
63 superstimulation response [9–11]. In rabbits, the effect of LH on superstimulation

66 has been studied using porcine FSH, obtaining highly variable results [11,12].
67 Because of break-throughs in recombinant technology, it is possible to easily
68 dispose of LH and FSH in an isolated manner, which facilitates the study of the role
69 of LH in ovarian response.

70

71 On the other hand, the usual parameters to evaluate embryo quality are the
72 morphologic appearance at recovery or the blastocyst rate after IVC. However, it
73 has been reported that messenger RNA (mRNA) expression is related with the
74 embryo quality [13–18]. In addition, the early steps in embryo development can be
75 affected by the culture media and conditions as well as the production procedure
76 itself, which alters gene expression [13,14,16,19–21]. Therefore, the evaluation of
77 gene expression may help assess the developmental quality of the embryos. It is
78 known that the transcription factor octamer-binding 4 (OCT4), NANOG homeobox
79 (NANOG), and sex-determining region Y-box 2 (SOX2) have essential roles in early
80 development and are considered key regulators of the pluripotency maintenance
81 system [22]. Thus, changes in their expression might trigger failures in embryo devel-
82 opment and implantation. The implications of super-stimulation experience acquired
83 in animal studies and its possible short-term and long-term consequences should be
84 taken into account when proposing gonadotropin treatment to increase the number
85 of oocytes or embryos produced for reproductive biotechnologies.

86

87 This study aimed to evaluate the relative importance of LH in ovarian stimulation with
88 recombinant human FSH (rhFSH) and the effect on the development ability of
89 embryos in vitro and the expression of transcription factors OCT4, SOX2, and
90 NANOG.

91 **2. MATERIAL AND METHODS**

92

93 Unless stated otherwise, all chemicals in this study were purchased from
94 Sigma-Aldrich Química S.A (Madrid, Spain). The proteomic analysis was carried out
95 in the CSIE_university of Valencia Proteomics Unit, a member of ISCIII ProteoRed
96 Proteomics Platform.

97

98

99 2.1. Animals

100 All animals were handled according to the principles of animal care published by
101 Spanish Royal Decree 53/2013 in the Official Spanish State Gazette (Boletín Oficial
102 del Estado) in 2013. The research was carried out at the experimental farm of the
103 Universidad Politécnica de Valencia. Forty-three nulliparous does and five bucks of
104 sexually mature New Zealand White rabbits were used. Animals were housed in flat-
105 deck cages, fed a standard pellet diet ad libitum, and had free access to water. An
106 alternating cycle of 16-hour light and 8-hour dark was used [23].

107

108 2.2. Hormonal treatment

109 Superstimulation was induced using rhFSH (GONAL-f 75; Serono Europe Ltd.,
110 London, UK) alone or in combination with recombinant human LH (rhLH; Luveris 75;
111 Serono Europe Ltd.). Does were treated intramuscularly with either 18.75 or 37.50 IU
112 rhFSH alone or in combination with 10% rhLH (1.87 and 3.75 IU, respectively)
113 dissolved in saline distributed in five equal doses at 12-hour intervals. Females were
114 assigned randomly to five treatment groups:

115

116 1. Control: females were not superstimulated but were treated with saline solution.
117 2. Low FSH: superstimulation was induced using 18.75-IU rhFSH.
118 3. Low FSH–LH: superstimulation was induced using 18.75-IU rhFSH p 10% rhLH
119 (1.87 IU).
120 4. High FSH: superstimulation was induced using 37.50-IU rhFSH.
121 5. High FSH–LH: superstimulation was induced using 37.50-IU rhFSH p 10% rhLH
122 (3.75 IU).
123
124 Does were inseminated with 0.5 mL of fresh heterospermic pool 10 to 12 hours after
125 the last gonadotropin injection. To prepare the heterospermic pool, ejaculates from
126 five males were collected using an artificial vagina. The 10-mL aliquot samples from
127 ejaculates were diluted 1:10 with a Tris-citrate-glucose extender (250-mM
128 Tris(hydroxymethyl)aminomethane, 83-mM citric acid, 50-mM glucose, pH 6.8–7.0,
129 300 mOsm/kg) for a prior motility rate evaluation. Ejaculates with an estimated
130 motility higher than 70% were pooled. From heterospermic pool, two aliquot samples
131 of 10 mL were taken; the first one was diluted 1:10 with the Tris-citrate-glucose
132 extender for motility rate evaluation in a computer-assisted analysis system, and the
133 second was diluted 1:10 with 1% of glutaraldehyde solution in PBS to calculate the
134 concentra- tion in a Thoma chamber and evaluate both the percent- ages of normal
135 intact acrosome and abnormal sperm by phase contrast at a magnification of 400.
136 Only hetero- spermic pools with more than 70% of motility rate, 85% of normal intact
137 acrosome, and less than 15% of abnormal sperm were used to inseminate the does.
138 Pooled semen was diluted to 40 million/mL by adding Tris-citrate-glucose extender to
139 prepare the seminal doses. Does were induced to ovulate with an intramuscularly
140 administration of 1-mg buserelin acetate at the same time as the insemination.

141 2.3. Embryo recovery

142 Females were slaughtered 38 to 40 hours after artificial insemination with an
143 intravenous injection of 0.6-g pentobarbital sodium (Dolethal; Vetoquinol, Madrid,
144 Spain), and the reproductive tract was immediately removed. Embryos were
145 recovered by perfusion of each oviduct with 5-mL Dulbecco's PBS without
146 calcium2.3.1. Seminal plasma preparation chloride and supplemented with 0.1% of
147 BSA. Embryos were scored by morphologic criteria according to Interna- tional
148 Embryo Transfer Society classification. Only embryos in the eight- to 16-cell stage
149 without morphologic abnor- malities in mucin coat, ZP, and embryo cells were
150 cataloged as normal embryos (Fig. 1).

151 The ovulation rate, number of embryos, and number of oocytes were
152 recorded. The recovery rate was calculated as the number of embryos and oocytes
153 recovered divided by the ovulation rate, multiplied by 100. The fertilization rate was
154 calculated as the number of embryos divided by the number of embryos and oocytes
155 recovered, multiplied by 100.

156

157 2.4. In vitro culture

158 A total of 429 embryos were cultured for 48 hours in medium TCM-199
159 (Sigma, St. Louis, MO, USA) supple- mented with 10% fetal bovine serum and
160 antibiotics (penicillin G sodium, 300,000 IU; penicillin G procaine, 700,000 IU; and
161 dihydrostreptomycin sulfate, 1250 mg; Penivet 1; Divasa Farmavic, Barcelona,
162 Spain) at 38.5 °C, 5% CO₂, and saturated humidity. The in vitro development ability
163 of embryos until blastocyst stage was recorded for subsequent analysis (Fig. 1).

164 2.5. RNA extraction and quantitative PCR analysis

165 PolyA RNA was extracted from pools consisting of 13 to 15 embryos using the
166 Dynabeads kit (Life Technologies, Carlsbad, CA, USA), following the manufacturer's
167 in- structions. Six independent embryo pools were produced for each experimental
168 group. Then, reverse transcription was carried out using SuperScript III Reverse
169 Transcriptase (Life Technologies AS, Oslo, Norway) according to the manufacturer's
170 instructions. Real-time polymerase chain reaction (PCR) reactions were conducted in
171 an Applied Biosystems 7500 system (Applied Biosystems). Every PCR was
172 performed from 5- μ L diluted 1:10 complementary DNA (cDNA) template, 250-nM of
173 forward and reverse specific primers (Table 1), and 10 μ L of Power SYBR Green
174 PCR Master Mix (Fermentas Gmbh, Madrid, Spain) in a final volume of 20 μ L. The
175 PCR protocol included an initial step of 50 $^{\circ}$ C (2 minutes), followed by 95 $^{\circ}$ C (10
176 minutes), and 42 cycles of 95 $^{\circ}$ C (15 seconds) and 60 $^{\circ}$ C (30 seconds). After
177 quantitative PCR, a melting curve analysis was performed by slowly increasing the
178 temperature from 65 $^{\circ}$ C to 95 $^{\circ}$ C, with continuous recording of changes in
179 fluorescent emission intensity. To avoid detection of genomic DNA (gDNA), at least
180 one primer per pair was designed to span an exon–exon boundary. All the primers
181 were tested on gDNA and RNA to confirm that potentially contaminant gDNA was not
182 amplified. The specificity was confirmed by melting curve analysis, gel
183 electrophoresis, and sequencing of the quantitative PCR products. Serial dilutions of
184 cDNA pool made from several samples were done to assess PCR efficiency. A ddCt
185 method adjusted for PCR efficiency was used, using the geometric average of
186 H2AFZ (H2A histone family member Z [24]) and GAPDH (glyceraldehyde- 3-
187 phosphate dehydrogenase [25]) as a housekeeping normalization factor [26].
188 Relative expression of cDNA pool from various samples was used as the calibrator to

189 normalize all samples within one PCR run or between several runs. SOX2 data were
190 normalized by a Napierian logarithm transformation.

191

192 2.6. Statistical analysis

193 The effect of superstimulation treatment (control, low FSH, low FSH–LH, high FSH,
194 or high FSH–LH) on the ovulation rate, number of recovered embryos, recovery rate,
195 and fertilization rate was analyzed by ANOVA using a general linear model
196 procedure. For the developmental rate, a probit link with binomial error distribution
197 was used. Data of relative mRNA abundance were normalized by a Napierian
198 logarithm transformation and analyzed using a general linear model. All statistical
199 analyses were performed with SPSS software (SPSS 16.0 software pack- age; SPSS
200 Inc., 2002, Chicago, IL, USA). Results were reported as least-square means with
201 standard error of the mean. Means were separated using Fisher’s protected least
202 significant difference test, with treatment effect declared significant at $P < 0.05$.

203

204

205

206 **3. RESULTS**

207 Effects of superstimulation treatments on recovery variables are shown in
208 Table 2. All the females treated with rhFSH increased the ovulation rate. However,
209 the lack of statistical difference among the low FSH concentration groups and control
210 may be due to the small sample size. Ovulation rate in high FSH concentration
211 groups was significantly higher than that in the control group. Addi- tionally, the
212 ovulation rate was significantly higher when the high FSH dose was supplemented
213 with LH. The number of recovered embryos showed a similar evolution to the

214 ovulation rate. A clear trend toward an increase in the number of recovered embryos
215 was observed in FSH-treated females, although there was no significant difference
216 among low FSH groups and the control. The high FSH groups showed significantly
217 higher numbers of recovered embryos than the control group, although the results
218 from high and low FSH groups without LH supplement were not statistically different.
219 The recovery rate was not affected by superstimulation treatment. Finally, the
220 fertilization rate

221 was significant decreased in the low FSH group supplemented with LH.

222 As shown in Table 3, the superstimulation treatment did not significantly affect
223 in vitro development rate until the expanded blastocyst stage. The proportion of
224 embryo development was very high.

225 Results of superstimulation treatment on embryos OCT4, NANOG, and SOX2
226 gene expression are shown in Figure 2. No significant differences among relative
227 values of OCT4, NANOG, and SOX2 gene expression from different superstimulation
228 treatment were detected.

229

230 **4. DISCUSSION**

231 In the present study, the ovulation rates of females treated with 37.50-IU
232 rhFSH supplemented with 10% rhLH were 3.5 times higher than those observed for
233 the control group and twice of those observed for 18.75-IU rhFSH alone or with 10%
234 rhLH groups. Similar results were reported by Ju et al. [27] with porcine FSH.
235 Nevertheless, the undefined LH concentration present in pituitary extract used by
236 these authors hampered the comparison of trials. In women, an LH concentration
237 window has been described [6,28] below which estradiol production is not adequate
238 and above which LH may be detrimental to follicular development. In a work

239 designed to simplify hormone administration and reduce the stress of treatment and
240 animal handling in rabbits, 25-IU rhFSH alone or in combination with rhLH was
241 diluted in a slow-release polymer (polyvinylpyrrolidone). It was shown that the
242 superovulatory response of does with

243

244 the rhFSH dose used was not dependent on rhLH supplementation [29], and
245 it was concluded that the endogenous LH concentration in rabbits seemed enough to
246 duplicate the ovulation rate of does treated with this FSH dose. A similar ovulation
247 rate was reported previously by other authors using FSH from pituitary glands
248 [11,12,30,31]. Therefore, results of the present work showed that 37.50-IU rhFSH
249 supplemented with 3.75-IU rhLH treatment was significantly more efficient than the
250 other treatments tested, showing an ovulation rate and a number of embryos
251 recovered three times higher than control females. During a normal follicular wave,
252 subordinate follicles regress because of decreasing concentration of circulating FSH.
253 Small follicles require FSH to continue their growth. In the superstimulation protocol,
254 the depressed endogenous FSH caused by the secretory products of dominant
255 follicles is replaced by exogenous FSH. However, a larger number of recruited
256 follicles implies an increase in the LH requirement of mural granulosa cells and,
257 consequently, a higher level of circulating FSH would result in an increased demand
258 of LH as a requirement to improve the number of follicles recruited. Ovulation rate
259 observed with 37.50-IU rhFSH was significantly higher when it was supplemented
260 with 3.75-IU rhLH. It seems that endogenous LH levels from females treated with
261 37.5-IU rhFSH alone were insufficient to increase follicular recruitment to the values
262 achieved when this concentration of FSH was supplemented with LH. This suggests
263 that the window of LH in rabbits is FSH dose dependent, and the higher the con-

264 centration of FSH used, the more the LH window shifts to higher concentrations.
265 When the FSH concentration used is low, 10% of LH supplement may exceed the LH
266 requirements and the fertilization and embryonic development is negatively
267 affected. In contrast, when the FSH concentration used is high, 10% of LH
268 supplementation is essential to increase the number of recruited follicles during the
269 selection phase. These results are in agreement with a study in women using rFSH
270 and rLH which suggest that LH actually sensitizes follicles to FSH and thereby
271 encourages follicular development and hormone secretion. In this study, the
272 number of developing follicles was much reduced or delayed in the group treated
273 with a fixed dose of rFSH without rLH [3]. The present findings seem to be consistent
274 with those of Filicori et al. [32], who found variable effects of LH and FSH on follicles
275 of different sizes. They showed that FSH alone affected the growth dynamic of small
276 follicles and the LH supplementation affected the medium or large follicles,
277 suggesting that expression of LH receptors may provide an additional source of
278 follicular support. In fact, at nonsaturating concentrations of FSH and LH, the
279 response is additive. Therefore, once the appropriate stage of follicular
280 development in response to treatment with FSH is reached, granulosa cells become
281 receptive to LH stimulation and LH becomes capable of exerting its actions on both
282 theca and granulosa cells [33].

283 Regarding embryo quality, to provide a more complete idea of the
284 development potential of the embryos in the present work, in addition to the in vitro
285 development ability of embryos, the OCT4, NANOG, and SOX2 gene expressions
286 were studied. Results of the present work showed that embryo quality, in terms of
287 ability to develop in vitro from

288 eight- to 16-cell stage to the blastocyst stage and OCT4, NANOG, and SOX2
289 gene expression, was not affected by superovulation treatment, with results similar to
290 nontreated females. In cows, it was shown that hormonal superstimulation with
291 porcine or ovine FSH, with high and low LH percentage, respectively, may affect the
292 mRNA profile of some genes [20,34]. Mundim et al. [20] showed differences in gene
293 expression profile of a group of genes related to embryo development between
294 superovulated and control embryos, but no differences were observed compared to
295 embryos produced in vitro. Chu et al. [34] observed that although the number of
296 genes influenced seemed low, the mRNA expression profile of potential factors of
297 developmental competence in the bovine oocyte was affected by follicular
298 stimulation. It is necessary to study whether the source of FSH may affect the
299 expression profile of these genes.

300 In conclusion, results from the present study showed that the treatment of
301 rabbit does with 37.5-IU rhFSH alone or supplemented with rhLH was effective in
302 stimulating superovulation without affecting the embryos ability to develop in vitro
303 until blastocyst and the embryonic expression of OCT4, NANOG, and SOX2 genes.
304 On the other hand, the present study suggests that the rhLH supplementation at
305 this rhFSH concentration increased the number of follicles recruited and the
306 superovulatory response in rabbits.

307

308

309

310 **Acknowledgements**

311

312 This research was supported in part by the Valencian regional government research
313 project PROMETEOII/2014/ 036. The authors would like to thank Neil Macowan
314 Language Services for revising the English version of the article.

315

316 **References**

317

318 [1] Fischer B, Chavatte-Palmer P, Viebahn C, Navarrete Santos A, Duranthon V.
319 Rabbit as a reproductive model for human health. *Reproduction* 2012;144:1–10.

320 [2] Lavara R, Baselga M, Vicente JS. Does storage time in LN2 influence survival
321 and pregnancy outcome of vitrified rabbit embryos? *Theriogenology* 2011;76:652–7.

322 [3] European Recombinant Human LH Study Group. Recombinant human luteinizing
323 hormone (LH) to support recombinant human follicle stimulating hormone (FSH)-
324 induced follicular development in LH- and FSH-deficient anovulatory women: a dose-
325 finding study. *J Clin Endocrinol Metab* 1998;83:1507–14.

326 [4] Shoham Z. The clinical therapeutic window for luteinizing hormone in controlled
327 ovarian stimulation. *Fertil Steril* 2002;77:1170–7.

328 [5] Filicori M, Cognigni GE, Samara A, Melappioni S, Perri T, Cantelli B,
329 et al. The use of LH activity to drive folliculogenesis: exploring un- charted territories
330 in ovulation induction. *Hum Reprod Update* 2002;8:543–57.

331 [6] Balasch J, Fábregues F. LH in the follicular phase: neither too high nor too low.
332 *Reprod Biomed Online* 2006;12:406–15.

333 [7] Aggarwal BB, Farmer SW, Papkoff H, Seidel Jr GE. Biochemical properties of
334 equine chorionic gonadotropin from two different pools of pregnant mare sera. Biol
335 Reprod 1980;23:570–6.

336 [8] Hesser MW, Morris JC, Gibbons JR. Advances in recombinant gonadotropin
337 production for use in bovine superovulation. Reprod Domest Anim 2011;46:933–42.

338 [9] Donaldson LE, Ward DN, Glenn SD. Use of porcine follicle- stimulating hormone
339 after chromatographic purification in super- ovulation of cattle. Theriogenology
340 1986;25:747–57.

341 [10] Zelinski Wooten MB, Hutchison JS, Hess DL, Wolf DP, Stouffer RL. Follicle-
342 stimulating hormone alone supports follicle growth and
343 oocyte development in gonadotropin-releasing hormone
344 antagonist-treated monkeys. Hum Reprod 1995;10:1658–66.

345 [11] Hashimoto S, Kuramochi T, Aoyagi K, Takahashi R, Ueda M, Hirao M, et al.
346 Refined porcine follicle-stimulating hormone promotes the responsiveness of rabbits
347 to multiple-ovulation treatment. Exp
348 Anim 2004;53:395–7.

349 [12] Salvetti P, Theau-Clément M, Beckers JF, Hurtaud J, Guerin P, Neto V,
350 et al. Effect of the luteinizing hormone on embryo production in
351 superovulated rabbit does. Theriogenology 2007;67:1185–93.

352 [13] Lonergan P, Rizos D, Kanka J, Nemcova L, Mbaye AM, Kingston M, et al.
353 Temporal sensitivity of bovine embryos to culture environ- ment after fertilization and
354 the implications for blastocyst quality.
355 *Reproduction* 2003;126:337–46.

356 [14] Gutiérrez-Adán A, Rizos D, Fair T, Moreira PN, Pintado B, De La
357 Fuente J, et al. Effect of speed of development on mRNA expression pattern in early
358 bovine embryos cultured in vivo or in vitro. *Mol Reprod Dev* 2004;68:441–8.

359 [15] Balasubramanian S, Son WJ, Mohana Kumar B, Ock SA, Yoo JG, Im GS, et al.
360 Expression pattern of oxygen and stress-responsive gene transcripts at various
361 developmental stages of in vitro and in vivo preimplantation bovine embryos.
362 *Theriogenology* 2007;68: 265–75.

363 [16] Wrenzycki C, Herrmann D, Niemann H. Messenger RNA in oocyte and embryos
364 in relation. *Theriogenology* 2007;68:77–83.

365 [17] Rizos D, Clemente M, Bermejo-Alvarez P, De La Fuente J, Lonergan P,
366 Gutierrez-Adán A. Consequences of in vitro culture conditions on embryo
367 development and quality. *Reprod Domest Anim* 2008;43: 44–50.

368 [18] Purpera MN, Giraldo AM, Ballard CB, Hylan D, Godke RA, Bondioli KR. Effects
369 of culture medium and protein supplementa- tion on mRNA expression of in vitro
370 produced bovine embryos. *Mol Reprod Dev* 2009;76:783–93.

371 [19] Rizos D, Lonergan P, Boland MP, Arroyo-Garcia R, Pintado B, de la Fuente J, et
372 al. Analysis of differential mRNA expression between bovine blastocysts produced in

373 different culture systems: implications for blastocyst quality. *Biol Reprod*
374 2002;66:589–95.

375 [20] Mundim TC, Ramos AF, Sartori R, Dode MA, Melo EO, Gomes LF, et al.
376 Changes in gene expression profiles of bovine embryos produced in vitro, by
377 natural ovulation, or hormonal superstimulation. *Genet Mol Res* 2009;8:1398–407.

378 [21] Saenz-De-Juano MD, Naturil-Alfonso C, Vicente JS, Marco-Jimenez F. Effect of
379 different culture systems on ARNm expression in developing rabbit embryos.
380 *Zygote* 2013;21:103–9.

381 [22] Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS, Zucker JP, et al. Core
382 transcriptional regulatory circuitry in human embryonic stem cells. *Cell*
383 2005;122:947–56.

384 [23] Theau-Clement M. Preparation of the rabbit doe to insemination: a review. *World*
385 *Rabbit Sci* 2007;15:61–80.

386 [24] Mamo S, Gal AB, Polgar Z, Dinnyes A. Expression profiles of the pluripotency
387 marker gene POU5F1 and validation of reference genes in rabbit oocytes and
388 preimplantation stage embryos. *BMC Mol Biol* 2008;9:67.

389 [25] Navarrete-Santos A, Tonak S, Kirstein M, Kietz S, Fischer B. Two insulin-
390 responsive glucose transporter isoforms and the insulin receptor are developmentally
391 expressed in rabbit preimplantation embryos. *Reprod Res* 2004;128:503–15.

392 [26] Weltzien FA, Pasqualinib C, Vernierb P, Dufoura S. A quantitative real-time RT-
393 PCR assay for European eel tyrosine hydroxylase. *Gen Comp Endocrinol*
394 2005;142:134–42.

395 [27] Ju JC, Chang YC, Huang WT, Tang PC, Cheng SP. Superovulation and
396 transplantation of demi- and aggregated embryos in rabbits. *Asian Australas J Anim*
397 *Sci* 2001;14:455–61.

398 [28] Borini A, Dal Prato L. Tailoring FSH and LH administration to indi- vidual
399 patients. *Reprod Biomed Online* 2005;11:283–93.

400 [29] Viudes-de-Castro MP, Cortell C, Mocé E, Marco-Jiménez F, Joly T, Vicente JS.
401 Effect of recombinant gonadotropins on embryo quality in superovulated rabbit does
402 and immune response after repeated treatments. *Theriogenology* 2009;72:655–62.

403 [30] Kauffman RD, Schmidt PM, Rall WF, Hoeg JM. Superovulation of rabbits with
404 FSH alters in vivo development of vitrified morulae. *Theriogenology* 1998;50:1081–
405 92.

406 [31] Mehaisen GM, Viudes-De-Castro MP, Vicente JS, Lavara R. In vitro and in vivo
407 viability of vitrified and non-vitrified embryos derived from eCG and FSH treatment in
408 rabbit does. *Theriogenology* 2006; 65:1279–91.

409 [32] Filicori M, Cognigni GE, Taraborelli S, Spettoli D, Ciampaglia W, Tabrelli de Fatis
410 C, et al. Luteinizing hormone activity supplementation en- hances follicle-stimulating
411 hormone efficacy and improves ovulation induction outcome. *J Clin Endocrinol Metab*
412 1999;84:2659–63.

413 [33] Oliveira JB, Mauri AN, Petersen CG, Martins AM, Cornicelli J, Cabaña M, et al.
414 Recombinant luteinizing hormone supplementation to recombinant follicle-stimulation
415 hormone during induced ovarian stimulation in the GnRH-agonist protocol: a meta-
416 analysis. *J Assist Reprod Genet* 2007;24:67–75.

417 [34] Chu T, Dufort I, Sirard MA. Effect of ovarian stimulation on oocyte gene
418 expression in cattle. *Theriogenology* 2012;77:1928–38.