

Document downloaded from:

<http://hdl.handle.net/10251/71246>

This paper must be cited as:

Bonet Solves, JA. (2015). The spectrum of Volterra operators on weighted spaces of entire functions. *Quarterly Journal of Mathematics*. 66:799-807. doi:10.1093/qmath/hav019.



The final publication is available at

<http://dx.doi.org/doi:10.1093/qmath/hav019>

Copyright Oxford University Press (OUP)

Additional Information

The spectrum of Volterra operators on weighted spaces of entire functions

José Bonet

Abstract

We investigate the spectrum of the Volterra operator V_g with symbol an entire function g , when it acts on weighted Banach spaces $H_v^\infty(\mathbb{C})$ of entire functions with sup-norms and when it acts on Hörmander algebras A_p or A_p^0 .

1 Introduction, notation and preliminaries

The aim of this note is to investigate the spectrum of the Volterra operator when it acts continuously on a weighted Banach space of entire functions $H_v^\infty(\mathbb{C})$. Our main result is Theorem 2.3. The characterizations of continuous Volterra operators on $H_v^\infty(\mathbb{C})$ obtained by Taskinen and the author in [13] play an important role. The present research originated with a question of A. Aleman in a meeting celebrated in Granada in February 2015 in which the results in [13] were presented. In Section 3 we study the spectrum of Volterra operators acting on a (DFN) Hörmander algebra A_p in Theorem 3.4, and on a Fréchet Hörmander algebra A_p^0 in Theorem 3.5.

In what follows $H(\mathbb{C})$ denotes the space of entire functions. The space $H(\mathbb{C})$ is a Fréchet space endowed with the compact open topology. The differentiation operator $Df(z) = f'(z)$, the integration operator $Jf(z) = \int_0^z f(\zeta)d\zeta$ and the multiplication operator $M_h(f) = hf$, $h \in H(\mathbb{C})$, are continuous on $H(\mathbb{C})$.

Given a non-constant entire function $g \in H(\mathbb{C})$ with $g(0) = 0$, the *Volterra operator* V_g with symbol g is defined on $H(\mathbb{C})$ by

$$V_g(f)(z) := \int_0^z f(\zeta)g'(\zeta)d\zeta \quad (z \in \mathbb{C}).$$

For $g(z) = z$ this reduces to the integration operator. Clearly V_g defines a continuous operator on $H(\mathbb{C})$. The Volterra operator for holomorphic functions on the unit disc was introduced by Pommerenke [28] and he proved that V_g is bounded on the Hardy space H^2 , if and only if $g \in BMOA$. Aleman and Siskakis [3] extended this result for H^p , $1 \leq p < \infty$, and they considered later in [4] the case of weighted Bergman spaces; see also [26]. We refer the reader to the memoir by Peláez and Rättyä [27] and the references therein. Volterra operators on weighted Banach spaces of holomorphic functions on the disc of type H^∞ have been investigated recently in [5]. Constantin started in [15] the study of the Volterra operator on spaces of entire functions. She characterized the continuity of V_g on the classical Fock spaces and investigated its spectrum. Constantin and Peláez

2010 Mathematics Subject Classification. Primary: 47G10, secondary: 30D15, 30D20, 46E15, 47B07, 47B37, 47B38.

Key words and phrases. Spectrum; integral operator; Volterra operator; entire functions; growth conditions; weighted Banach spaces of entire functions; Hörmander algebras

[16] characterize the entire functions $g \in H(\mathbb{C})$ such that V_g is bounded or compact on a large class of Fock spaces induced by smooth radial weights. See also [13]. Aleman and Constantin [1] and Aleman and Peláez [2] investigate the spectra of Volterra operators on several spaces of holomorphic functions on the disc.

A *weight* v is a continuous function $v : [0, \infty[\rightarrow]0, \infty[$, which is non-increasing on $[0, \infty[$ and satisfies $\lim_{r \rightarrow \infty} r^m v(r) = 0$ for each $m \in \mathbb{N}$. We extend v to \mathbb{C} by $v(z) := v(|z|)$. For such a weight, the *weighted Banach space of entire functions* is defined by

$$H_v^\infty(\mathbb{C}) := \{f \in H(\mathbb{C}) \mid \|f\|_v := \sup_{z \in \mathbb{C}} v(|z|)|f(z)| < \infty\},$$

and it is endowed with the weighted sup norm $\|\cdot\|_v$. Changing the value of v on a compact interval does not change the space and gives an equivalent norm. Spaces of this type appear in the study of growth conditions of analytic functions and have been investigated in various articles, see e.g. [10, 11, 22] and the references therein.

For an entire function $f \in H(\mathbb{C})$, we denote $M(f, r) := \max\{|f(z)| \mid |z| = r\}$. Using the notation O and o of Landau, $f \in H_v^\infty(\mathbb{C})$ if and only if $M(f, r) = O(1/v(r))$, $r \rightarrow \infty$. By C, C', c etc. we denote positive constants, the value of which may vary from place to place.

Let $T : X \rightarrow X$ be a continuous operator on a space X . The *resolvent* of T on X is the set $\rho(T, X)$ of all $\lambda \in \mathbb{C}$ such that $T - \lambda I : X \rightarrow X$ is bijective and has a continuous inverse. Here I stands for the identity operator on X . The *spectrum* $\sigma(T, X)$ of T is the complement in \mathbb{C} of the resolvent. The point spectrum is the set $\sigma_{pt}(T, X)$ of those $\lambda \in \mathbb{C}$ such that $T - \lambda I$ is not injective. Observe that in this paper we consider operators defined not only on Banach spaces, but also on more general spaces: $H(\mathbb{C})$ and the Hörmander algebra A_p^0 are Fréchet spaces and the Hörmander algebra A_p is the dual of a Fréchet space. Accordingly, when we refer to a space, we mean a Hausdorff locally convex space. We refer the reader to [25] for results and terminology about functional analysis.

At this point we include some preliminary results, in particular about the spectrum of the Volterra operator on $H(\mathbb{C})$ that are inspired by [1] and [2]. Let $g \in H(\mathbb{C})$ be a non-constant entire function such that $g(0) = 0$ and let $V_g f(z) = \int_0^z f(\zeta)g'(\zeta)d\zeta$, $z \in \mathbb{C}$, denote the Volterra operator associated with g , that acts continuously on $H(\mathbb{C})$.

Proposition 1.1 *The operator $V_g - \lambda I : H(\mathbb{C}) \rightarrow H(\mathbb{C})$ is injective for each $\lambda \in \mathbb{C}$. In particular $\sigma_{pt}(V_g, H(\mathbb{C})) = \emptyset$. Moreover, $0 \in \sigma(V_g, H(\mathbb{C}))$.*

Proof. If $(V_g - \lambda I)f = 0$, then $\lambda f(0) = 0$ and $f(z)g'(z) = \lambda f'(z)$, $z \in \mathbb{C}$. If $\lambda = 0$, we get $f = 0$, since g is not constant. In case $\lambda \neq 0$, we get $f(z) = C \exp(g(z)/\lambda)$, which implies $f = 0$, since $f(0) = 0$.

The operator V_g is not surjective on $H(\mathbb{C})$ because $V_g f(0) = 0$ for each $f \in H(\mathbb{C})$. Thus $0 \in \sigma(V_g, H(\mathbb{C}))$. \square

Lemma 1.2 *Given $\lambda \in \mathbb{C}$, $\lambda \neq 0$, and $h \in H(\mathbb{C})$, the equation $f - (1/\lambda)V_g f = h$ has a unique solution given by*

$$f(z) = R_{\lambda, g} h(z) = h(0)e^{\frac{g(z)}{\lambda}} + e^{\frac{g(z)}{\lambda}} \int_0^z e^{-\frac{g(\zeta)}{\lambda}} h'(\zeta) d\zeta, \quad z \in \mathbb{C}.$$

Proof. This is well known; see e.g. [1, p. 200] or [2, p. 2]. The uniqueness follows from Proposition 1.1. It is enough to substitute in the equation to check the result. Alternatively, the equation implies $f(0) = h(0)$ and that f is the solution of the equation $y' - (g'(z)/\lambda)y = h'(z)$. The result is also obtained solving this linear equation. \square

Proposition 1.3 *Let $g \in H(\mathbb{C})$ be a non-constant entire function such that $g(0) = 0$. The Volterra operator V_g satisfies $\sigma(V_g, H(\mathbb{C})) = \{0\}$ and $\sigma_{pt}(V_g, H(\mathbb{C})) = \emptyset$.*

Proof. This is a direct consequence of Proposition 1.1 and the continuity of the operator $R_{\lambda, g} : H(\mathbb{C}) \rightarrow H(\mathbb{C})$, $\lambda \neq 0$. \square

Lemma 1.4 *Let $X \subset H(\mathbb{C})$ be a locally convex space that contains the constants and such that the inclusion $X \subset H(\mathbb{C})$ is continuous. Assume that $V_g : X \rightarrow X$ is continuous for some non-constant entire function g such that $g(0) = 0$. Then*

$$\{0\} \cup \{\lambda \in \mathbb{C} \setminus \{0\} \mid e^{\frac{g}{\lambda}} \notin X\} \subset \sigma(V_g, X).$$

If X is a Banach space, then

$$\{0\} \cup \overline{\{\lambda \in \mathbb{C} \setminus \{0\} \mid e^{\frac{g}{\lambda}} \notin X\}} \subset \sigma(V_g, X).$$

Proof. Since X contains the constants, V_g is not surjective and $0 \in \sigma(V_g, X)$. If $\lambda \notin \sigma(V_g, X)$, then $\lambda \neq 0$ and the operator $R_{\lambda, g} : X \rightarrow X$ defined in Lemma 1.2 is continuous. In particular, $R_{\lambda, g}(1) = e^{\frac{g}{\lambda}} \in X$. This implies the desired inclusion. Recall that in case X is a Banach space, $\sigma(V_g, X)$ is compact. \square

Lemma 1.5 *Let $X \subset H(\mathbb{C})$ be a locally convex space that contains the constants and such that the inclusion $X \rightarrow H(\mathbb{C})$ is continuous. Assume that $V_g : X \rightarrow X$ is continuous for some non-constant entire function g such that $g(0) = 0$. The following conditions are equivalent:*

- (i) $\lambda \in \rho(V_g, X)$.
- (ii) $R_{\lambda, g} : X \rightarrow X$ is continuous.
- (iii) (a) $e^{\frac{g}{\lambda}} \in X$, and
 - (b) $S_{\lambda, g} : X_0 \rightarrow X_0$, $S_{\lambda, g}h(z) := e^{\frac{g(z)}{\lambda}} \int_0^z h'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta$, $z \in \mathbb{C}$, is continuous on the subspace X_0 of X of all the functions $h \in X$ with $h(0) = 0$.

Proof. This follows directly from the definitions. \square

Lemma 1.6 *Let $X \subset H(\mathbb{C})$ be a locally convex space that contains the constants and such that the inclusion $X \rightarrow H(\mathbb{C})$ is continuous. Let X_0 be the subspace of X of all the functions $h \in X$ with $h(0) = 0$. The following conditions are equivalent for $\lambda \in \mathbb{C} \setminus \{0\}$.*

- (i) $S_{\lambda, g} : X_0 \rightarrow X_0$, $S_{\lambda, g}h(z) := e^{\frac{g(z)}{\lambda}} \int_0^z h'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta$, $z \in \mathbb{C}$, is continuous.
- (ii) $T : X_0 \rightarrow X_0$, $Th(z) := e^{\frac{g(z)}{\lambda}} \int_0^z h(\zeta) g'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta$, $z \in \mathbb{C}$, is continuous.

Proof. This is a direct consequence of the identity

$$e^{\frac{g(z)}{\lambda}} \int_0^z h'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta = h(z) + (1/\lambda) e^{\frac{g(z)}{\lambda}} \int_0^z h(\zeta) g'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta,$$

valid for $h \in H(\mathbb{C})$, $h(0) = 0$, that can be seen integrating by parts. \square

2 Spectra of Volterra operators on $H_v^\infty(\mathbb{C})$

We concentrate our attention in the Volterra operator acting on the Banach space $H_v^\infty(\mathbb{C})$, with $v(r) = \exp(-\alpha r^p)$, where $\alpha, p > 0$. According to [13, Corollaries 3.12 and 3.13], we have the following result.

Proposition 2.1 *Assume that $v(r) = \exp(-\alpha r^p)$, $\alpha > 0$, $p > 0$.*

- (i) $V_g : H_v^\infty(\mathbb{C}) \rightarrow H_v^\infty(\mathbb{C})$ is continuous if and only if g is a polynomial of degree less than or equal to the integer part of p .
- (ii) $V_g : H_v^\infty(\mathbb{C}) \rightarrow H_v^\infty(\mathbb{C})$ is compact if and only if g is a polynomial of degree less than or equal to the integer part of $p - 1$.

Lemma 2.2 *Let v be a weight such that $v(r)e^{\alpha r^n}$ is non-increasing on $[r_0, \infty[$ for some $r_0 > 0$, $\alpha > 0$ and $n \in \mathbb{N}$. The operator $T : H_v^\infty(\mathbb{C}) \rightarrow H_v^\infty(\mathbb{C})$ defined by*

$$T_\gamma h(z) := e^{\gamma z^n} \int_0^z \zeta^{n-1} h(\zeta) e^{-\gamma \zeta^n} d\zeta, \quad z \in \mathbb{C},$$

is continuous if $|\gamma| < \alpha$.

Proof. Changing the value of $v(r)$ on a compact interval, we may assume that $v(r)e^{\alpha r^n}$ is non-increasing on $[0, \infty[$. Fix $z \in \mathbb{C}, z \neq 0$. For each $0 \leq t \leq 1$ we have

$$v(|z|) \leq v(t|z|) e^{\alpha |z|^n (t^n - 1)}.$$

Therefore we can estimate

$$\begin{aligned} v(|z|) |T_\gamma h(z)| &= v(|z|) \left| \int_0^1 z^n t^{n-1} h(tz) e^{\gamma z^n (1-t^n)} dt \right| \leq \\ &\leq |z|^n \int_0^1 t^{n-1} |h(tz)| v(tz) e^{|\gamma| |z|^n (1-t^n)} e^{\alpha |z|^n (t^n - 1)} dt \leq \\ &\leq \|h\|_v |z|^n \int_0^1 t^{n-1} e^{(\alpha - |\gamma|) |z|^n (t^n - 1)} dt = \\ &(1/n) \|h\|_v (\alpha - |\gamma|)^{-1} (1 - e^{-(\alpha - |\gamma|) |z|^n}) \leq (1/n) (\alpha - |\gamma|)^{-1} \|h\|_v, \end{aligned}$$

since $\alpha - |\gamma| > 0$. □

Theorem 2.3 *Assume that $v(r) = \exp(-\alpha r^p)$, $\alpha > 0$, $p > 0$. Let g be a polynomial of degree n less than or equal to the integer part of p with $g(0) = 0$.*

- (i) *If the degree n of g satisfies $n < p$, then $\sigma(V_g, H_v^\infty(\mathbb{C})) = \{0\}$.*
- (ii) *If $p = n \in \mathbb{N}$ and $g(z) = \beta z^n + k(z)$, k a polynomial of degree strictly less than n , then $\sigma(V_g, H_v^\infty(\mathbb{C})) = \{\lambda \in \mathbb{C} \mid |\lambda| \leq |\beta|/\alpha\}$.*

Moreover, we have $\sigma(V_g, H_v^\infty(\mathbb{C})) = \{0\} \cup \overline{\{\lambda \in \mathbb{C} \setminus \{0\} \mid e^{\frac{g}{\lambda}} \notin H_v^\infty(\mathbb{C})\}}$.

Proof. (i) If n is less than or equal to the integer part of $p - 1$, then $V_g : H_v^\infty(\mathbb{C}) \rightarrow H_v^\infty(\mathbb{C})$ is compact. This implies that $\sigma(V_g, H_v^\infty(\mathbb{C})) = \{0\}$, since V_g has no eigenvalues by Proposition 1.1.

Assume now that $p - 1 < n < p$. For each $\lambda \neq 0$, $e^{\frac{g}{\lambda}} \in H_v^\infty(\mathbb{C})$ as it is easy to check.

Suppose first that $g(z) = \beta z^n$ for some $\beta \neq 0$. For $\lambda \neq 0$, take $\alpha > |\beta|/|\lambda|$. Clearly $v(r)e^{\alpha r^n}$ is non-increasing on $[r_0, \infty[$ for some $r_0 > 0$. We can apply Lemma 2.2 to conclude that

$$T_{\beta/\lambda}h(z) := e^{\frac{\beta z^n}{\lambda}} \int_0^z \zeta^{n-1} h(\zeta) e^{-\frac{\beta \zeta^n}{\lambda}} d\zeta, \quad z \in \mathbb{C},$$

is continuous on the subspace of $H_v^\infty(\mathbb{C})$ of the functions vanishing at 0. By Lemma 1.6, the operator

$$S_{\lambda,g}h(z) := e^{\frac{g(z)}{\lambda}} \int_0^z h'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta, \quad z \in \mathbb{C},$$

is continuous in the same Banach space. Finally, Lemma 1.5 implies $\lambda \in \rho(V_g, H_v^\infty(\mathbb{C}))$. This completes the proof in this case.

Suppose now that $g(z) = \beta z^n + k(z)$ for some $\beta \neq 0$ and some polynomial k of degree strictly less than n . Setting $g_1(z) := \beta z^n$, we have $V_g = V_{g_1} + V_k$, and V_k is a compact injective operator on $H_v^\infty(\mathbb{C})$. If $\lambda \neq 0$, we have $V_g - \lambda I = (V_{g_1} - \lambda I) + V_k$, and $V_g - \lambda I$ is bijective if and only if $V_{g_1} - \lambda I$ is bijective. This is a consequence e.g. of [20, Corollary 34.14] keeping in mind that both $V_g - \lambda I$ and $V_{g_1} - \lambda I$ are injective by Proposition 1.1. Therefore, we conclude $\sigma(V_g, H_v^\infty(\mathbb{C})) = \sigma(V_{g_1}, H_v^\infty(\mathbb{C})) = \{0\}$.

(ii) We suppose now that $v(r) = \exp(-\alpha r^n)$, $\alpha > 0$, and that g is a polynomial of degree exactly n .

Again we consider first the case $g(z) = \beta z^n$. For $\lambda \in \mathbb{C} \setminus \{0\}$, we have $e^{\frac{g}{\lambda}} \in H_v^\infty(\mathbb{C})$ if and only if $|\beta|/|\lambda| \leq \alpha$. Therefore, we can apply Lemma 1.4 to conclude that $\{\lambda \mid |\lambda| \leq |\beta|/\alpha\} \subset \sigma(V_g, H_v^\infty(\mathbb{C}))$. Now take $\lambda \in \mathbb{C}$ with $|\lambda| > |\beta|/\alpha$. Since $v(r) \exp(\alpha r^n) = 1$, we apply Lemma 2.2 and Lemma 1.6 to get that the operator

$$S_{\lambda,g}h(z) := e^{\frac{g(z)}{\lambda}} \int_0^z h'(\zeta) e^{-\frac{g(\zeta)}{\lambda}} d\zeta, \quad z \in \mathbb{C},$$

is continuous on the subspace of $H_v^\infty(\mathbb{C})$ of the functions vanishing at 0. Consequently $\lambda \in \rho(V_g, H_v^\infty(\mathbb{C}))$ by Lemma 1.5, which yields $\sigma(V_g, H_v^\infty(\mathbb{C})) = \{\lambda \in \mathbb{C} \mid |\lambda| \leq |\beta|/\alpha\}$ in the present case.

In the general case $g(z) = \beta z^n + k(z)$, $\beta \neq 0$ and some polynomial k of degree strictly less than n , we proceed as in the proof of part (i) to conclude $\sigma(V_g, H_v^\infty(\mathbb{C})) = \sigma(V_{g_1}, H_v^\infty(\mathbb{C}))$, $g_1(z) = \beta z^n$. □

Given a weight v , one defines the following closed subspace of $H_v^\infty(\mathbb{C})$:

$$H_v^0(\mathbb{C}) := \{f \in H(\mathbb{C}) \mid \lim_{|z| \rightarrow \infty} v(|z|)|f(z)| = 0\}.$$

The polynomials are contained and dense in $H_v^0(\mathbb{C})$. By [12, Ex 2.2], the bidual of $H_v^0(\mathbb{C})$ is isometrically isomorphic to $H_v^\infty(\mathbb{C})$. By [13, Corollaries 3.8 and 3.12] V_g is bounded on $H_v^0(\mathbb{C})$ if and only if it is bounded on $H_v^\infty(\mathbb{C})$ when $v(r) = \exp(-\alpha r^p)$, $\alpha, p > 0$. Now, proceeding as in the proof of [5, Lemma 1] or [6, Lemma 2.1] one can show, for $v(r) = \exp(-\alpha r^p)$, that if V_g is bounded on $H_v^\infty(\mathbb{C})$, then it coincides with the bi-transpose $(V_g)''$ of $V_g : H_v^0(\mathbb{C}) \rightarrow H_v^0(\mathbb{C})$. Accordingly, $\sigma(V_g, H_v^0(\mathbb{C})) = \sigma(V_g, H_v^\infty(\mathbb{C}))$.

3 Spectra of Volterra operators on Hörmander algebras

A function $p : \mathbb{C} \rightarrow [0, \infty[$ is called a *growth condition* if it is continuous, subharmonic, radial, increases with $|z|$ and satisfies:

$$(\alpha) \log(1 + |z|^2) = o(p(|z|)) \text{ as } |z| \rightarrow \infty,$$

$$(\beta) p(2|z|) = O(p(|z|)) \text{ as } |z| \rightarrow \infty.$$

Given a growth condition p , we define the following weighted spaces of entire functions (see e.g. [8] and [9]):

$$A_p := \{f \in H(\mathbb{C}) \mid \text{there is } A > 0 : \sup_{z \in \mathbb{C}} |f(z)| \exp(-Ap(z)) < \infty\},$$

endowed with the inductive limit topology, for which it is a (DFN)-algebra (cf. [23]). We also define

$$A_p^0 := \{f \in H(\mathbb{C}) \mid \text{for all } \varepsilon > 0 : \sup_{z \in \mathbb{C}} |f(z)| \exp(-\varepsilon p(z)) < \infty\},$$

endowed with the projective limit topology, for which it is a nuclear Fréchet algebra (cf. [24]).

Clearly $A_p^0 \subset A_p$. Condition (α) implies that the polynomials are dense in A_p and in A_p^0 . Condition (β) implies that both spaces are stable under differentiation. By the closed graph theorem, the differentiation operator D is continuous on A_p and on A_p^0 . It was observed in [7, Lemma 4.1] that the integration operator J is continuous on both spaces.

Weighted algebras of entire functions of this type, usually known as *Hörmander algebras*, have been considered since the work of Berenstein and Taylor [9] by many authors; see e.g. [8] and the references therein. Braun, Meise and Taylor studied in [14], [23] and [24] the structure of (complemented) ideals in these algebras. As an example, when $p(z) = |z|^a$, then A_p consists of all entire functions of order a and finite type or order less than a , and A_p^0 is the space of all entire functions of order at most a and type 0. For $a = 1$, A_p is the space of all entire functions of exponential type, and A_p^0 is the space of entire functions of infraexponential type.

The following Lemma, that is a consequence of condition (β) for the growth condition p , is well known.

Lemma 3.1 *If $p : \mathbb{C} \rightarrow [0, \infty[$ is a growth condition, then there are $M > 0$ and $s > 0$ such that $p(r) \leq Mr^s + M$ for each $r \in [0, \infty[$.*

Proposition 3.2 *Let g be an entire function.*

$$(i) \ V_g : A_p \rightarrow A_p \text{ is continuous if and only if } g \in A_p.$$

$$(ii) \ V_g : A_p^0 \rightarrow A_p^0 \text{ is continuous if and only if } g \in A_p^0.$$

Proof. We show (i), the proof of (ii) being similar. The identities $D \circ V_g = M_{g'}$, $V_g = J \circ M_{g'}$ and the continuity of D and J on A_p imply that $V_g : A_p \rightarrow A_p$ is continuous if and only if the operator of multiplication $M_{g'} : A_p \rightarrow A_p$ is continuous. Since A_p is an algebra containing the constants, this holds if and only if $g' \in A_p$. We can apply again that D and J are continuous on A_p to conclude that this statement is equivalent to $g \in A_p$. \square

The spectrum of the integration operator J on A_p and A_p^0 was investigated in [7, Proposition 4.7]. This result, that corresponds to V_g for $g(z) = z$, is extended in this section.

Lemma 3.3 *Let $p : \mathbb{C} \rightarrow [0, \infty[$ be a growth condition and let h be an entire function.*

- (i) *The function e^h belongs to A_p if and only if $M(h, r) = O(p(r))$ as $r \rightarrow \infty$. If this is the case, then h is a polynomial.*
- (ii) *The function e^h belongs to A_p^0 if and only if $M(h, r) = o(p(r))$ as $r \rightarrow \infty$. If this is the case, then h is a polynomial.*

Proof. (i) If $M(h, r) = O(p(r))$ as $r \rightarrow \infty$, then e^h clearly belongs to A_p . To prove the converse, denote by $A(h, r)$ the maximum of the real part of h in the circle $|z| \leq r$ for $r > 0$. As $e^h \in A_p$, there is $C > 0$ such that $A(h, r) \leq Cp(r) + C$ for each $r > 0$. We apply Carathéodory inequality for the circle $|z| \leq 2r$ (c.f. [21, Theorem 6.8]) and property (β) of p to get

$$M(h, r) \leq 2(A(h, 2r) - \operatorname{Re}h(0)) + |h(0)| \leq 2Cp(2r) + 2(C + |h(0)|) \leq C'p(r) + C'.$$

Now Lemma 3.1 and a standard argument imply that h is a polynomial.

(ii) By [24, Lemma 2.3], $e^h \in A_p^0$ if and only if there is a growth condition q such that $e^h \in A_q$ and $q(r) = o(p(r)), r \rightarrow \infty$. By part (i), $M(h, r) = O(q(r)) = o(p(r)), r \rightarrow \infty$. The reverse implication is trivial. □

Theorem 3.4 *Let $p : \mathbb{C} \rightarrow [0, \infty[$ be a growth condition and let $g \in A_p$ be non-constant.*

- (i) *If $M(g, r) = O(p(r)), r \rightarrow \infty$, is not satisfied (which happens in particular when $p(r) = o(r), r \rightarrow \infty$), then $\sigma(V_g, A_p) = \mathbb{C}$.*
- (ii) *If $M(g, r) = O(p(r)), r \rightarrow \infty$, then $\sigma(V_g, A_p) = \{0\}$. In this case g is a polynomial and $r = O(p(r)), r \rightarrow \infty$.*

Moreover, in both cases we have $\sigma(V_g, A_p) = \{0\} \cup \overline{\{\lambda \in \mathbb{C} \setminus \{0\} \mid e^{\frac{g}{\lambda}} \notin A_p\}}$.

Proof. First observe that $M(g/\lambda, r) = (1/|\lambda|)M(g, r)$ for each $\lambda \in \mathbb{C} \setminus \{0\}$ and $r > 0$. Therefore, it follows from Lemma 3.3 (i) that $e^{\frac{g}{\lambda}} \in A_p$ for some (all) $\lambda \neq 0$, if and only if $e^g \in A_p$, that is equivalent to $M(g, r) = O(p(r))$ as $r \rightarrow \infty$.

(i) If $M(g, r) = O(p(r))$ as $r \rightarrow \infty$ is not satisfied, then $e^{\frac{g}{\lambda}} \notin A_p$ for each $\lambda \neq 0$ by our remarks above. We can apply Lemma 1.4 to conclude $\sigma(V_g, A_p) = \mathbb{C}$.

Observe that in case $p(r) = o(r), r \rightarrow \infty$, then $M(g, r) = O(p(r))$ as $r \rightarrow \infty$ is not satisfied, since otherwise we would have $M(g, r) = O(p(r)) = o(r)$ as $r \rightarrow \infty$, that implies that g is constant; a contradiction.

(ii) If $M(g, r) = O(p(r)), r \rightarrow \infty$, then $e^{\frac{g}{\lambda}} \in A_p$ for each $\lambda \neq 0$. Given $\lambda \in \mathbb{C} \setminus \{0\}$, setting $G = e^{\frac{g}{\lambda}}$, $1/G = e^{-\frac{g}{\lambda}}$, the operator $S_{\lambda, g}$ of Lemma 1.5 (iii) satisfies $S_{\lambda, g} = M_G \circ J \circ M_{1/G} \circ D$. These four operators are continuous on the algebra A_p . Therefore $\lambda \in \rho(V_g, A_p)$, and $\sigma(V_g, A_p) = \{0\}$.

In this case, since g must be a non constant polynomial, the assumption in (ii) implies $r = O(p(r)), r \rightarrow \infty$. □

Theorem 3.5 Let $p : \mathbb{C} \rightarrow [0, \infty[$ be a growth condition and let $g \in A_p^0$ be non-constant.

- (i) If $M(g, r) = o(p(r)), r \rightarrow \infty$, is not satisfied (which happens in case $p(r) = O(r), r \rightarrow \infty$), then $\sigma(V_g, A_p^0) = \mathbb{C}$.
- (ii) If $M(g, r) = o(p(r)), r \rightarrow \infty$, then $\sigma(V_g, A_p^0) = \{0\}$. In this case g is a polynomial and $r = o(p(r)), r \rightarrow \infty$.

Moreover, in both cases, we have $\sigma(V_g, A_p^0) = \{0\} \cup \overline{\{\lambda \in \mathbb{C} \setminus \{0\} \mid e^{\frac{g}{\lambda}} \notin A_p^0\}}$.

Proof. Since $M(g/\lambda, r) = (1/|\lambda|)M(g, r)$ for each $\lambda \in \mathbb{C} \setminus \{0\}$ and $r > 0$, Lemma 3.3 (ii) implies that $e^{\frac{g}{\lambda}} \in A_p^0$ for some (or all) $\lambda \neq 0$ if and only if $e^g \in A_p^0$.

(i) If $M(g, r) = o(p(r))$ as $r \rightarrow \infty$ is not satisfied, then $e^{\frac{g}{\lambda}} \notin A_p^0$ for each $\lambda \neq 0$ by our comments above. By Lemma 1.4 we have $\sigma(V_g, A_p) = \mathbb{C}$.

Observe that in case $p(r) = O(r), r \rightarrow \infty$, then $M(g, r) = o(p(r))$ as $r \rightarrow \infty$ does not hold. Otherwise, $M(g, r) = o(p(r)) = O(r)$ as $r \rightarrow \infty$, which implies that g is constant; a contradiction.

(ii) If $M(g, r) = o(p(r)), r \rightarrow \infty$, then $e^{\frac{g}{\lambda}} \in A_p^0$ for each $\lambda \neq 0$ by our comments above. Given $\lambda \in \mathbb{C} \setminus \{0\}$, setting $G = e^{\frac{g}{\lambda}}$, the operator $S_{\lambda, g}$ of Lemma 1.5 (iii) satisfies $S_{\lambda, g} = M_G \circ J \circ M_{1/G} \circ D$ and is continuous on the algebra A_p^0 . Therefore $\lambda \in \rho(V_g, A_p^0)$.

Since g must be a non constant polynomial, the assumption in (ii) yields $r = o(p(r))$ as $r \rightarrow \infty$. □

Acknowledgement. This research was partially supported by MINECO Project MTM2013-43540-P and by GV Project Prometeo II/2013/013.

References

- [1] A. Aleman, O. Constantin, *Spectra of integration operators on weighted Bergman spaces*, J. Anal. Math. **109** (2009), 199-231.
- [2] A. Aleman, J.A. Peláez, *Spectra of integration operators and weighted square functions*, Indiana Univ. Math. J. **61** (2012), 1-19.
- [3] A. Aleman, A.G. Siskakis, *An integral operator on H^p* , Complex Var. Theory Appl. **28** (1995), 149-158.
- [4] A. Aleman, A.G. Siskakis, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. **46** (1997), 337-356.
- [5] M. Basallote, M.D. Contreras, C. Hernández-Mancera, M.J. Martín, P.J. Paúl, *Volterra operators and semigroups in weighted Banach spaces of analytic functions*, Collect. Math. **65** (2014), 233-249.
- [6] M.J. Beltrán, J. Bonet, C. Fernández, *Classical operators on weighted Banach spaces of entire functions*, Proc. Amer. Math. Soc. **141** (2013), 4293-4303.
- [7] M.J. Beltrán, J. Bonet, C. Fernández, *Classical operators on the Hörmander algebras*, Discr. Cont. Dynam. Syst. **35** (2015), 637-652.
- [8] C. A. Berenstein and R. Gay. *Complex analysis and special topics in harmonic analysis*. Springer-Verlag, New York, 1995.

- [9] C. Berenstein and B. A. Taylor, *A new look at interpolation theory for entire functions of one variable*, Adv. in Math. **33** (1979), 109–143.
- [10] K.D. Bierstedt, J. Bonet, A. Galbis, *Weighted spaces of holomorphic functions on balanced domains*, Michigan Math. J. **40** (1993), no. 2, 271–297.
- [11] K.D. Bierstedt, J. Bonet, J. Taskinen, *Associated weights and spaces of holomorphic functions*, Studia Math. **127** (1998), 137–168.
- [12] K.D. Bierstedt, W. H. Summers, *Biduals of weighted Banach spaces of analytic functions*, J. Austral. Math. Soc. Ser. A **54** (1993), no. 1, 70–79.
- [13] J. Bonet, J. Taskinen, *A note about Volterra operators on weighted Banach spaces of entire functions*, Math. Nachr. (to appear).
- [14] R. W. Braun, *Weighted algebras of entire functions in which each closed ideal admits two algebraic generators*, Michigan Math. J. **34** (1987), 441–450.
- [15] O. Constantin, *A Volterra-type integration operator on Fock spaces*, Proc. Amer. Math. Soc. **140** (2012), 4247–4257.
- [16] O. Constantin, J.A. Peláez, *Integral operators, embedding theorems and a Littlewood-Paley formula on weighted Fock spaces*, arXiv:1304.7501v1.
- [17] O. Constantin, J.A. Peláez, *Boundedness of the Bergman projection on L^p spaces with exponential weights*, arXiv: 1309.6071v2.
- [18] M. Contreras, A.G. Hernández-Díaz, *Weighted composition operators in weighted Banach spaces of analytic functions*, J. Austral. Math. Soc. (Series A) **69** (2000), 41–60.
- [19] A. Harutyunyan, W. Lusky, *On the boundedness of the differentiation operator between weighted spaces of holomorphic functions*, Studia Math. **184** (2008), 233–247.
- [20] G.J.O. Jameson, *Topology and Normed Spaces*, Chapman Hall, London, 1974.
- [21] B. Ya. Levin, *Distribution of zeros of entire functions*. Translated from the Russian by R. P. Boas, J. M. Danskin, F. M. Goodspeed, J. Korevaar, A. L. Shields and H. P. Thielman. Revised edition. Translations of Mathematical Monographs, 5. American Mathematical Society, Providence, R.I., 1980.
- [22] W. Lusky, *On the isomorphism classes of weighted spaces of harmonic and holomorphic functions*, Studia Math. **75** (2006) 19–45.
- [23] R. Meise, *Sequence space representations for (DFN)-algebras of entire functions modulo closed ideals*, J. Reine Angew. Math. **363** (1985), 59–95.
- [24] R. Meise and B. A. Taylor, *Sequence space representations for (FN)-algebras of entire functions modulo closed ideals*, Studia Math. **85** (1987), 203–227.
- [25] R. Meise and D. Vogt. *Introduction to functional analysis*, volume 2 of *Oxford Graduate Texts in Mathematics*. The Clarendon Press Oxford University Press, New York, 1997.
- [26] J. Pau, J.A. Peláez, *Embedding theorems and integration operators on Bergman spaces with rapidly decreasing weights*, J. Funct. Anal. **259** (2010), 2727–2756.

- [27] J.A. Peláez, J. Rättyä, *Weighted Bergman Spaces induced by rapidly decreasing weights*, *Memoirs Amer. Math. Soc.* **227**, 2014.
- [28] Ch. Pommerenke, *Schlichte Funktionen un analytische Functionen von beschränkter mittlerer Oszilation*, *Comment. Math. Helv.* **52** (1977), 591-602.
- [29] J.H. Shapiro, *Composition Operators and Classical Function Theorey*, Springer, New York, 1993.

Author's address:

José Bonet: Instituto Universitario de Matemática Pura y Aplicada IUMPA, Universitat Politècnica de València, E-46071 Valencia, Spain
email: jbonet@mat.upv.es