

Document downloaded from:

<http://hdl.handle.net/10251/78362>

This paper must be cited as:

Casals Missio, J.; Cebolla Cornejo, J.; Rosello Ripolles, S.; Beltran Arandes, J.; Casanas, F.; Nuez Viñals, F. (2011). Long-term postharvest aroma evolution of tomatoes with the alcobaça (alc) mutation. *European Food Research and Technology*. 233(2):331-342. doi:10.1007/s00217-011-1517-6.



The final publication is available at

<http://dx.doi.org/10.1007/s00217-011-1517-6>

Copyright Springer

Additional Information

1 **TITLE**

2 Long-term postharvest aroma evolution of tomatoes with the alcobaça (*alc*) mutation

3

4 **AUTHORS**

5 Joan Casals¹, Jaime Cebolla-Cornejo², Salvador Roselló³, Joaquim Beltán⁴, Francesc Casañas¹, Fernando
6 Nuez²

7

8 ¹ Departament d' Enginyeria Agroalimentària i Biotecnologia, Universitat Politècnica de Catalunya, c/
9 Esteve Terrades 8, Campus del Baix Llobregat, 08860 Castelldefels, Spain.

10 ² Instituto Universitario de Conservación y Mejora de la Agrodiversidad Valenciana (COMAV).
11 Universitat Politècnica de València. Cno. de Vera, s.n. 46022. València. Spain.

12 ³ Departamento de Ciencias Agrarias y del Medio Natural, Universitat Jaume I, Campus de Riu Sec,
13 12071 Castellón, Spain.

14 ⁴ Instituto Universitario de Plaguicidas y Aguas (IUPA), Universitat Jaume I, Campus de Riu Sec, 12071
15 Castellón, Spain.

16

17 **Corresponding author: F. nuez**

18 **Tel. +34-963877421**

19 **Fax: +34-963879422**

20 **e-mail: fnuez@btc.upv.es**

21

22

23

24 **Abstract**

25 The postharvest evolution of Penjar tomatoes has been studied in four accessions representative of the
26 variability of the varietal type. The long term shelf life of these materials, which carry the *alc* allele, was
27 confirmed with 31.2 to 59.1% of commercial fruits after 6 months of effective conservation at room
28 temperature and a limited loss of weight (21.1% to 27.9%). Aroma in Penjar tomatoes is differentiated
29 from other tomato varieties by a characteristic ‘sharp-floral’ aroma descriptor. The evolution of the
30 ‘sharp-floral’ aroma during postharvest showed a peak of intensity at 2 months of postharvest, though in
31 one accession a delay of 2 months in this response was detected. Out of 25 volatiles analysed, including
32 main and background notes, a reverse iPLS variable selection revealed that the main candidates behind
33 this aromatic behaviour are α -terpineol, *trans*-2-hexenal, 6-methyl-5-hepten-2-one, *trans*-2-octenal, α -
34 pinene, β -ionone, 2+3-methylbutanol and phenylacetaldehyde. Between harvest and 2 months postharvest
35 most compounds reduced considerably their concentration, while the intensity of the ‘sharp-floral’
36 descriptor increased, which means that probably there is a rearrangement of the relative concentrations
37 among volatiles that may lead to masking/unmasking processes.

38

39 **Keywords**

40 Alcobaça, aroma, postharvest, ripening mutants, sensory analysis, tomato landrace

41

42 **1. Introduction**

43 More than 400 volatiles have been reported in tomato (*Solanum lycopersicum* L.) [1] and at least 10 of
44 these compounds are required to reproduce its aroma: *cis*-3-hexenal, *cis*-3-hexenol, hexanal, 1-penten-3-
45 one, 3-methylbutanal, *trans*-2-hexenal, 6-methyl-5-hepten-2-one, methyl salicylate, 2-isobutylthiazole
46 and β -ionone [2].

47 The deficient aroma profile of fruits being commercialized at the moment [3] is mainly due to three
48 factors: first, the aroma is a complex polygenic trait with a difficult selection and is usually neglected in
49 breeding programs. Nevertheless it should be noted that the elucidation of volatile precursors [3] and of
50 genes related to the accumulation of volatiles [4,5] open promising opportunities to tomato breeders.
51 Second, handling procedures might play an important role in the aroma profile. In this sense, harvesting
52 in mature-green stage [6] and low temperature storage procedures [7] lead to a decrease in fruit volatile

53 concentrations. Third, breeding for shelf life has had collateral effects, and at the moment it is one of the
54 main causes of the lower aroma levels in modern varieties.

55 In fact, the use of ripening mutants *rin* (ripening inhibitor) [8] and *nor* (non-ripening) [9], which operate
56 upstream of ethylene biosynthesis, increase shelf life with a delay in the ripening process but in return
57 they cause negative effects on aroma profiles lowering the levels of many important volatiles in the red
58 ripe (RR) stage [10-12]. This effect may be a consequence of the impairment of ethylene and lycopene
59 biosynthesis, compounds implied in the metabolic pathways of a great number of volatile compounds [13,
60 14]. Alcobaça (*alc*) is another mutation with a similar effect on ripening [15] and it is allelic to *nor* [16].
61 But this mutation seems to have a lower negative impact on fruit quality [15] and the use of *alc* has been
62 described as a more appropriate strategy than the use of *rin* and *nor* in the development of long-shelf life
63 quality cultivars of tomato [17]. Despite this potential benefit, this mutation has been disregarded in
64 breeding programs, which have been focused on the use of the *rin* mutant mainly in the development of
65 large-sized fresh-market cultivars, and of the *nor* mutant in the case of cherry cultivars [18].

66 In the North East of Spain the *alc* allele is widely distributed in different genetic backgrounds making up
67 a varietal type called Penjar. These tomatoes are characterized by a long shelf life (mean storage ability of
68 126.8 days) and a reduced fruit size (mean fruit weight of 64.1 g). In a recent analysis of the genetic
69 diversity in the varietal type using Amplified Fragment Length Polymorphism (AFLP) a 18.07% of
70 polymorphism was found, revealing the broad genetic base of Penjar landrace [16]. Considering the
71 importance of the genetic background in the aroma profile of tomato fruits, it would be logical to expect
72 that the great diversity found in the Penjar type might lead to considerable differences in the aroma
73 profiles of different accessions, even though all of them carry the *alc* allele.

74 This type of tomatoes is mainly used to prepare ‘pan con tomate’, a traditional dish prepared rubbing the
75 tomato on a slice of toasted bread, and to cook fried tomato sauces. It is usually grown in the open field,
76 harvested during August-October, and it is commercialised during the traditional low-temperature and
77 non-producing period ranging from December to March. This time span represents a conservation period
78 between 2 and 6 months, with storage at room temperature. Local consumers usually consider that Penjar
79 tomatoes have better aroma properties when compared to other tomato varieties, a consideration quite
80 unusual in the appreciation of the aroma of the ripening mutants, and this fact justifies higher selling
81 prices in the local market.

82 There are no detailed works on the effect of the ripening mutant *alc* on tomato aroma, and studies
83 regarding aroma evolution during storage in other varieties are carried only on a short-term basis. The
84 Penjar tomato is a good model to analyse both effects, as it includes a variety of genetic backgrounds and
85 more than 6 months of effective conservation [16]. In this context, the main purpose of this work is to
86 obtain a sensory and analytical description of the aroma of Penjar tomatoes and to track its evolution
87 during its storage (0 to 6 months).

88

89 **2. Material and Methods**

90 **2.1. Plant Material**

91 In previous works, an extensive prospection and collection of accessions belonging to the traditional
92 varietal type Penjar was carried out in its area of cultivation on the East coast of Spain. The collected
93 accessions were characterized examining their morphologic, agronomic and genetic diversity [16]. Using
94 this information four accessions, conserved at the COMAV Seedbank, with an outstanding long shelf life
95 and representing different shapes, colours and agronomic characteristics were selected (Table 1). All
96 these accessions had previously been genetically analysed and the presence of the *alc* allele was
97 confirmed [16].

98

99 **2.2. Field trials**

100 The accessions were cultivated in open field conditions in Castellar del Vallès (UTM: Latitude 41° 36'
101 57''; Longitude 2° 4' 15''; Zone 31). In order to check the homogeneity of growing conditions a
102 randomized complete block design was selected with 4 repetitions and 20 plants per plot. Cultivation was
103 carried out using the traditional practices applied for tomato cultivation in the area, including drip
104 irrigation, staking, fortnight pruning, integrated pest management and initial manure fertilization. The
105 characteristics of the accessions were checked and mean yield, mean fruit weight, soluble solids (°Brix),
106 fruit colour (visual estimation), fruit shape, fruit blossom end shape and other interesting traits were
107 recorded. Yield was recorded in 20 randomly selected plants per accession, while fruit traits were
108 evaluated in 20 randomly selected fruits from different plants per accession. All the fruits from the second
109 to the fourth truss were harvested and stored in darkness at room temperature ($20 \pm 5^\circ\text{C}$) and humidity
110 (68-75% relative humidity). During postharvest, a screening of the fruits was performed every two weeks.
111 Fruits were discarded if they showed external signs of desiccation, loss of turgor or fungal infection, the

112 rest of the fruits were considered commercial. Shelf life was calculated as the percentage of commercial
113 fruits at 6 months of postharvest storage. The percentage loss of weight was determined at 2, 4 and 6
114 months of postharvest storage using 16 fruits per accession, on a per fruit basis.

115

116 **2.3. Sample preparation and aroma analysis**

117 2.3.1. Sample preparation

118 Samples were obtained at harvest (0 months postharvest) and at 2, 4 and 6 months of postharvest storage.
119 Each sample was kept frozen in order to analyse the aromatic profile of the whole collection at the same
120 time and in the same conditions. Each sample was made up by 10 fruits with good conservation (without
121 external signs of deterioration) and with weights near to the estimated mean weight calculated for the
122 accession (Table 1). The lack of internal bruising was established as an additional criterion in order to
123 select the fruits for the sample [19]. The lignified area surrounding the pedicel scar was discarded and the
124 fruits were ground and homogenized, adding a saturated solution of CaCl₂ to inactivate volatile degrading
125 enzymes [20]. Samples were instantly kept frozen at -80°C until analysis.

126

127 2.3.2. Sensory analysis

128 Sensory analysis was conducted to discriminate the odour between accessions and between postharvest
129 storages (0, 2, 4 and 6 months). Sensory analysis was performed with 10 trained panelists with previous
130 experience in tomato and bean evaluation [21]. The panelists were specifically trained to evaluate tomato
131 odour descriptors using Penjar populations. Firstly, in order to reach a consensus in the odour descriptors
132 more appropriate for Penjar tomatoes, the panelists were presented during 4 sessions with Penjar tomato
133 samples with 2 and 4 months of postharvest storage, as well as with samples belonging to commercial
134 fresh tomatoes obtained from the local market (4 sessions). These sessions enabled an initial consensus on
135 a limited set of odour descriptors. During other 8 sessions, the panelists were presented with numerous
136 samples including different genotypes and storage periods in order to get familiar with the range of
137 variation in the intensity of the selected descriptors. Finally during 2 additional sessions the optimal
138 serving temperature was evaluated. Four collections with 0, 2, 4 and 6 months of postharvest storage were
139 evaluated at four different serving temperatures: 15, 17.5, 20 and 25 °C.

140 Once the best serving temperature was selected, the following thawing procedure was adopted: samples
141 were taken out of the ultra-low freezer (-80°C) the day before the evaluation session and hermetically

142 sealed and placed in a refrigerator (8°C) during 12 hours. Three hours before the evaluation session, the
143 samples were introduced in a chamber at 20°C.

144 Tasting sessions were carried out twice a week in a room designed for sensory analyses (ISO 8589) that
145 was illuminated with green light to mask the color of the samples. Accessions were evaluated in
146 quadruplicate and were randomly distributed in 16 sessions (4 accessions per session). The samples were
147 presented in sealed cylindrical vials (diameter: 50 mm; height: 43 mm). Vials were unsealed 2 minutes
148 before starting the sensory analysis. All scoring took place on a semi-structured scale ranging from 0 to
149 10 with the endpoints anchored and marked with the descriptors.

150

151 2.3.3. Volatile analysis

152 Twenty five tomato volatiles were chromatographically determined in the samples: 2-phenylethanol,
153 *trans*-2-hexenal, 2-isobutylthiazole, 6-methyl-5-hepten-2-one, 2+3-methyl-1-butanol, hexanal, 1-hexanol,
154 *cis*-3-hexenol, *cis*-3-hexenal, *trans*-2-heptenal, R-limonene, nonanal, eugenol, geranyl acetone, methyl
155 salicylate, linalool, guaiacol, β -ionone, *trans*-2-octenal, α -pinene, phenylacetaldehyde, benzaldehyde, α -
156 terpineol, camphor, and β -cyclocitral. Reference aroma compounds were obtained from Sigma-Aldrich
157 Química S.A. (Madrid, Spain) as pure compounds. Stock solutions of the aroma standards at 500 mg L⁻¹
158 were prepared in acetone and stored at -18°C. Working solutions were prepared by volume dilution in
159 diethyl ether-hexane (1:1). The internal standard methyl salicylate-D₄ of 99.5% purity was purchased
160 from Sigma-Aldrich Química S.A. (Madrid, Spain). Calcium chloride 97% (Riedel de Haen) was
161 purchased from Supelco (Sigma-Aldrich Química S.A., Madrid, Spain). Organic solvents (hexane, ethyl
162 acetate, diethyl ether) of trace residue analysis quality were purchased from Scharlab (Barcelona, Spain).
163 SPE cartridges (Supelco, Sigma-Aldrich Química S.A., Madrid, Spain) were prepared by the
164 manufacturer packing 500 mg of Tenax TA (80-100 mesh,) in 6 mL polyethylene cartridges retained
165 using two polietilene frits.

166 The extraction system developed in a previous work [22] consisted in a 50 mL Erlenmeyer flask attached
167 to a glass cap with two connexion tubes: the inlet connected to a dry N₂ gas supply, and the outlet fitted to
168 the Tenax trap. Dry nitrogen (99.7%) was used to carry out the purge process, and was led to flow into the
169 flask at a flow of 1 L min⁻¹. 30 g of tomato sample together with a 5 % (w:w) of CaCl₂ and with addition
170 of 50 μ L of 15 μ g mL⁻¹ methyl salicylate-D₄ (surrogate/internal standard) were magnetically stirred (350
171 rpm) and heated at 35 °C for 120 min in order to allow the volatile analytes to be retained in the Tenax

172 trap (maintained at ambient temperature). The trap was removed and eluted with 3.5 mL of hexane-ether
173 (1:1) mixture. The final volume extract was adjusted to 1 mL by means of a gentle stream of nitrogen.
174 Chromatographic determination was carried out using a Varian CP-3800 gas chromatograph (Varian Inc.
175 Palo Alto, USA) coupled to an ion trap mass spectrometry detector (Saturn 4000, Varian Inc. Palo Alto,
176 USA). Separation of the analytes was carried out on a 30 m x 0.25 mm DB-5MS (0.25 μm film thickness)
177 Varian capillary column, using helium at a constant flow of 1 mL min^{-1} as carrier gas. The temperature
178 program was as follows: 45 $^{\circ}\text{C}$ for 5 min, then raised to 96 $^{\circ}\text{C}$ at a rate of 3 $^{\circ}\text{C min}^{-1}$, then raised to 150 $^{\circ}\text{C}$
179 at a rate of 6 $^{\circ}\text{C min}^{-1}$, and finally raised up to 240 $^{\circ}\text{C}$ at a rate of 30 $^{\circ}\text{C min}^{-1}$, with a final isothermal stage
180 of 1.5 min (total chromatographic analysis time of 36 min). Injection in the splitless mode of a volume of
181 1 μL (injection port temperature 200 $^{\circ}\text{C}$, splitless time 1 minute) was carried out using an autosampler
182 Varian 8400 (Varian Inc. Palo Alto, USA) equipped with a 10 μL syringe. The gas-chromatograph was
183 directly interfaced with the Varian 4000 mass-spectrometer, ion trap, (Varian Inc. Palo Alto, USA) in the
184 external ionization mode with an electron ionization energy of 70 eV in the positive ion mode. Transfer
185 line temperature was established at 250 $^{\circ}\text{C}$ and ion source and trap temperatures were adjusted to 200 $^{\circ}\text{C}$.
186 Quantitation of analytes in the sample extracts was performed using a external calibration curve obtained
187 after direct injection of solvent standards containing internal standard and plotting relative areas to
188 internal standard methyl salicylate-D4 against concentration (ng mL^{-1}) as described by Beltran et al. [22].
189 Quantitation ion used for the internal standard methyl salicylate-D4 was 155. This ion corresponded to
190 the molecular mass of the compound after having changed the deuterium in the alcohol group by
191 hydrogen, which occurs due to the contact with the aqueous sample.

192

193 **2.4. Statistical analysis**

194 For sensory data analysis ANOVA procedure was conducted using SAS statistical package v.8.02 (SAS
195 Institute Inc, Cary, NC, USA). A lineal model considering all the factors and their interactions was
196 selected: $x_{ijk} = \mu + \alpha_i + \beta_j + \gamma_k + s_l + \alpha\beta_{ij} + \alpha_i\gamma_k + \beta_j\gamma_k + \alpha_i\beta_j\gamma_k + \epsilon_{ijk}$, where α_i = panelist, β_j =accession,
197 γ_k =postharvest storage, s_l = session (random factor) and $\alpha\beta_{ij}$, $\alpha_i\gamma_k$, $\beta_j\gamma_k$ and $\alpha_i\beta_j\gamma_k$ are the interactions
198 between fixed factors. A Student-Newman-Keuls mean comparison test was performed after checking
199 effect significance with the ANOVA.

200 To perform the statistical analysis of the concentrations of the volatile compounds being determined,
201 logodor units were calculated using commonly accepted odour thresholds for all volatiles. This

202 transformation was selected to scale the relative importance of each compound in aroma perception. In
203 order to study the relation between sensory data and volatile composition a Partial Least Square (PLS)
204 regression was used [23]. Prior to the PLS regression, the data were autoscaled with mean-centering and
205 division by the standard deviation of the variable [24] to avoid the distortion caused by different variable
206 scaling. The PLS regression model was calculated using full crossvalidation resampling method. The
207 goodness of the model fit was tested using the Root Mean Square Error of Calibration (RMSEC) and the
208 Root Mean Square Error of Cross Validation (RMSECV).

209 In order to select the number of latent variables of the PLS model two criteria were used: an additional
210 latent variable was only chosen when the RMSECV was improved by at least 2% and the number of new
211 variables was minimized as possible. In order to improve model precision an aromatic variable selection
212 was performed using an Interval PLS (iPLS) variable selection which performs a hierarchical, sequential
213 and exhaustive search for the best combinations of variables. iPLS was performed in reverse mode, with
214 intervals successively removed from the analysis [24].

215 The calculations of PLS regressions were made using PLS_Toolbox v 6.0 (Eigenvector Research Inc,
216 Wenatchee, WA, USA) for Matlab v 7.6.0 (Mathworks Inc, Natick, MA, USA).

217

218 **3. Results**

219 **3.1. Shelf life evolution**

220 Field trials confirmed that there were no statistical agro-morphological differences between blocks, thus
221 samples from the same accession were pooled. Postharvest storage behaviour (Table 2) showed
222 significant differences between accessions. The highest shelf life was recorded in accession CDP-1245,
223 which showed 59.1 % of commercial fruits after 6 months of conservation. A value that was significantly
224 different to that of accession CDP-5468, which showed the lowest shelf life (31.2 %). Accessions CDP-
225 1240 (42.4 %) and CDP-8268 (42.8 %) showed no significant differences between them and between the
226 rest of accessions. The higher weight loss was detected in the accession CDP-1245, with 12.1%, 19.2%
227 and 27.9% of weight loss at 2, 4 and 6 months postharvest respectively, values significantly higher than
228 the weight loss recorded for CDP-1240 and CDP-5468 and CDP-8268 at 6 months postharvest.

229

230 **3.2. Panel training and consensus of odour attributes**

231 With the lexicon proposed by Hongsoongnern and Chambers [25] as a starting point, different descriptors
232 were suggested by the panel to describe the odour perceived in the accessions assayed. Panelists
233 identified a characteristic odour in most of the Penjar tomatoes samples, and it was described as ‘sharp’
234 with ‘floral notes’. Other descriptors cited by the panelists in the Penjar samples were: ‘green’,
235 ‘fermented’, ‘pharmaceutical’ and ‘earthy’. Out of all these descriptors, only the odours ‘sharp-floral’ and
236 ‘earthy’ were not found in the samples of commercial standard fresh tomatoes. These descriptors also
237 appeared in different intensities in the different accessions and storage periods. The odour descriptor
238 ‘sharp-floral’ was the most cited by the panelists during the training sessions. Other suggested descriptors
239 were discarded: ‘earthy’ was considered as important but not frequent, the odour descriptors ‘fermented’
240 and ‘pharmaceutical’ were judged as negative and the odour descriptor ‘green’ was judged as occasional.
241 Therefore, the rest of the training and the evaluation sessions were performed using only the descriptor
242 ‘sharp-floral’. During the training, all the panelists indicated that the aromas were better perceived at
243 20°C among the four temperatures tested, and this serving temperature was selected for the sensory
244 analysis.

245

246 **3.3. Sensory analysis**

247 The odour descriptor ‘sharp-floral’ increased its intensity during postharvest storage of the Penjar
248 tomatoes ($p < 0.0001$), with a maximum observed at 2 months of postharvest storage (Figure 1). After this
249 peak (4 months postharvest) the intensity of this descriptor decreased to similar values to those recorded
250 at the harvest (0 months postharvest). Finally, at 6 months postharvest the intensity of the ‘sharp-floral’
251 descriptor was very low in all the accessions. Out of the four accessions assayed, accessions CDP-1240
252 and CDP-5468 recorded the highest intensities of the ‘sharp-floral’ descriptor with higher values than
253 CDP-1245 at 0, 2 and 4 months postharvest and to CDP-8268 at 2 months postharvest ($p < 0.0001$). Only
254 accession CDP-8268 showed a different pattern in the evolution of aroma perception, with a maximum
255 intensity of the ‘sharp-floral’ descriptor at 4 months postharvest. This unusual delay caused the
256 significance of the accession x postharvest storage interaction ($p = 0.0229$).

257

258 **3.2. Volatile compounds**

259 Twenty four volatiles were detected in the samples analyzed. *Cis*-3-hexenal remained under detection
260 limits in all the accessions and storage periods. This absence was unusual as it has been considered as one
261 of the main aroma volatiles in other tomato varieties [2].

262 At the harvest (0 months postharvest storage), the compound with the highest concentration was 2-
263 phenylethanol (Table 3). Other abundant compounds were *trans*-2-hexenal, *cis*-3-hexenol, hexanal and 2-
264 isobutylthiazole. Accessions CDP-5468 and CDP-1240 registered the higher concentrations of volatiles at
265 harvest, and 4 of the most important volatiles, including, *cis*-3-hexenol, *trans*-2-hexenal, hexanal and 2-
266 isobutylthiazole, reached a concentration more than 5 times higher than those found in the accessions
267 CDP-1245 and CDP-8268.

268 The data obtained for postharvest storages of 2, 4 and 6 months showed that there is a generalized
269 decrease in the concentration of all the volatiles determined, excluding some cases such as nonanal and α -
270 pinene, with very low concentration at harvest. The most important reduction in the concentration
271 occurred during the period between harvest and 2 months postharvest, when a mean reduction of 50%
272 was registered (Table 3), except for accession CDP-1245 where, in average, no considerable reduction
273 was recorded in this period, a result probably related to the smaller concentrations detected at harvest in
274 this accession. After this initial reduction, between 2 and 4 months postharvest the decrease in
275 concentration was small. Finally, in most cases concentration remained stable between 4 and 6 months.

276 In order to obtain a better interpretation of the relation between volatile composition and the sensory
277 perception by the panelists, a PLS analysis using all the detected volatile components was carried out.
278 The two first latent variables were selected to minimize calibration (RMSEC) and crossvalidation
279 (RMSECV) errors. With the first two latent variables the model captured a 64.53% of the variation of
280 sensory panel response using 62.89% of the variation in the volatiles composition matrix. The
281 determination coefficient obtained in the calibration model was moderate ($R^2=0.63$) with a REMSEC of
282 1.08 and a RMSECV of 1.69 sensory units. The first latent variable was positively correlated with all
283 the volatiles with similar loadings, but negatively correlated with α -pinene. The second latent variable
284 was positively correlated mainly with volatiles 1-hexanol, hexanal and phenylacetaldehyde mainly and
285 negatively correlated with volatiles camphor, α -terpineol, 2-phenylethanol, linalool and β -ionone.

286 Despite the good prediction response, the model still could not clearly establish which of the original
287 variables were really important to explain the variability of the sensory panel response. Therefore, a
288 selection of a subset of aromatic compounds were performed using reverse Interval PLS (iPLS) [26] in

289 order to obtain a superior prediction model. The results of the iPLS variable selection indicated that the
290 main volatiles related with the variation in the sensory matrix were α -terpineol, *trans*-2-hexenal, 6-methyl-
291 5-hepten-2-one, *trans*-2-octenal, α -pinene, β -ionone, 2+3-methylbutanol and phenylacetaldehyde. Using
292 these set of volatiles, the model minimized RMSEC and RMSECV with the two first latent variables,
293 which captured 65.19% of the variation in the sensory matrix using 73% of the variation in the volatiles
294 matrix. A higher determination coefficient was obtained ($R^2=0.73$) with lower errors (RMSEC=0.93
295 sensory units and RMSECV=1.33 sensory units). Thus, the reduction in the number of initial volatiles
296 enabled the development of a better model, confirming the good selection of the main volatiles involved
297 in the sensory matrix variation. This time, the first component was positively correlated with similar
298 loadings with volatiles *trans*-2-hexenal, 6-methyl-5-hepten-2-one, *trans*-2-octenal, 2+3-methylbutanol,
299 phenylacetaldehyde and β -ionone and with a lower loading with α -terpineol and again negatively
300 correlated with volatile α -pinene (Table 4). The second latent variable was positively correlated with
301 volatiles α -pinene, 2+3-methylbutanol and phenylacetaldehyde and negatively with volatiles 6-methyl-5-
302 hepten-2-one, *trans*-2-octenal and β -ionone; a value close to 0 was obtained for volatile *trans*-2-hexenal
303 (Table 4).

304 In the PLS model obtained (Figure 2) it was easier to identify clusters of points associated to postharvest
305 storage duration than to accessions. The points corresponding to the peaks of intensity of the odour
306 descriptor 'sharp-floral' were clustered in the upper right quarter of the graph, even the point
307 corresponding to the intensity peak of the accession CDP-8268 that showed an unusual delay in the
308 response was in the same area. Other samples with high values of 'sharp-floral' intensity (Figure 1) were
309 also clustered in the same quarter (Figure 2). This was the case of the accession CDP-1240 at 4 months
310 postharvest and of the accession CDP-5468 at harvest. Accession CDP-1240 at harvest with high
311 intensity in the descriptor (Figure 1) was placed in the lower-right quarter, but close to the other samples
312 with high intensity. In the upper right quarter of the model only accessions with high 'sharp-floral'
313 intensity could be found (Figure 2).

314

315 **4. Discussion**

316 As expected, a considerable variation in shelf life was detected among the accessions assayed. Although
317 all of them offered good conservation in long-term storage, it was possible to identify outstanding
318 accessions such as CDP-1245 with almost 59.1% commercial fruits after 6 months of storage at room

319 temperature. The differences detected confirmed the good selection of the materials as the objective was
320 to evaluate a representative sample of the variation in the varietal type. It should be noted the good
321 response of the Penjar tomatoes, especially if the loss of weight is compared with results provided by
322 other authors. In this sense, Javanmardi and Kubota [27] reported a loss of weight ratio at room
323 temperature of 0.68% per day, and that would mean a 40.8% in two months, while in our study Penjar
324 tomatoes showed only a 9.0%-12.1% reduction in this period.

325 Despite different aroma notes such as 'green', 'sharp', 'floral', 'earthy', 'fermented' and 'pharmaceutical'
326 being identified in the collection of Penjar tomatoes with the *alc* mutation, it was the 'sharp with floral
327 notes' descriptor the one that clearly and continuously was associated to this particular varietal type. This
328 descriptor would represent an 'identification mark' for the varietal type as it was not found in reference
329 commercial fresh tomato varieties. The intensity of this descriptor, as expected, varied during postharvest
330 storage, reaching a maximum not at harvest, but generally at 2 months postharvest. This is an unusual but
331 interesting result, as it is usually suggested that a reduction of postharvest storage minimizes the typical
332 loss of the characteristic tomato aroma [28, 29].

333 The existence of a characteristic odour descriptor possibly contributes to the preservation of a local
334 market associated to this varietal type, as well as to the association of the variety with traditional dishes.
335 On the other hand, the identification of intensity peaks for the descriptor enables the determination of the
336 best moment to release the stored materials with the maximum quality. In general, the best aromatic
337 properties would be obtained at 2 months postharvest.

338 The fact that Penjar varietal type is formed by a wide variety of genetic backgrounds, in which the *alc*
339 allele has been inserted, enabled the identification of accessions with high odour scores, such as CDP-
340 1240 and CDP-5468. It also enabled the identification of unusual patterns of aroma evolution. In this
341 sense, the accession CDP-8268 showed a delay in the 'sharp-floral' descriptor intensity at 4 months
342 instead of the 2 months peak identified in the rest of the accessions.

343 The existence of genotypic variability among the Penjar tomatoes, as odour intensity is concerned, also
344 leads to a further conclusion related to the structure of traditional or landrace populations. It is known that
345 these materials are usually configured as population varieties with a high level of diversity, maintained
346 through mass selection processes. It is also known that the materials that have survived the genetic
347 erosion processes are usually related to quality markets because the consumer identifies in them a higher
348 level of organoleptic quality. In the case of the Penjar tomato, the main morpho-agronomic characteristic

349 of the varietal type is due to its long shelf life as a consequence of the introgression of the *alc* allele in
350 different varietal types [16]. Therefore this is the characteristic that has been traditionally associated with
351 a higher organoleptic quality. But, the considerable variation in odour intensity detected in this work
352 results in the existence of low quality populations, which are probably maintained in the market through
353 the generalization of a higher quality traditionally assigned to the varietal type. The association of the
354 ideas 'traditional' and 'high quality' is not always true, especially in species such as the tomato where the
355 existence of a certain degree of cross-pollination may contribute to varietal degeneration. Therefore, in
356 order to consolidate quality markets and to promote on-farm conservation of these genetic resources it is
357 necessary to purge the existing populations, fostering those with better organoleptic profiles.

358 Regarding volatile concentration, it is unusual to find tomato fruits with low levels of *cis*-3-hexenal as in
359 this case. This compound has been described as the most important in tomato in several studies [20, 30,
360 31], with a major contribution to the aroma descriptors 'fresh green', 'sweet' [30] and 'tomato-like' [31].
361 It has been reported the instability of *cis*-3-hexenal and its isomerization to *trans*-2-hexenal during
362 isolation and analysis [20], though it does not seem that this is the case of this study. In fact, we have
363 found *cis*-3-hexenal using exactly the same methodology in other tomato varieties [32]. The absence of
364 this compound may be important in the characteristic aroma of the Penjar tomatoes, as it may be related
365 to the emergence or unveil of other compounds which typically-show lower logodor units.

366 Apart from the deficiency in *cis*-3-hexenal it does not seem that the introgression of the *alc* allele affects
367 the concentration of other volatiles, as it has been reported in the ripening mutant *nor* [10-12], which is
368 allelic to *alc* [16]. The comparison of the results obtained in this study and the analyses performed with
369 the same methodology or the previously published results by other groups in other varietal types [2, 33,
370 34], apart from the lack of *cis*-3-hexenal, only evidenced reduced levels of hexenal and
371 phenylacetaldehyde.

372 The lightness of the external colour typical of this varietal type made logical to expect reduced levels of
373 volatiles derived from the carotenoid degradation pathway such as 6-methyl-5-hepten-2-one and geranyl
374 acetone[14], especially considering that the *alc* mutation has been related to low levels of this carotenoid
375 [15]. But on the contrary, the values obtained in the Penjar tomatoes at harvest (Table 1) were similar to
376 those reported by other authors in conventional varieties: 0.13 mg kg⁻¹ [2], 0.1 to 0.3 mg kg⁻¹ [20] or 0.05
377 to 0.2 mg kg⁻¹ [33] in the case of 6-methyl-5-hepten-2-one, and 0.057 mg kg⁻¹ [2] in the case of geranyl
378 acetone. It should also be highlighted that the concentration obtained of 2-isobutylthiazole at harvest in

379 the accessions CDP-1240 and CDP-5468 (Table 1) is more than 10 times higher than the previously
380 reported in other varieties: 0.04 mg kg⁻¹ [2], 0.01 mg kg⁻¹ [6] or 0.03 mg kg⁻¹ [33].

381 In some fruits a single compound dominates aroma perception, but in tomato no single compound
382 dominates and more than 10 volatiles have been described as having positive log odour units. Even
383 compounds with negative logodor units should not be neglected, as they may still contribute to the overall
384 flavour as background notes [11]. It has even been determined that some of the last, such as eugenol, may
385 have an impact on tomato aroma upon release from their glycosidic conjugates [6].

386 In this complex context, with so many compounds, and relations between them, conditioning odour
387 perception, it is extremely difficult to elucidate a direct relation between aroma perception by the
388 panelists and volatile composition of the fruit, and its evolution during storage period. The best alternative
389 found was to carry out Partial Least Square regression (PLS) analysis. PLS attempts to find factors which
390 both capture the greatest amount of variance in the aromatic composition and achieve the best correlation
391 between the panel 'striking' odour intensity evaluation (predicted variable) and the volatile composition
392 matrix (predictor variables) including storage evolution. In other words, PLS maximize covariance
393 between predictor and predicted variables. This statistical procedure is frequently used in several complex
394 chemometric applications, and has also been applied to identify the most important descriptors in aroma
395 perception [35]. Following this methodology, optimized with iPLS variable selection, the volatiles α -
396 terpineol, *trans*-2-hexenal, 6-metyl-5-hepten-2-one, *trans*-2-octenal, α -pinene, β -ionone, 2+3-
397 methylbutanol and phenylacetaldehyde were identified as important compounds to consider in order to
398 explain the postharvest odour evolution of the Penjar tomatoes.

399 The contribution of each compound to the descriptor is really difficult to ascertain. Several compounds
400 may change the induced aroma perception at different concentrations and some of them may interact with
401 others masking or unmasking aroma notes [1]. Additionally, not only each compound may be responsible
402 for different attributes at different concentrations, but their perception may vary with changes in alcohol
403 content such as the increase in ethanol during ripening and this may add complexity to tomato aroma
404 evaluation [31].

405 Regarding the perception of the selected volatiles, α -terpineol has been described as 'floral/fruity' [36],
406 *trans*-2-hexenal might induce a 'green' or 'stale' perception [31], 6-metyl-5-hepten-2-one as 'sweet-
407 floral' [31], *trans*-2-octenal as 'sweet/phenolic' [37], α -pinene as 'stem-like' [38], β -ionone as 'sweet

408 fruity' [31], 2+3-methylbutanol as 'tomato-like' [39], and phenylacetaldehyde as 'sweet' [30]. In short,
409 most of them may contribute to the 'sharp-floral' descriptor found in the Penjar tomatoes.

410 In the PLS model the first latent variable had positive and similar loadings with almost all these selected
411 volatiles and it may be related with overall volatile content, while in the second latent variable 5 volatiles
412 had negative loadings and 3 had positive loadings, and it would be related to aroma nuance. As the
413 samples corresponding to the higher 'sharp-floral' intensity had positive values of the first two latent
414 variables of the optimized PLS model (figure 2), a higher impact would be ascribed to volatiles with high
415 loadings in both latent variables. This was the case of 2+3-methylbutanol and phenylacetaldehyde (Table
416 4). Nevertheless it may also be possible that some of the compounds with negative loadings in the second
417 latent variable might be masking other compounds, and thus should not be disregarded. It should also be
418 pointed that between harvest and 2 months postharvest most compounds reduced considerably their
419 concentration, while the intensity of the 'sharp-floral' descriptor increased, which means that probably
420 there is a rearrangement of the relative concentrations among volatiles that may lead to
421 masking/unmasking processes.

422 Berna et al. [38] studying the evolution of aroma profiles from harvest to 19 days postharvest storage
423 reported an initial shift with terpenoids, produced in the stem, holding an important participation in the
424 overall aroma at the beginning of conservation, to a more important role of compounds such as 1-
425 nitropentane and 6-methyl-5-hepten-2-one related to fresh tomato and fruity aroma respectively as storage
426 progressed. They also found an increase of 2-methylbutanol at ending stages of maturity.

427 It is difficult to extrapolate similarities between these findings related to the first weeks of conservation
428 and our work, as the Penjar tomatoes are adapted to longer storage periods and therefore time span
429 evaluated is much larger. Nevertheless it is interesting to see that compounds selected as important in the
430 evolution of the aroma profiles with the reverse-iPLS such as 6-methyl-5-hepten-2-one and 2+3-
431 methylbutanol are highlighted in both studies.

432 Krumbein et al. [40] monitoring the postharvest aroma evolution during 21 days on different cultivars,
433 some of them with reported long shelf life, found that the increase in hexanal and 2-isobutylthiazole
434 during postharvest was connected with an increase of the mouldy descriptor, whereas the attribute
435 tomato-like increased simultaneously, maybe linked with the concentration of geranyl acetone, a
436 compound related to this attribute. In the present study, the content of hexanal evolved differently in each
437 accession, but 2-isobutylthiazole decreased rapidly. Nevertheless, it is important to highlight that β -

438 ionone and 6-methyl-5-hepten-2-one, compounds derived from carotenoid metabolism as geranyl acetone
439 were also selected as important in the explanation of the aroma evolution of Penjar tomatoes.

440 The evaluation of aroma profiles in tomato is extremely complex. Despite the attempts to generalize the
441 volatile and aroma profiles correlation as a common model for all the tomato varieties, it seems clear that
442 at least in the varieties with long-term conservation such as the Penjar tomatoes, the standard conclusions
443 are not justified. Specific aroma notes may be variety dependent and masking/unmasking relations may
444 reveal the effect of volatiles usually disregarded in the evaluation of tomato aroma.

445

446 **5. Conclusions**

447 The aroma of Penjar tomatoes is mainly characterized by the ‘sharp-floral’ descriptor, although other
448 notes as ‘earthy’ contribute to its typical aroma. The ‘sharp-floral’ aroma note evolves during postharvest
449 (0 to 6 months), increasing during the period 0 to 2 months, when it reaches its maximum. The broad
450 genetic basis of this varietal type results in considerable differences between accessions: two of the 4
451 accessions studied (CDP-1240 and CDP-5468) showed a significantly higher ‘sharp-floral’ intensity, and
452 one accession (CDP-8268) showed a delay in the development of the intensity peak of the ‘sharp-floral’
453 note. These results are very interesting in order to emphasize the added value of this landrace and to
454 determine the better time for its commercialization (2 months).

455 Despite the volatile concentration decrease during the first two months of conservation, there is an
456 increase in ‘sharp-floral’ aroma perception, a result with difficult explanation. The use of iPLS variable
457 selection revealed that 8 of the 24 volatiles detected play a prevalent role, and it seems that the
458 rearrangement of the relative concentrations during the postharvest period and the consequent
459 masking/unmasking processes is the most plausible explanation for the changes in odour intensity during
460 the postharvest of the Penjar tomato.

461

462 **Acknowledgments**

463 This work was supported by grants from the Conselleria de Agricultura, Pesca y Alimentació de la
464 Comunidad Valenciana, the Fundación de la Comunidad Valenciana para la Investigación
465 Agroalimentaria (AGROALIMED) and from the Departament d’Agricultura, Alimentació i Acció Rural
466 (DAR) de la Generalitat de Catalunya.

467

468 **References**

- 469 1. Petro-Turza M (1987) Flavor of tomato and tomato products. *Food Rev Int* 2:309-351
470
- 471 2. Buttery RG (1993) Quantitative and sensory aspects of flavor of tomato and other vegetables and
472 fruits. In: Acree TE, Teranishi R (eds) *Flavor science: sensible principles and techniques*. American
473 Chemical Society, Washington
474
- 475 3. Goff SA, Klee HJ (2006) Plant volatile compounds: sensory cues for health and nutritional value?
476 *Science* 311:815-819
477
- 478 4. Tieman DM, Zeigler M, Schmelz EA, Taylor MG, Bliss P, Kirst M, Klee MJ (2006) Identification of
479 loci affecting flavour volatile emissions in tomato fruits. *J Exp Bot* 57:887-896
480
- 481 5. Zanor MI, Rambla JL, Chaïb J, Steppa A, Medina A, Granell A, Fernie AR, Causse M (2009)
482 Metabolic characterization of loci affecting sensory attributes allows an assessment of the influence of the
483 levels of primary metabolites and volatile organic contents. *J Exp Bot* 60:2139-2154
484
- 485 6. Ortiz-Serrano P, Gil JV (2010) Quantitative comparison of free and bound volatiles of two commercial
486 tomato cultivars (*Solanum lycopersicum* L.) during ripening. *J Agric Food Chem* 58:1106-1114
487
- 488 7. Boukobza F, Taylor AJ (2002) Effect of postharvest treatment on flavour volatiles of tomatoes.
489 *Postharvest Biol Technol* 25:321-331
- 490 8. Vrebalov J, Ruezinsky D, Padmanabhan V, White R, Medrano D, Drake R, Schuch W, Giovannoni J
491 (2002) A MADS-box gene necessary for fruit ripening at the tomato ripening-inhibitor (*rin*) locus.
492 *Science* 296:343-346
- 493 9. Giovannoni JJ, Tanksley SD, Vrebalov J, Noensie E (2004) NOR gene for use in manipulation of fruit
494 quality and ethylene response. US Patent No 5,234,834 issued 13 July 2004
- 495 10. McGlasson WB, Last JH, Shaw KJ, Meldrum SK (1987) Influence of the non-ripening mutants *rin*
496 and *nor* on the aroma of tomato fruit. *HortScience* 22:632-634
- 497 11. Baldwin EA, Scott JW, Shewmaker CK, Schuch W (2000) Flavor trivia and tomato aroma:
498 biochemistry and possible mechanisms for control of important aroma components. *HortScience*,
499 35:1013-1022
- 500 12. Kovács K, Rupert CF, Tikunov Y, Graham N, Bradley G, Seymour GB, Bovy AG, Grierson D (2009)
501 Effect of pleiotropic ripening mutations on flavour volatile biosynthesis. *Phytochemistry* 70:1003-1008
- 502 13. Gao HY, Zhu BZ, Zhu HL, Zhang YL, Xie YH, Li YC, Luo YB (2007) Effect of suppression of
503 ethylene biosynthesis on flavour products in tomato fruits. *Russ J Plant Physiol* 54:80-88
- 504 14. Lewinsohn E, Sitrit Y, Bar E, Azulay Y, Meir A, Zamir D, Tadmor Y (2005) Carotenoid
505 pigmentation affects the volatile composition of tomato and watermelon fruits, as revealed by
506 comparative genetic analyses. *J Agric Food Chem* 53:3142-3148
507
- 508 15. Kopeliovitch E, Mizrahi Y, Rabinowitch D, Kedar N (1980) Physiology of the mutant *alcobaca*.
509 *Physiol Plant* 48:307-311
510
- 511 16. Casals J, Pacual L, Cañizares J, Cebolla J, Casañas F, Nuez F (2011) Genetic basis of long shelf life
512 and variability into Penjar tomato. *Genet Resour Crop Evol*. Doi: 10.1007/s10722-011-9677-6
513
- 514 17. Kuzyomenskii AV (2007) Effect of cumulative polymery of tomato keeping life genes. *Cytology and*
515 *Genetics* 41:268-275
516
- 517 18. Paran I, van der Knaap E (2007) Genetic and molecular regulation of fruit and plant domestication
518 traits in tomato and pepper. *J Exp Bot* 58:3841-3852

- 519
520 19. Moretti CL, Baldwin EA, Sargent SA, Huber DJ (2002) Internal bruising alters aroma volatile profiles
521 in tomato fruit tissues. *HortScience* 37:378-382
522
- 523 20. Buttery RG, Teranishi R, Ling LC (1987) Fresh tomato aroma volatiles: a qualitative study. *J Agric*
524 *Food Chem* 35:540-544
525
- 526 21. Romero del Castillo R, Valero J, Casañas F, Costell E (2008) Training, validation and maintenance of
527 a panel to evaluate the texture of dry beans (*Phaseolus vulgaris* L.). *J Sens Stud* 23:303-319
- 528 22. Beltran J, Serrano E, López FJ, A. Peruga, Valcárcel M, Roselló S (2006) Comparison of two
529 quantitative GC-MS methods for analysis of tomato aroma based on purge-and-trap and on solid-phase
530 microextraction. *Anal Bioanal Chem* 385:1255-1264
- 531 23. Martens H, Naes T (1989) *Multivariate Calibration*. John Wiley and Sons, New York
532
- 533 24. Wise BM, Gallagher NB, Bro R, Shaver JM, Windig W, Koch RS (2006) *Chemometrics tutorial for*
534 *PLS_Toolbox and Solo*. Eigenvector Research Inc, Wenatchee
535
- 536 25. Hongsoongnern P, Chambers E (2008) A lexicon for texture and flavor characteristics of fresh and
537 processed tomatoes. *J Sens Stud* 23:583-599
- 538 26. Norgaard L; Saudland A, Wagner J, Nielsen JP, Munck L, Engelsen SB (2000) Interval partial least-
539 squares regression (iPLS): A comparative chemometric study with an example from near-infrared
540 spectroscopy. *Appl Spectrosc* 54: 413-419
- 541 27. Javanmardi J, Kubota C (2006) Variation of lycopene, antioxidant activity, total soluble solids and
542 weight loss of tomato during postharvest storage. *Postharvest Biol Technol* 41:151-155
543
- 544 28. Kader AA (1986) Effects of postharvest handling procedures on tomato quality. *Acta Hort* 190:209-
545 222
546
- 547 29. Maul F, Sargent SA, Sims CA, Baldwin EA, Balaban MO, Huber DJ (2000) Tomato flavor and aroma
548 quality as affected by storage temperature. *J Food Sci* 65:1228-1237
549
- 550 30. Krumbein A, Auerswald H (1998) Characterization of aroma volatiles in tomatoes by sensory
551 analyses. *Nahrung* 6:S395-S399
552
- 553 31. Tandon KS, Baldwin EA, Shewfelt RL (2000) Aroma perception of individual volatile compounds in
554 fresh tomatoes (*Lycopersicon esculentum*, Mill.) as affected by the medium of evaluation. *Postharvest*
555 *Biol Technol* 20:261-268
- 556 32. Cebolla-Cornejo J, Roselló S, Valcárcel M, Serrano E, Beltran J, Nuez F (2011) Evaluation of
557 genotype and environment effects on taste and aroma flavour components of Spanish fresh tomato
558 varieties. *J Agric Food Chem* (accepted, in production process)
- 559 33. Carbonell-Barrachina AA, Agustí A, Ruiz JJ (2006) Analysis of flavor volatile compounds by
560 dynamic headspace in traditional and hybrid cultivars of Spanish tomatoes. *Eur Food Res Technol*
561 222:536-542
562
- 563 34. Alonso A, Vázquez-Araújo L, García-Martínez S, Ruiz JJ, Carbonell Barrachina AA (2009) Volatile
564 compounds of traditional and virus-resistant breeding lines of *Muchamiel* tomatoes. *Eur Food Res*
565 *Technol* 230:315-323
566
- 567 35. Liggett E, Drake MA, Delwiche JF (2008) Impact of flavor attributes on consumer liking of Swiss
568 cheese. *J.Dairy Sci* 91:466-476
569

570 36. Ortiz-Serrano P, Gil JV (2007) Quantitation of free and glycosidically bound volatiles in and effect of
571 glycosidase addition on three tomato varieties (*Solanum lycopersicum* L.). J Agric Food Chem 55:9170-
572 9176
573
574 37. Xu Y, Barringer S (2010) Comparison of tomatillo and tomato volatile compounds in the headspace
575 by selected ion flow tube mass spectrometry (SIFT-MS). J Food Sci 75:C268-C273
576
577 38. Berna AZ, Lammertyn J, Saevels S, Di Natale C, Nicolai BM (2004) Electronic nose systems to study
578 shelf life and cultivar effect on tomato aroma profile. Sensors and Actuators B, Chemical 97: 324-333
579
580 39. Baldwin EA, Scott JW, Einstein MA, Malundo TMM, Carr BT, Shewfelt RL, Tandon KS (1998)
581 Relationship between sensory and instrumental analysis for tomato flavor. J Am Soc Hortic Sci 12:906-
582 915
583
584 40. Krumbein A, Peters P, Brückner B (2004) Flavour compounds and a quantitative descriptive analysis
585 of tomatoes (*Lycopersicon esculentum* Mill.) of different cultivars in short-term storage. Postharvest Biol
586 Technol 32:15-28
587
588
589
590
591
592
593

594
595

596 **Table 1** Agronomic and morphologic characteristics of the Penjar accessions assayed (mean \pm standard
597 deviation).
598

Accession	Yield (kg plant ⁻¹) ^A	Fruit weight (g) ^B	Soluble solids ($^{\circ}$ Brix) ^B	Fruit colour	Fruit shape	Fruit blossom end shape	Other traits
CDP-1245	2.31 \pm 0.33	61.7 \pm 8.2	4.8 \pm 0.8	Yellow	Flattened	Flat	Potato-leaf
CDP-1240	2.07 \pm 0.66	115.8 \pm 31.8	4.9 \pm 1.0	Orange-red	Heart-shaped	Pointed	High sensibility to fruit cracking
CDP-8268	3.06 \pm 0.86	59.2 \pm 17.4	4.7 \pm 0.4	Orange-red	Heart-shaped	Pointed	Multiparous inflorescence
CDP-5468	1.71 \pm 0.11	31.4 \pm 4.1	6.6 \pm 0.7	Pink	Heart-shaped	Pointed	Multiparous inflorescence

599 ^AMean from 16 plants.

600 ^BFruit traits were evaluated on a random sample of 20 fruits from different plants.

601
602
603
604
605

Table 2 Mean values for postharvest traits. In the same column, different letters indicate significant differences (Student Newman Keuls, at $p \leq 0.05$)

Accession	Shelf life (%) ^A	Loss of weight 2 months (%) ^B	Loss of weight 4 months (%) ^B	Loss of weight 6 months (%) ^B
CDP-1245	59.1 a	12.1 a	19.2 a	27.9 a
CDP-8268	42.8 ab	10.4 ab	16.6 ab	23.9 b
CDP-1240	42.4 ab	9.0 b	14.8 b	21.1 b
CDP-5468	31.2 b	9.8 b	15.9 b	24.0 b

606
607
608
609

^A % commercial fruits at 6 months postharvest

^B % of weight loss with respect to initial weight at harvest

610 **Table 3** Mean concentration (mg kg⁻¹) of main volatiles related to tomato aroma at different postharvest storage periods
 611

Months:	CD-P1245				CDP-1240				CDP-8268				CDP-5468			
	0	2	4	6	0	2	4	6	0	2	4	6	0	2	4	6
2-Phenylethanol	0.7950	0.4337	0.1975	0.2573	0.9388	0.3878	0.1882	0.3787	0.7580	0.3760	0.3859	0.2282	0.3505	0.3563	0.4130	0.3712
<i>trans</i> -2-Hexenal	0.0120	0.1136	0.0260	0.0743	0.6158	0.0823	0.0209	0.0033	0.0442	0.0099	0.0324	0.0283	0.9818	0.3072	0.0103	0.0103
2-Isobutylthiazole	0.0154	0.0208	0.0038	0.0059	0.4603	0.1160	0.0279	0.0012	0.0380	0.0004	0.0007	0.0008	0.2904	0.1153	0.0012	0.0012
6-Methyl-5-hepten-2-one	0.0471	0.0686	0.0328	0.0534	0.2911	0.0724	0.0502	0.0356	0.1800	0.0432	0.0472	0.0474	0.2265	0.1045	0.0471	0.0471
2+3-Methylbutanol	n.d.	0.0145	n.d.	n.d.	0.2537	0.0785	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	0.0270	n.d.	n.d.
Hexanal	0.0424	0.2790	0.0718	0.1640	0.2383	0.2007	0.1150	0.0553	0.0409	0.0509	0.1130	0.0660	0.5090	0.2867	0.0569	0.0569
1-Hexanol	0.0141	0.0350	0.0159	0.0841	0.1658	0.0482	0.0678	0.0145	0.0135	0.0531	0.0281	0.0838	0.2091	0.1611	0.0563	0.0563
<i>cis</i> -3-Hexenol	0.0051	0.0351	0.0057	0.0312	0.1580	0.0469	0.0121	0.0048	0.0087	0.0043	0.0159	0.0105	0.5440	0.0670	n.d.	n.d.
<i>trans</i> -2-heptenal	0.0594	0.0701	0.0122	0.0389	0.1382	0.0700	0.0695	0.0125	0.0473	0.0360	0.0130	0.0120	0.0575	0.0517	0.0671	0.0671
R-Limonene	0.0216	0.0371	0.0081	0.0148	0.1079	0.0330	0.0087	0.0142	0.0158	0.0343	0.0086	0.0128	0.0119	0.0115	0.0352	0.0352
Nonanal	0.0283	0.0255	0.0245	0.0307	0.0641	0.0283	0.0253	0.0250	0.0246	0.0250	0.0244	0.0223	0.0252	0.0316	0.0279	0.0279
Eugenol	0.0276	0.0135	0.0030	0.0151	0.0604	0.0132	0.0146	0.0118	0.0497	0.0094	0.0423	0.0111	0.0338	0.0225	0.0088	0.0088
Geranyl acetone	0.0212	0.0141	0.0010	0.0179	0.0490	0.0171	0.0012	0.0042	0.0403	n.d.	0.0133	0.0109	0.0331	0.0406	0.0036	0.0036
Methyl salicylate	0.0013	0.0247	0.0091	0.0186	0.0486	0.0178	0.0356	0.0110	0.0647	0.0016	0.0330	0.0098	0.0312	0.0273	0.0131	0.0131
Linalool	0.0176	0.0081	0.0047	0.0126	0.0337	0.0084	0.0037	0.0020	0.0395	0.0033	0.0073	0.0034	0.0134	0.0066	0.0041	0.0041
Guaiacol	0.0274	0.0108	0.0026	0.0115	0.0317	0.0099	0.0108	0.0063	0.0642	0.0050	0.0173	0.0083	0.0888	0.0198	0.0049	0.0049
Benzaldehyde	0.0151	0.0129	0.0123	0.0197	0.0293	0.0196	0.0132	0.0122	0.0251	0.0112	0.0125	0.0098	0.0189	0.0224	0.0126	0.0126
α -Terpineol	0.0126	0.0064	0.0037	0.0105	0.0267	0.0056	0.0027	0.0013	0.0313	0.0026	0.0051	0.0028	0.0011	0.0053	0.0031	0.0031
β -Cyclocitral	0.0069	0.0029	0.0020	0.0031	0.0120	0.0041	0.0027	0.0015	0.0087	0.0012	0.0022	0.0011	0.0043	0.0027	0.0015	0.0015
β -Ionone	0.0086	0.0025	0.0016	0.0025	0.0101	0.0031	0.0020	0.0011	0.0060	0.0009	0.0017	0.0011	0.0042	0.0026	0.0012	0.0012
<i>trans</i> -2-Octenal	0.0037	0.0038	0.0025	0.0039	0.0073	0.0041	0.0044	0.0025	0.0062	0.0029	0.0033	0.0022	0.0035	0.0050	0.0035	0.0035
α -Pinene	0.0077	0.0087	0.0077	0.0061	0.0065	0.0091	0.0090	0.0085	0.0085	0.0065	0.0077	0.0060	0.0062	0.0063	0.0080	0.0080
Camphor	0.0019	0.0011	0.0012	0.0018	0.0035	0.0018	0.0013	0.0010	0.0036	0.0008	0.0011	0.0008	0.0019	0.0018	0.0011	0.0011
Phenylacetaldehyde	n.d.	n.d.	n.d.	0.0011	0.0013	0.0029	0.0005	n.d.	n.d.	n.d.	0.0009	n.d.	0.0005	0.0005	n.d.	n.d.
Total	1.192	1.2425	0.4497	0.879	3.7521	1.2808	0.6873	0.6085	1.5188	0.6785	0.8169	0.5794	3.4468	1.6833	0.7805	0.7387

612 n.d.: not detected

613 **Table 4** Loadings of the volatiles included in the PLS model optimized with reverse iPLS variable
 614 selection considering the first two latent variables
 615

Volatile	Loading on latent variable 1	Loading on latent variable 2
α -Terpineol	0.255	-0.582
<i>trans</i> -2-Hexenal	0.426	-0.046
6-Metyl-5-hepten-2-one	0.413	-0.276
<i>trans</i> -2-Octenal	0.413	-0.243
α -Pinene	-0.061	0.359
β -Ionone	0.366	-0.473
2+3-Methylbutanol	0.379	0.239
Phenylacetaldehyde	0.361	0.338

616

617 **Figure captions**

618

619 **Fig. 1** Evolution of the intensity of the ‘sharp-floral’ odour descriptor during postharvest of four Penjar
620 accessions. Inferior abscise legend indicates mean intensity for each postharvest period (different letters
621 indicate significant differences, Student Newman Keuls at $p < 0.05$). Inside the figure, different letters
622 indicate significant differences between accessions within each postharvest time (same statistical
623 procedure)

624

625 **Fig. 2** PLS model optimized with reverse iPLS variable selection relating volatile concentration and
626 sensory evaluation. First latent positively correlated with similar loadings with volatiles *trans*-2-hexenal,
627 6-metyl-5-hepten-2-one, *trans*-2-octenal, 2+3-methylbutanol, phenylacetaldehyde and β -ionone, and with
628 a lower loading with α -terpineol and negatively correlated with α -pinene. Second latent variable
629 positively correlated with volatiles α -pinene, 2+3-methylbutanol and phenylacetaldehyde, and negatively
630 with volatiles 6-metyl-5-hepten-2-one, *trans*-2-octenal and β -ionone. Postharvest storage: ▼ 0 months, *2
631 months, ■ 4 months, + 6 months

632

633

634

635