

Document downloaded from:

<http://hdl.handle.net/10251/101673>

This paper must be cited as:



The final publication is available at

<http://doi.org/10.1039/C4PP00231H>

Copyright The Royal Society of Chemistry

Additional Information

**Photophysics and Photochemistry of the β -Lapachone Derived
Diphenyldihydrodioxin: Generation and Characterization of Its Cation
Radical**

José Carlos Netto-Ferreira^{1,2*}, Virginie Lhiaubet-Vallet², Bauer Bernardes¹, Aurelio
Baird Buarque Ferreira¹ and Miguel Ángel Miranda²

¹ Departamento de Química – Universidade Federal Rural do Rio de Janeiro, Antiga Rio
São Paulo, km 47, Seropédica 23970-000 RJ, Brazil.

² Instituto de Tecnología Química UPV-CSIC, Universitat Politècnica de València,
Avda. de los Naranjos s/n, 46022 Valencia, Spain.

Electronic Supplementary Information

*Corresponding author: jcnetto@ufrj.br

Table of Contents

	page
Synthesis of 2,3-diphenyl-1,4-dioxene (2)	2
Synthesis of the diphenyldihydrodioxin 3	2
Figure 1S. Infrared absorption spectrum for 3 (in KBr).....	4
Figure 2S. Mass spectrum for 3	5
Figure 3S. ¹ H Nuclear Magnetic Resonance for 3 , in CDCl ₃	5
Figure 4S. ¹³ C Nuclear Magnetic Resonance for 3 , in CDCl ₃	6
Figure 5S. UV-visible absorption spectra for 3 , in acetonitrile: (—) 1.2x10 ⁻³ mol/L; (- -) 3.1x10 ⁻⁵ mol/L.....	6
Figure 6S. Normalized absorption (—), excitation (—) and fluorescence emission (- -) spectra for 3 , in acetonitrile.....	7
Figure 7S. Normalized fluorescence emission spectra for 3 in different solvents.....	7
Figure 8S. Chromatogram (GC) for a solution of β-lapachone (1), in ACN.....	8
Figure 9S. Chromatogram (GC) for a solution of 2,3-diphenyldioxene (2), in ACN.....	8
Figure 10S. Chromatogram (GC) for the irradiation of a solution of 3 under N ₂ , in ACN.....	9
Figure 11S. Chromatogram (GC) for the irradiation of an oxygenated solution of 3 , in ACN, showing the formation of a new product with retention time at 12.18 min.....	9
Figure 12S. Absorption spectra for the thermal generation of the diphenyldihydrodioxin cation radical 4 upon successive additions of μL aliquots (from 0 to 600 μL) of a 1.0 mmol L ⁻¹ solution of BAHA to a 1.0 mmol L ⁻¹ solution of diphenyldihydrodioxin 3 , in dichloromethane.....	10
Figure 13S. Superposition of the absorption spectra for thermal and photochemically generated 4	11
Figure 14S. Stern-Volmer plot for the quenching of benzophenone triplet by diphenyldihydrodioxin 3 , in acetonitrile.....	11
References	12

Synthesis of 2,3-diphenyl-1,4-dioxene (2)

This olefin was synthesized following a procedure from the literature.¹ Briefly, 10.6 g (0.05 mol) of benzoin and 0.2 g of *para*-toluenesulfonic acid were heated under reflux with 125 ml of ethylene glycol for 20 h in a dry atmosphere using a calcium chloride tube. The solution was cooled, filtered and washed with an ethylene glycol/water mixture and then dried at low pressure. Yellowish crystals were obtained and TLC analysis showed only one spot, distinct from the substrate. The yield of the crude product was 28%, m.p. = 94-5°C.

IR (KBr, cm⁻¹): 3078, 3052 (aromatic C-H), 2981, 2925, 2878 (C-H), 1634 (C=C), 1598 (aromatic C=C), 1287, 1258 (C-O-C=), 759 e 693 (monosubstituted aromatic ring).
RMN ¹H (CDCl₃) δ (ppm): 7.33 (m, 10H), 4.4 (s, 4H).

Synthesis of the diphenyldihydrodioxin 3

The synthesis of **3** was already described in the literature.² Briefly: β-lapachone (1 mmol) was dissolved in benzene (20 mL) and 2.5 mmol of 2,3-diphenyl-1,4-dioxene was added. After deoxygenating with N₂, the tubes were irradiated with a 500 W Hg lamp. This mixture was filtered under reduced pressure, the colorless solid was collected and recrystallized from ethanol; m.p.: 209–211 °C, yield 69%. Spectra (IR, MS, ¹H NMR and ¹³C NMR) for this compound are shown in Supporting Information (Figures 1S-4S).

UV (acetonitrile, λ_{max} nm (ε, Lmol⁻¹cm⁻¹): 212 (3.9x10⁴), 245.5 (3.08x10⁴), and a broad band from 270 to 360 nm, with maximum at 317 (5.8x10³) (Figure 5S)

IR (KBr) (cm^{-1}): 3065.4, 2972.7, 2935.9, 1646.2, 1586.0, 1495.4, 1450.1, 1413.3, 1389.4, 1326.2, 1264.9, 1240.4, 1180.6, 1160.1, 1105.0, 1068.7, 1042.0, 1018.4, 953.3, 914.2, 854.2, 765.1, 725.9.

MS m/z (abundance): 480 (<1%), 238 (11%), 214 (1%), 199 (1%), 181 (1%), 159 (1%), 130(1%), 105 (100%), 77 (17%), 51 (2%).

HRMS: m/z 480.2020 (theoretical 480.2036)

^1H NMR (CDCl_3) δ (ppm): 8.16 (1H, m); 8.12 (1H, m); 7.77–7.68 (4H, m); 7.44 (1H, dt, $J = 7.02$ and 1.36 Hz); 7.32 (1H, dt, $J = 6.20$ and 1.36 Hz); 7.24–7.18 (6H, m); 4.34–4.14 (2H, m); 3.96–3.89 (2H, m); 3.02–2.76 (2H, m), 1.87 (2H, $J = 6.48$ and 1.62 Hz); 1.42 (3H, s); 1.38 (3H, s).

^{13}C NMR (CDCl_3) δ (ppm): 17.39; 26.63; 26.82; 32.09; 61.41; 61.78; 74.03; 94.43; 95.10; 106.6; 119.75; 121.54; 123.27; 123.99; 125.57; 127.27; 127.67; 128.52; 134.46; 137.46; 137.79; 144.10.




Figure 1S. Infrared absorption spectrum for **3** (in KBr).




Figure 2S. Mass spectrum for **3**.




Figure 3S. ^1H Nuclear Magnetic Resonance for **3**, in CDCl_3




Figure 4S. ^{13}C Nuclear Magnetic Resonance for **3**, in CDCl_3 .




Figure 5S. UV-visible absorption spectra for **3**, in acetonitrile: (—) 1.2×10^{-3} mol/L; (—) 3.1×10^{-5} mol/L.




Figure 6S. Normalized absorption (—), excitation (—) and fluorescence emission (—) spectra for **3**, in acetonitrile.




Figure 7S. Normalized fluorescence emission spectra for **3** in different solvents.




Figure 8S. Chromatogram (GC) for a solution of β -lapachone (**1**), in ACN.




Figure 9S. Chromatogram (GC) for a solution of 2,3-diphenyldioxene (**2**), in ACN.




Figure 10S. Chromatogram (GC) for the irradiation of a solution of **3** under N_2 , in ACN.




Figure 11S. Chromatogram (GC) for the irradiation of an oxygenated solution of **3**, in ACN, showing the formation of a new product with retention time at 12.18 min.




Figure 12S. Absorption spectra for the thermal generation of the diphenylhydrodioxin cation radical **4** upon successive additions of μL aliquots (from 0 to 600 μL) of a 1.0 mmol L^{-1} solution of BAHA to a 1.0 mmol L^{-1} solution of diphenylhydrodioxin **3**, in dichloromethane.




Figure 13S. Superposition of the absorption spectra for thermal and photochemically generated **4**.




Figure 14S. Stern-Volmer plot for the quenching of benzophenone triplet by diphenylhydrodioxin **3**, in acetonitrile.

References

1. R. K. Summerbell, D. R. Berger, *J. Am. Chem. Soc.* 1959, **81**, 633.
2. B. O. Bernardes, A. B. B. Ferreira, J. L. Wardell, S. M. S. V. Wardell, J. C. Netto-Ferreira, E. R. T. Tiekink, *Acta Cryst.* 2013, **E69**, o1487.