

CONTENTS

| | |
|--|-----------|
| CHAPTER 1. GENERAL INTRODUCTION | 1 |
| 1. Global aquaculture sector | 2 |
| 2. Feed and Feeding | 5 |
| 2.1. Requirements..... | 5 |
| 2.1.1. Protein requirements | 6 |
| 2.1.2. Lipid requirements..... | 7 |
| 2.1.3. Carbohydrates requirements | 8 |
| 2.1.4. Vitamins and minerals | 12 |
| 2.2. Effect of feed on rainbow trout histology..... | 13 |
| 2.3. Effect of feed on stress | 14 |
| 2.4. Effect of feed on rainbow trout quality | 15 |
| 2.4.1. Effect of feed on nutritional parameters | 16 |
| 2.4.2. Effect of feed on rainbow trout colour..... | 17 |
| 2.4.3. Effect of feed on rainbow trout fillet texture..... | 18 |
| 2.4.4. Effect of feed on rainbow trout fillet lipid oxidation..... | 18 |
| 2.4.5. Effect of feed on rainbow trout sensorial properties | 19 |
| References..... | 20 |
| CHAPTER 2. OBJECTIVES..... | 44 |
| CHAPTER 3. RESUM OF THE EXPERIMENTS | 47 |
| CHAPTER 4. ENHANCEMENT OF QUALITY OF RAINBOW TROUT (<i>Oncorhynchus mykiss</i>) FLESH INCORPORATING BARLEY ON DIET WITHOUT NEGATIVE EFFECT ON REARING PARAMETERS..... | 50 |

| | |
|---|----|
| 1. INTRODUCTION | 52 |
| 2. MATERIAL AND METHODS | 53 |
| 2.1. <i>Production system</i> | 53 |
| 2.2. <i>Fish and experimental design</i> | 54 |
| 2.3. <i>Diets and feeding</i> | 55 |
| 2.4. <i>Apparent digestibility coefficients</i> | 57 |
| 2.5. <i>Biochemical parameters in blood plasma</i> | 57 |
| 2.6. <i>Quality markers of fish meat</i> | 59 |
| 2.6.1. <i>Proximate composition analysis</i> | 59 |
| 2.6.2. <i>Water activity (a_w)</i> | 60 |
| 2.6.3. <i>Colour</i> | 60 |
| 2.6.4. <i>Texture analysis</i> | 60 |
| 2.6.5. <i>Thiobarbituric Acid Reactive Substances (TBARS)</i> | 61 |
| 2.7. <i>Sensory analysis</i> | 61 |
| 2.8. <i>Statistical analyses</i> | 62 |
| 2.9. <i>Ethical statement</i> | 63 |
| 3. RESULTS | 64 |
| 3.1. <i>Rearing parameters: growth and biometric analysis</i> | 64 |
| 3.2. <i>Apparent digestibility coefficients (ADC)</i> | 64 |
| 3.3. <i>Biochemical parameters</i> | 65 |
| 3.4. <i>Quality markers of fish meat</i> | 67 |
| 3.4.1. <i>Proximate composition</i> | 67 |
| 3.4.2. <i>Water activity (a_w)</i> | 67 |
| 3.4.3. <i>Colour</i> | 68 |
| 3.4.4. <i>Texture</i> | 69 |

| | |
|--|-----------|
| 3.4.5. Thiobarbituric Acid Reactive Substances (TBARS)..... | 70 |
| 3.5. Sensory analysis..... | 71 |
| 4. DISCUSSION | 74 |
| 5. CONCLUSIONS..... | 79 |
| REFERENCES | 80 |
| | |
| CHAPTER 5. EFFECTS OF DIETARY BARLEY ON RAINBOW TROUT EXPOSED TO AN ACUTE STRESS CHALLENGE..... | 91 |
| 1. INTRODUCTION | 93 |
| 2. MATERIAL AND METHODS | 95 |
| 2.1. Diets..... | 95 |
| 2.2. Production system..... | 96 |
| 2.3. Stress challenge: Acute stress (hypoxia)..... | 96 |
| 2.3.1. Sampling | 97 |
| 2.3.2. Cortisol analysis | 97 |
| 2.3.3. Glucose and lactate..... | 97 |
| 2.3.4. MDA..... | 98 |
| 2.4. Statistical analyses..... | 98 |
| 2.5. Ethical statement..... | 99 |
| 3.1. Cortisol | 99 |
| 3.2. Plasma glucose and lactate concentrations..... | 101 |
| 3.3. MDA..... | 102 |
| 4. DISCUSSION | 103 |
| 5. CONCLUSIONS..... | 107 |
| REFERENCES | 107 |

CHAPTER 6. EFFECTS ON HISTOLOGICAL PARAMETERS OF RAINBOW TROUT FED WITH DIFFERENT BARLEY CONCENTRATIONS BEFORE AND AFTER AN HYPOXIA CHALLENGE 116

1. INTRODUCTION 118

2. MATERIAL AND METHODS 120

 2.1. *Fish and rearing conditions*..... 120

 2.2. *Stress challenge: acute stress (hypoxia)*..... 121

 2.3. *Sampling procedure*..... 121

 2.4. *Statistical Analysis*..... 122

3. RESULTS 123

 3.1. *Effect of barley on liver and gut histology and morphology* 123

 3.2. *Effect of stress on liver and gut histology and morphology*..... 128

 3.3. *Interaction between diets and hypoxia on liver and gut histology and morphology*
..... 129

4. DISCUSSION 132

5. CONCLUSION 135

REFERENCES 136

CHAPTER 7. EFFECTS ON LIPID OXIDATION AND BIOACTIVE PROPERTIES OF RAINBOW TROUT FILLETS FED WITH BARLEY. 149

1. INTRODUCTION 151

2. MATERIAL AND METHODS 152

 2.1. *Experimental design*..... 152

 2.2. *Fatty acid profile (FAME)*..... 152

 2.3. *Alpha-tocopherol content*..... 153

 2.4. *Oxidative parameters*..... 153

| | |
|---|------------|
| 2.4.1 Peroxide value (PV)..... | 153 |
| 2.4.2. Conjugated hydroperoxides (dienes and trienes) | 153 |
| 2.5. Antioxidant markers..... | 154 |
| 2.5.1. Total Flavonoid determination (TFC) | 154 |
| 2.5.2. Extract preparation..... | 154 |
| 2.5.3.Total phenols (TP) | 154 |
| 2.5.4. Phenolic characterization using high-performance liquid chromatography (HPLC)..... | 154 |
| 2.5.5. Determination of the oxygen radical absorbance capacity (ORAC) | 155 |
| 2.6. Statistical analysis | 156 |
| 3. RESULTS AND DISCUSSION | 156 |
| 3.1. Fatty acid profile..... | 156 |
| 3.2. Alpha-tocopherol content..... | 158 |
| 3.3. Oxidative parameters..... | 159 |
| 3.3.1. Peroxide value (PV) and conjugated hydroperoxides (dienes and trienes) ... | 159 |
| 3.4. Antioxidant markers..... | 160 |
| 3.4.1. Total flavonoid content (TFC) and total phenolic content (TP)..... | 160 |
| 3.4.2. Individual phenolic compounds | 162 |
| 3.4.3. Fillets antioxidant activity | 164 |
| 4. CONCLUSION..... | 164 |
| REFERENCES | 165 |
| CHAPTER 8. RED BEET AND BETAINE AS INGREDIENTS IN DIETS OF RAINBOW TROUT (<i>Oncorhynchus mykiss</i>): EFFECTS ON GROWTH PERFORMANCE, NUTRIENT RETENTION AND FLESH QUALITY. | 173 |
| 1. INTRODUCTION | 175 |

| | |
|---|-----|
| 2. MATERIAL AND METHODS | 176 |
| 2.1. Diets | 176 |
| 2.2. Rearing markers..... | 178 |
| 2.2.1. Growth trial and fish sampling | 178 |
| 2.2.2. Apparent digestibility coefficients (ADCs) | 181 |
| 2.3. Proximate composition analysis | 181 |
| 2.4. Quality markers of fish flesh and fish sampling..... | 182 |
| 2.4.1. Water activity (a_w)..... | 182 |
| 2.4.2. Colour | 182 |
| 2.4.3. Texture analysis | 183 |
| 2.4.4. Thiobarbituric Acid Reactive Substances (TBARS)..... | 184 |
| 2.4.5. Sensory analysis..... | 184 |
| 2.5. Statistical analysis | 185 |
| 2.6. Ethical statement..... | 185 |
| 3. RESULTS | 186 |
| 3.1. Diets | 186 |
| 3.2. Rearing markers..... | 186 |
| 3.2.1. Growth performance, biometric parameters, body composition and nutrient retention efficiencies | 186 |
| 3.2.2. Apparent digestibility coefficients (ADC)..... | 190 |
| 3.3. Fish flesh proximate composition..... | 192 |
| 3.4. Fish flesh quality markers..... | 194 |
| 3.4.1. Water activity (a_w)..... | 194 |
| 3.4.2. Colour | 195 |
| 3.4.3. Texture | 196 |

| | |
|---|------------|
| 3.4.4. Thiobarbituric Acid Reactive Substances (TBARS)..... | 197 |
| 3.4.5. Sensory analysis..... | 198 |
| 4. DISCUSSION..... | 199 |
| 5. CONCLUSIONS..... | 202 |
| REFERENCES | 203 |
| | |
| CHAPTER 9. EFFECTS OF DIETARY INCLUSIONS OF RED BEET AND BETAINE ON THE ACUTE STRESS RESPONSE AND MUSCLE LIPID PEROXIDATION IN RAINBOW TROUT | 210 |
| 1. INTRODUCTION | 212 |
| 2. MATERIAL AND METHODS | 213 |
| 2.1. Production system..... | 213 |
| 2.2. Fish, diets and feeding..... | 214 |
| 2.3. Stress trial..... | 215 |
| 2.3.1. Sampling | 215 |
| 2.3.2. Cortisol analysis | 216 |
| 2.3.3. Glucose and lactate..... | 216 |
| 2.3.4. Lipid peroxidation..... | 217 |
| 2.4. Statistical analyses..... | 217 |
| 2.5. Ethical statement..... | 218 |
| 3. RESULTS | 218 |
| 3.1. Cortisol | 218 |
| 3.2. Plasma glucose and lactate | 221 |
| 3.3. Lipid peroxidation..... | 223 |
| 4. DISCUSSION | 224 |
| 5. CONCLUSIONS..... | 227 |

| | |
|------------------|-----|
| REFERENCES | 227 |
|------------------|-----|

CHAPTER 10. RED BEET AND BETAINE HAVE A POSITIVE EFFECT ON HISTOLOGICAL PARAMETERS OF RAINBOW TROUT BEFORE AND AFTER AN ACUTE HYPOXIA CHALLENGE.....237

| | |
|---|-----|
| 1. INTRODUCTION | 239 |
| 2. MATERIAL AND METHODS | 241 |
| 2.1. <i>Experimental design</i> | 241 |
| 2.2. <i>Diets</i> | 242 |
| 2.3. <i>Rearing conditions and feeding</i> | 242 |
| 2.4. <i>Acute hypoxia</i> | 243 |
| 2.5. <i>Sampling procedure</i> | 243 |
| 2.6. <i>Statistical analysis</i> | 244 |
| 3. RESULTS | 244 |
| 3.1. <i>Effect of red beet and betaine on liver and gut histology and morphology</i> | 244 |
| 3.2. <i>Effect of the stress on liver and gut histology and morphology</i> | 246 |
| 3.3. <i>Interaction between diets and stress on liver and gut histology and morphology</i> | 247 |
| 4. DISCUSSION | 250 |
| 5. CONCLUSION | 253 |
| REFERENCES | 254 |

CHAPTER 11. REDUCTION OF LIPID OXIDATION AND ENHANCEMENT OF BIOACTIVITY THROUGH THE INCLUSION OF RED BEET AND BETAINE ON RAINBOW TROUT DIETS.270

| | |
|-------------------------------|-----|
| 1. INTRODUCTION | 272 |
| 2. MATERIAL AND METHODS | 274 |

| | |
|--|-----|
| 2.1. Experimental design..... | 274 |
| 2.2. Fatty acid profile (FAME) | 274 |
| 2.3. Alpha-tocopherol content..... | 275 |
| 2.4. Oxidative parameters..... | 275 |
| 2.4.1. Peroxide value (PV)..... | 275 |
| 2.4.2. Conjugated hydroperoxides (dienes and trienes) | 275 |
| 2.5. Antioxidant markers..... | 276 |
| 2.5.1. Extract preparation..... | 276 |
| 2.5.2. Phenolic characterization using HPLC | 276 |
| 2.5.3. Total Flavonoid determination (TFC) | 277 |
| 2.5.4. Total phenols (TP) | 277 |
| 2.5.5. Determination of the oxygen radical absorbance capacity (ORAC)..... | 277 |
| 2.5.6. Trolox Equivalent Antioxidant Capacity (TEAC) and DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity..... | 277 |
| 2.5.7. Relative Antioxidant Capacity Index (RACI)..... | 278 |
| 2.6. Statistical analysis | 278 |
| 3. RESULTS AND DISCUSSION | 278 |
| 3.1. Fatty acid profile..... | 278 |
| 3.2. α -tocopherol content | 282 |
| 3.3. Oxidative parameters: peroxide value (PV) and conjugated hydroperoxides (dienes and trienes)..... | 282 |
| 3.4. Antioxidant activity..... | 282 |
| 3.4.1. Total flavonoid content (TFC) and total phenolic content (TP)..... | 282 |
| 3.4.2. Individual phenolic compounds | 284 |
| 3.4.3. Antioxidant activity of fish fillets | 286 |

| | |
|--|------------|
| 4. CONCLUSION..... | 287 |
| REFERENCES | 288 |
| CHAPTER 12. GENERAL DISCUSSION..... | 298 |
| 1. Growth parameters..... | 299 |
| 2. Stress..... | 302 |
| 3. Histology..... | 304 |
| 4. Flesh quality..... | 309 |
| 5. Sensory analysis..... | 312 |
| 6. Economic analysis | 313 |
| References..... | 316 |
| CHAPTER 13. CONCLUSIONS..... | 328 |

List of Tables

| | |
|---|------------|
| Table 1. Vitamin requirements of rainbow trout (mg/kg dry diet) | 12 |
| Table 2. Mineral requirements of rainbow trout^a | 13 |
| Table 3. Different experiments carried out on the current Doctoral Thesis | 48 |
| Table 4. Formulation and proximate composition of the experimental diets | 56 |
| Table 5. Growth and biometric indexes of rainbow trout fed with different experimental diet for 84 days (n=4)..... | 64 |
| Table 6. Apparent digestibility coefficients (ADC) of protein, fat and carbohydrate in rainbow trout fed with five experimental diets differing on the source of carbohydrate (wheat and barley) (n=3). | 65 |
| Table 7. Proximate composition of rainbow trout meat fed with increasing levels of barley at the end of the experimental period (data are expressed as % of dry matter) (n=4)..... | 67 |
| Table 8. Effect of barley concentration on the CIELAB parameters of fish meat at the end of the experimental growth period (n=4)..... | 69 |
| Table 9. Effect of barley concentration on hardness, cohesiveness, elasticity and gumminess of fish meat fed with diets with increasing concentrations of barley at the end of the experimental growth period (n=4). | 69 |
| Table 10. Areas, maximum (Lmax) and minimum (Lmin) lengths of hepatocytes of fish fed different experimental diets (n=4)..... | 123 |
| Table 11. Areas, maximum (Lmax) and minimum (Lmin) lengths of hepatocytes of fish fed different experimental diets after the stress challenge (n=4)..... | 130 |
| Table 12. Fatty acid profiles (FA) of barley and experimental diets..... | 157 |
| Table 13. Fatty acid profiles (FA) of rainbow trout fillets when fish were fed with increasing barley levels. (data are expressed as % of dry matter) (n=12). | 158 |

| | |
|---|------------|
| Table 14. Effect of barley on the peroxide value (PV) and conjugated hydroperoxides (dienes and trienes) of rainbow trout fillet of fish fed at different barley concentrations (data are expressed as % of dry matter) (n=12). | 160 |
| Table 15. Content in individual phenolic compounds in barley and experimental diets ($\mu\text{g mL}^{-1}$). | 163 |
| Table 16. Effect of barley on antioxidant properties of rainbow trout fillets of fish fed at different barley concentrations (data are expressed as % of dry matter) (n=12). | 164 |
| Table 17. Formulation and proximate composition of the experimental diets. | 177 |
| Table 18. Effect of red beet and total betaine level on growth and nutritive parameters of rainbow trout (values are least-squares means \pm SEM, n=3). ... | 188 |
| Table 19. Effects of red beet and total betaine level on biometric parameters, body composition and nutrient retention of rainbow trout (values are least-squares means \pm SEM, n=3). | 189 |
| Table 20. Apparent digestibility coefficients (ADCs) of protein (ADC_{protein}), lipid (ADC_{lipid}) and carbohydrates (ADC_{CHO}) in rainbow trout fed the experimental diets differing on the source of carbohydrate (wheat and barley) (values are least-squares means \pm SEM, n=3). | 191 |
| Table 21. Proximate composition of rainbow trout flesh fed with increasing red beet and betaine levels at the end of the experimental growth period (data are expressed as % of dry matter) (values are least-squares means \pm SEM, n=3)..... | 193 |
| Table 22. Effect of red beet and betaine on hepatocytes areas, maximum (L_{Max}) and minimum (L_{Min}) lengths (μm) before the stress. (values are least-squares means \pm SEM, n=4). | 245 |

| | |
|---|------------|
| Table 23. Effect of red beet and betaine on hepatocytes areas, maximum (LMax) and minimum (LMin) lengths (μm) after the stress (values are least-squares means \pm SEM, n=4). | 248 |
| Table 24. Red beet and experimental diets fatty acid profiles. | 279 |
| Table 25. Effect of red beet and betaine on the fatty acid profile of rainbow trout filets. Data are shown as least-squares means \pm standard error of the mean (SEM) of triplicate groups (n=9) | 281 |
| Table 26. Effect of red beet and betaine on the antioxidant activity of rainbow trout filets. Data are shown as least-squares means \pm standard error of the mean (SEM) of triplicate groups (n=9) | 287 |
| Table 27. Growth and biometric parameters of rainbow trout fed at different barley and red beet and betaine concentrations. | 300 |
| Table 29. Results of economic parameters at the end of the trial using barley as experimental ingredient (n = 4). | 314 |
| Table 30. Results of economic parameters at the end of the trial using red beet and betaine as experimental ingredients (n = 4). | 315 |

List of Figures

| | |
|--|------------|
| Figure 1. World capture and aquaculture production (FAO 2016. The state of world fisheries and aquaculture) | 2 |
| Figure 2. Share of aquaculture in total European production of aquatic animals (FAO 2016. The state of world fisheries and aquaculture)..... | 4 |
| Figure 3. Overview of factor affecting fish quality parameters (Source: Lie (2001), Flesh quality - the role of nutrition. Aquaculture Research 32, 341-348, with some modifications) | 16 |
| Figure 5. Effect of barley concentration on water activity (aw) of fish at the end of the experimental growth period. | 68 |
| Figure 6. Effect of barley on meat gumminess of fish fed different experimental diets | 70 |
| Figure 7. Effect of barley concentration on lipid oxidation (TBARS) of fish meat at the end of experimental growth period. | 71 |
| Figure 8. Effect of barley concentration on the gill colour of fish from sensory analysis (QIM) at the end of the experimental growth period. | 72 |
| Figure 9. Effect of barley concentration on the colour of fish meat from sensory analysis (QDM) at the end of the experimental growth period. | 73 |
| Figure 10. Effect of barley concentration on the texture of fish meat from sensory analysis (QDM) at the end of the experimental growth period. | 73 |
| Figure 11. Effect of barley on plasma cortisol (A) and fish cortisol (B) content of rainbow trout under normoxia (basal), 30 minutes, 6 and 12 hours after stress. | 100 |

| | |
|---|------------|
| Figure 12. Effect of barley on plasma glucose (A) and lactate (B) content of rainbow trout under normoxia (basal), 30 minutes, 6 and 12 hours after stress. | 102 |
| Figure 13. Effect of barley on MDA values (nM malonaldehyde per mg of protein) of rainbow trout under normoxia (basal), 30 minutes, 6 and 12 hours after stress. | 103 |
| Figure 14. Hepatocytes vacuolation of fish fed different experimental diets. | 124 |
| Figure 15. Incidence of lymphocytic foci appearance in hepatocytes of fish fed different experimental diets. | 125 |
| Figure 16. Cell infiltration on the posterior intestine of fish fed different experimental diets. | 126 |
| Figure 17. Mucus cells on the posterior intestine of fish fed different experimental diets. | 127 |
| Figure 18. Gut vacuolation (posterior intestine) of fish fed different experimental diets. | 128 |
| Figure 19. Effect of stress on hepatocytes size (areas and lengths). | 129 |
| Figure 20. Hepatocytes vacuolation of fish fed different experimental diets after the stress challenge. | 130 |
| Figure 21. Incidence of lymphocytic foci appearance in hepatocytes of fish fed different experimental diets after the stress challenge. | 131 |
| Figure 22. Gut vacuolation (posterior intestine) of fish fed different experimental diets after the stress challenge. | 132 |
| Figure 23. Fillet alpha-tocopherol content of fish fed different experimental diets (n=12). | 159 |
| Figure 24. Barley and experimental diets total flavonoid contents (TFC). | 161 |

| | |
|--|------------|
| Figure 25. Barley and experimental diets total phenolic contents (TP)..... | 162 |
| Figure 26. HPLC chromatogram of phenolic compounds profile in barley extracts | 163 |
| Figure 27. Effect of increasing levels of red beet and betaine on fish flesh betaine content..... | 194 |
| Figure 28. Effect of red beet and betaine concentration on water activity (aw) of fish meat at the end of the experimental growth period..... | 195 |
| Figure 29. Effect of red beet and betaine on fish flesh redness (A* values) at the end of the experimental growth period..... | 196 |
| Figure 30. Effect of red beet and betaine on fish flesh elasticity at the end of the experimental growth period..... | 197 |
| Figure 31. Effect of red beet and betaine concentration on lipid oxidation (TBARS) measured as μg malonaldehyde g^{-1} of fish meat at the end of the experimental growth period..... | 198 |
| Figure 32. Effect of red beet and betaine on plasma cortisol (A) and fin cortisol (B) content of rainbow trout before the acute stress challenge (basal), 30 minutes after stress and 6 and 12 hours after stress. | 220 |
| Figure 33. Effect of red beet and betaine on plasma glucose (A) and lactate (B) content of rainbow trout before the acute stress challenge (basal), 30 minutes after stress and 6 and 12 hours after stress. | 222 |
| Figure 34. Effect of red beet and betaine on MDA (nmol per g of tissue) of rainbow trout before the acute stress challenge (basal), 30 minutes after stress and 6 and 12 hours after stress..... | 223 |
| Figure 35. Hepatocytes vacuolization of fish fed different experimental diets. | 245 |

| | |
|---|------------|
| Figure 36. Incidence of lymphocytic foci appearance in hepatocytes of fish fed different experimental diets before the hypoxia challenge. | 246 |
| Figure 37. Effect of the stress on hepatocytes areas, maximum (L_{max}) and minimum (L_{min}) lengths. | 247 |
| Figure 38. Hepatocytes vacuolization of fish fed different experimental diets 30 minutes after the hypoxia challenge. | 249 |
| Figure 39. Incidence of lymphocytic foci appearance in hepatocytes of fish fed different experimental diets after the hypoxia challenge. | 250 |
| Figure 40. Red beet and experimental diets total flavonoid content (TFC). | 283 |
| Figure 41. Red beet and experimental diets total phenolic content (TP). | 284 |
| Figure 42. Fillets total phenolic content (TP) of fish fed with different experimental diets (n=9). | 284 |
| Figure 43. HPLC chromatogram of phenolic compounds in red beet extracts. ... | 286 |
| Figure 44. Summary of the different aspects studied in the current doctoral thesis. | 299 |
| Figure 45. Growth curve in barley and red beet experiments. | 300 |
| Figure 46. Hepatocytes of fish fed with diets containing red beet and betaine before the stress trial. | 305 |
| Figure 47. Liver degeneration of fish fed with diets containing barley before the stress challenge. | 306 |
| Figure 48. Gut cell infiltration of fish fed with diets containing 8% barley before the stress challenge. | 307 |