

Regulation of the nitric oxide synthesis and signaling by post-translational modifications and N-end rule pathway-mediated proteolysis in *Arabidopsis thaliana*

**TABLE OF CONTENTS**

<b>1. <u>INTRODUCTION</u></b>	1
1.1. <u><i>Arabidopsis thaliana</i> as model organism</u>	3
1.2. <u>Biosynthesis and metabolism of nitric oxide</u>	3
1.3. <u>NO as a regulator</u>	7
1.4. <u>NO sensing</u>	10
1.5. <u>NO and stress-triggered responses. The role on plant tolerance to freezing</u>	12
<b>2. <u>OBJECTIVES</u></b>	15
<b>3. <u>MATERIALS AND METHODS</u></b>	19
3.1. <u>Biological material</u>	21
3.1.1. <u>Plant material</u>	21
3.1.1.1. Plant species used	21
3.1.1.2. Plant growth	22
3.1.1.2.1. Growing media for plants	22
3.1.1.2.2. Growing conditions for plants	22
3.1.2. <u>Microbiological material</u>	23
3.1.2.1. Bacterial strains used	23
3.1.2.2. Bacterial growth	23
3.1.2.2.1. Growing media for bacteria	23
3.1.2.2.2. Growing conditions for bacteria	23
3.2. <u>Manipulation of the biological material</u>	23
3.2.1. <u>Plant treatments</u>	23
3.2.1.1. Plant transformation by floral dipping	23
3.2.1.2. Generation of a <i>nir1</i> mutant by CRISPR-Cas9 technology	23
3.2.1.3. NO treatments	26
3.2.1.4. Freezing tolerance assays	26
3.2.1.5. Staining methods	27
3.2.1.5.1. NO detection by fluorescence and confocal microscopy	27
3.2.1.5.2. Cuticle permeability tests	27
3.2.1.5.3. Staining for cell permeability, cell death, superoxide and starch	28
3.2.1.6. Change of nitrogen source for plant growth	28
3.2.1.7. Inhibition of protein synthesis or degradation	28
3.2.2. <u>Bacterial manipulation</u>	28
3.2.2.1. Genetic transformation by heat shock	28

<b>3.3. <u>Nucleic acid methods</u></b>	29
<b>3.3.1. Extraction and purification of nucleic acids</b>	29
3.3.1.1. Extraction of genomic DNA from leaves or seedlings	29
3.3.1.2. Cleaning of PCR products	29
3.3.1.3. Extraction of DNA from agarose gel	29
3.3.1.4. Extraction and isolation of plasmidic DNA from bacteria	29
3.3.1.5. Extraction and isolation of total RNA from seedlings	30
<b>3.3.2. Manipulation of nucleic acids</b>	30
3.3.2.1. Methods for quantification and concentration	30
3.3.2.2. Methods for DNA amplification	30
3.3.2.3. Reverse transcription (RT) of mRNA	30
3.3.2.4. Quantification of transcripts by quantitative PCR (qPCR)	31
3.3.2.5. Transcriptomic analysis by microarrays	31
3.3.2.6. DNA sequencing	32
3.3.2.7. Electrophoretic techniques	32
3.3.2.8. Enzymatic digestion of DNA with restriction enzymes	33
3.3.2.9. Southern blot	33
3.3.2.10. Cloning techniques in bacteria	33
<b>3.4. <u>Protein methods</u></b>	34
<b>3.4.1. Extraction and purification of proteins</b>	34
3.4.1.1. Extraction of total proteins from plant tissues	34
3.4.1.2. Immunopurification of tagged proteins with magnetic beads	34
3.4.1.3. Precipitation of proteins	34
<b>3.4.2. Manipulation of proteins</b>	34
3.4.2.1. Quantification of total proteins	34
3.4.2.2. Protein separation by SDS-PAGE	34
3.4.2.3. Protein detection by Western blot analyses	35
3.4.2.4. LC-MS/MS-based proteomic analyses	35
3.4.2.5. NR and NIR activity assays	35
<b>3.5. <u>Metabolic methods</u></b>	36
<b>3.5.1. LC/MS- and GC/MS-based analyses of the metabolome</b>	36
<b>3.5.2. Quantification of anthocyanins</b>	38
<b>3.5.3. Phytohormone quantification</b>	38
<b>3.6. <u>In silico and statistical methods</u></b>	38
<b>3.6.1. In silico analyses</b>	38
<b>3.6.2. Prediction of PTMs</b>	39
<b>3.6.3. Statistical analyses</b>	39
<b>4. <u>RESULTS</u></b>	41
<b><u>4.1. Regulation of NR-dependent NO biosynthesis. Post-translational modifications of the biosynthetic enzymes</u></b>	43
<b>4.1.1. Abstract</b>	43
<b>4.1.2. Nitrate Reductase is Regulated by Proteasome-mediated Degradation and Nitrate Signaling</b>	43

4.1.3. NR and NiR1 proteins are post-translationally regulated by NO	49
4.2. <u>NO-triggered transcriptomic and metabolomic responses.</u>	55
<u>Role of hormones in NO sensing</u>	
4.2.1. Abstract (1)	55
4.2.2. Overrepresentation of hormone- and oxygen-related regulatory components in the early NO-responsive transcriptome	55
4.2.3. A sensing test based on inhibition of hypocotyl elongation allowed identification of hormone-related transcription factors modulating NO sensitivity	57
4.2.4. Ethylene perception and signaling as well as salicylate and strigolactone biosynthesis are required for NO sensing	60
4.2.5. NO sensing and ABA signalling	64
4.2.6. NO sensing and jasmonate signaling	66
4.2.7. Involvement of brassinosteroids in NO sensing	67
4.2.8. Abstract (2)	70
4.2.9. Metabolomic analyses reflect a transient reprogramming response to exogenous NO	70
4.2.10. Altered levels of amino acids and dipeptides suggests NO treatment increased protein breakdown	72
4.2.11. Acute exposure to NO triggers the accumulation of polyamines and responses to oxidative stress	74
4.2.12. Altered lipidome reflects changes in lipidic structures such as membranes and cuticle in NO-exposed plants	76
4.2.13. Altered purine, pyrimidine and chlorophyll metabolism suggest NO enhanced nucleic acid turnover, chlorophyll degradation and non-programmed cell death	78
4.2.14. Exposure to NO affected photorespiration and central carbon metabolism	80
4.3. <u>Function of the ERFVII RAP2.3 on the regulation of NO homeostasis and signaling</u>	83
4.3.1. Abstract	83
4.3.2. A rheostat-like mechanism based on RAP2.3 degradation controls endogenous NO content	83
4.3.3. RAP2.3 modulates NO sensing in shoots and roots	86
4.3.4. Genome-wide transcriptome analyses revealed RAP2.3 as a general negative regulator of NO-triggered responses	87
4.3.5. The NO-responsive RAP2.3-independent transcriptome includes jasmonic acid and ethylene core signaling gene sets	91
4.3.6. The NO-responsive RAP2.3-dependent transcriptome suggest the existence of NO-sensitive gene-specific hormone signaling pathways	91
4.4. <u>Role of NO in constitutive and cold acclimation induced freezing tolerance</u>	96

4.4.1. Abstract (1)	96
4.4.2. The transcriptome of NO-deficient <i>nia1,2noa1-2</i> mutant plants is enriched in cold-related transcripts	96
4.4.3. Enhanced biosynthesis of ABA, JA and osmoprotective metabolites in NO-deficient plants	98
4.4.4. Increased levels of antioxidant metabolites in <i>nia1,2noa1-2</i> plants	101
4.4.5. NO negatively regulates constitutive freezing tolerance of Arabidopsis	101
4.4.6. Abstract (2)	104
4.4.7. NO functions as a positive regulator of cold acclimation in Arabidopsis	104
4.4.8. NO promotes the cold-induced expression of <i>CBF</i> genes in Arabidopsis	107
4.4.9. NO regulates the sensitivity of Arabidopsis to ABA in response to low temperature	108
4.4.10. NO activates anthocyanin accumulation in Arabidopsis during cold acclimation	109
5. <u>DISCUSSION</u>	113
6. <u>CONCLUSIONS</u>	135
7. <u>BIBLIOGRAPHY</u>	139
A. <u>ANNEXES</u>	i

## FIGURES

Figure 1.1. Scheme of the main biosynthetic and metabolic pathways of NO	5
Figure 1.2. Scheme of the Cys-Arg/N-end rule pathway-mediated degradation of ERFVII transcription factors as sensor mechanism of NO in plants	11
Figure 1.3. Scheme summarizing the CBF-dependent pathway of cold acclimation-induced freezing tolerance	14
Figure 3.1. Generation of <i>nir1</i> mutant plants by CRISPR-Cas9 technology	25
Figure 4.1. Effect of nitrogen source and proteasome-mediated degradation on NR and NiR	44
Figure 4.2. Regulation of NR and NiR by NLP7 nitrate signaling factor	46
Figure 4.3. Complementation of <i>nlp7-1</i> and <i>prt6-1</i> related phenotypes and NO production by <i>NLP7</i> over-expression	47
Figure 4.4. Generation of transgenic plants overexpressing HA-tagged versions of NR1, NR2 and NiR1	48

Figure 4.5. NO accumulation in plants altered in nitrite reduction to ammonium	48
Figure 4.6. NR protein in nitrate signaling and Arg/N-end rule pathway mutants.	48
Figure 4.7. Effect of exogenous NO on NR protein stability	49
Figure 4.8. Post-translational modifications identified in planta for NR1/NIA1 and NIR2/NIA2	50
Figure 4.9. Post-translationally modified residues in aligned NR1/NIA1 and NR2/NIA2 proteins	52
Figure 4.10. Post-translational modifications identified in planta for NiR1	53
Figure 4.11. Location of S-nitrosylated cysteines and nitrated/aminated tyrosines in the 3D model of NiR1	54
Figure 4.12. 3D modelling of Arabidopsis NRs showing the position of nitrated conserved tyrosine residues close to the FAD binding site	54
Figure 4.13. Identification and characterization of the NO-responsive transcriptome	56
Figure 4.14. Screening of TPT transgenic lines conditionally expressing transcription factor encoding genes through a NO-triggered hypocotyl shortening assay in etiolated seedlings	58
Figure 4.15. $\beta$ -estradiol induced transcript accumulation in randomly selected TRANSPLANTA transgenic lines	59
Figure 4.16. Ethylene perception and signaling is required for NO sensing	61
Figure 4.17. Effect of NO treatment on the transcript levels of A, ethylene, B, jasmonate and C, brassinosteroid biosynthetic/or signaling encoding genes	62
Figure 4.18. Involvement of salicylates and strigolactones in NO sensing	63
Figure 4.19. Involvement of ABA signaling in NO sensing	65
Figure. 4.20. Involvement of jasmonate signaling in NO sensing	67
Figure 4.21. Genes up-regulated by either brassinolide (BL) or NO treatments	68
Figure 4.22. Involvement of brassinosteroid signaling in NO sensing	69
Figure 4.23. Metabolomic Changes after Applying an NO Pulse to Arabidopsis Plants	71
Figure 4.24. Superoxide content reduction and post-translational modifications induced by NO	74
Figure 4.25. Enhanced polyamine content in plants exposed to NO	75
Figure 4.26. Altered phospholipid catabolism in plants after exposure to NO	77

Figure 4.27. Alterations in the permeability of lipidic structures	78
Figure 4.28. NO triggers DNA and chlorophyll degradation	79
Figure 4.29. Endogenous NO content and cell death upon exposure to exogenous NO	80
Figure 4.30. Metabolites of glycolysis and TCA cycle in plants exposed to NO	81
Figure 4.31. NO in plants overexpressing MC- and MA-RAP2.3 versions	84
Figure 4.32. NO in N-end rule pathway mutant and wild type plants	85
Figure 4.33. NO in roots of mutant and transgenic plants	86
Figure 4.34. Overexpression of non-degradable MA-RAP2.3 confers hyposensitivity to NO	88
Figure 4.35. Transcriptome analysis of $\beta$ -estradiol inducible RAP2.3 transgenic lines	90
Figure 4.36. Effect of NO and RAP2.3 on the expression of JA biosynthesis and signaling genes	93
Figure 4.37. Effect of NO and RAP2.3 on the expression of ABA biosynthesis and signaling genes	95
Figure 4.38. Levels of cold-inducible transcripts in Col-0 and <i>nia1,2noa1-2</i> plants	97
Figure 4.39. Levels of hormones and osmoprotective metabolites in Col-0 and <i>nia1,2noa1-2</i> plants	99
Figure 4.40. Glycolysis and TCA cycle metabolite ratios between <i>nia1,2noa1-2</i> and Col-0 plants	100
Figure 4.41. Endogenous content of ascorbate-glutathione cycle metabolites in wild type and NO-deficient plants	102
Figure 4.42. Constitutive freezing tolerance of Col-0 and <i>nia1,2noa1-2</i> plants	103
Figure 4.43. NO levels in cotyledons of Col-0 and <i>nia1,2noa1-2</i> plants after cold acclimation	104
Figure 4.44. NO levels in Col-0 and <i>nia1,2noa1-2</i> plants at different times after cold treatment	105
Figure 4.45. Cold acclimation-induced freezing tolerance of Col-0 and <i>nia1,2noa1-2</i> plants	106
Figure 4.46. Effect of different nitrogen sources on NO production	107
Figure 4.47. Effects of cold acclimation on the CBFs and their targets in Col-0 and <i>nia1,2noa1-2</i> plants	108
Figure 4.48. ABA homeostasis and signaling in cold acclimated wild-type and NO-deficient plants	109
Figure 4.49. Effects of cold acclimation on anthocyanin synthesis in Col-0 and <i>nia1,2noa1-2</i> plants	110

Figure 5.1. Scheme showing the direct involvement of ethylene (ET), strigolactones (SL), salicylates (SA), abscisic acid (ABA) and brassinosteroids (BR) in NO-triggered inhibition of hypocotyl elongation.	118
Figure 5.2. Model of NO involvement in the regulation of constitutive freezing tolerance	120
Figure 5.3. Scheme showing the endogenous and environmental factors involved in the positive regulation exerted by NO on cold acclimation-induced freezing tolerance	122
Figure 5.4. Scheme summarizing the regulation that high concentrations of NO exert over different metabolite categories and cell processes	125
Figure 5.5. RAP2.3-dependent and –independent NO-regulated pathways	128
Figure 5.6. Regulation of NRs, NiR1 and NO production.	131
Figure 5.7. Scheme showing processes investigated in this work that regulate NO or are regulated by it	133
Figure A1. Map of pCR8/GW/TOPO vector (Invitrogen)	iv
Figure A2. Map of pALLIGATOR2 vector	v
Figure A3. Map of pMTN2982 (pDONR207) vector	vi
Figure A4. Map of pHEE2E-TRI vector	vii
Figure A5. <i>In silico</i> analyses of <i>NLP7</i> promoter	viii

## TABLES

Table 4.1. TPT lines showing hypo- or hyper-sensitivity to NO on hypocotyl elongation of etiolated seedlings upon conditional expression of TF encoding genes	60
Table 4.2. Statistical comparisons in the metabolomic analyses	72
Table A1. Abbreviations and acronyms	ix
Table A2. Buffers and solutions used in this work	xii
Table A3. Oligonucleotides used in this work	xiii
Table A4. Proteins with the MC- N-terminal amino acid sequence in the Arabidopsis proteome	xvi
Table A5. Genotyping of T1 generation of <i>pU6::gNIR1-1</i>	xx
Table A6. Prediction of S-nitrosylation and nitration sites of potential NO targets	xxi
Table A7. <i>In silico</i> genome-wide analysis of putative <i>RAP2.3</i> -binding motifs and the intersection with NO-regulated genes	xxiii
Table A8. Genes that were up-regulated in non-acclimated <i>nia1,2noa1-2</i> plants and were reported to be related to cold-triggered responses	xxix

**Table A9. Differential transcript levels in TPT-RAP2.3 lines not- treated with  $\beta$ -estradiol** xxxiii

## **EQUATIONS**

**Equation 3.1. Normalized relative expression** 31