

Fixed point sets in digital topology, 1

LAURENCE BOXER^a AND P. CHRISTOPHER STAECKER^b

^a Department of Computer and Information Sciences, Niagara University, Niagara University, NY 14109, USA; and Department of Computer Science and Engineering, SUNY at Buffalo (boxer@niagara.edu)

^b Department of Mathematics, Fairfield University, Fairfield, CT 06823-5195, USA. (cstaecker@fairfield.edu)

Communicated by V. Gregori

ABSTRACT

In this paper, we examine some properties of the fixed point set of a digitally continuous function. The digital setting requires new methods that are not analogous to those of classical topological fixed point theory, and we obtain results that often differ greatly from standard results in classical topology.

We introduce several measures related to fixed points for continuous self-maps on digital images, and study their properties. Perhaps the most important of these is the fixed point spectrum $F(X)$ of a digital image: that is, the set of all numbers that can appear as the number of fixed points for some continuous self-map. We give a complete computation of $F(C_n)$ where C_n is the digital cycle of n points. For other digital images, we show that, if X has at least 4 points, then $F(X)$ always contains the numbers 0, 1, 2, 3, and the cardinality of X . We give several examples, including C_n , in which $F(X)$ does not equal $\{0, 1, \dots, \#X\}$.

We examine how fixed point sets are affected by rigidity, retraction, deformation retraction, and the formation of wedges and Cartesian products. We also study how fixed point sets in digital images can be arranged; e.g., for some digital images the fixed point set is always connected.

2010 MSC: 54H25.

KEYWORDS: *digital image; fixed point; retraction.*

1. INTRODUCTION

Digital images are often used as mathematical models of real-world objects. A digital model of the notion of a continuous function, borrowed from the study of topology, is often useful for the study of digital images. However, a digital image is typically a finite, discrete point set. Thus, it is often necessary to study digital images using methods not directly derived from topology. In this paper, we introduce several such methods to study properties of the fixed point set of a continuous self-map on a digital image.

Many of our results have elementary proofs. Their importance is, in part, due to the following. Digital topology has been successful in showing that digital images resemble the Euclidean objects they model with respect to topological properties such as connectedness, homotopy, covering maps, fundamental groups, retractions, and homology; however, we see in this paper that the fixed point properties of digital images and the Euclidean objects they model can be very different.

Some of the results of this paper were presented in [7].

2. PRELIMINARIES

Let \mathbb{N} denote the set of natural numbers; and \mathbb{Z} , the set of integers. $\#X$ will be used for the number of elements of a set X .

2.1. Adjacencies. A digital image is a pair (X, κ) where $X \subset \mathbb{Z}^n$ for some n and κ is an adjacency on X . Thus, (X, κ) is a graph for which X is the vertex set and κ determines the edge set. Usually, X is finite, although there are papers that consider infinite X . Usually, adjacency reflects some type of “closeness” in \mathbb{Z}^n of the adjacent points. When these “usual” conditions are satisfied, one may consider the digital image as a model of a black-and-white “real world” digital image in which the black points (foreground) are the members of X and the white points (background) are members of $\mathbb{Z}^n \setminus X$.

We write $x \leftrightarrow_{\kappa} y$, or $x \leftrightarrow y$ when κ is understood or when it is unnecessary to mention κ , to indicate that x and y are κ -adjacent. Notations $x \simeq_{\kappa} y$, or $x \simeq y$ when κ is understood, indicate that x and y are κ -adjacent or are equal.

The most commonly used adjacencies are the c_u adjacencies, defined as follows. Let $X \subset \mathbb{Z}^n$ and let $u \in \mathbb{Z}$, $1 \leq u \leq n$. Then for points

$$x = (x_1, \dots, x_n) \neq (y_1, \dots, y_n) = y$$

we have $x \leftrightarrow_{c_u} y$ if and only if

- for at most u indices i we have $|x_i - y_i| = 1$, and
- for all indices j , $|x_j - y_j| \neq 1$ implies $x_j = y_j$.

The c_u -adjacencies are often denoted by the number of adjacent points a point can have in the adjacency. E.g.,

- in \mathbb{Z} , c_1 -adjacency is 2-adjacency;
- in \mathbb{Z}^2 , c_1 -adjacency is 4-adjacency and c_2 -adjacency is 8-adjacency;

- in \mathbb{Z}^3 , c_1 -adjacency is 8-adjacency, c_2 -adjacency is 18-adjacency, and c_3 -adjacency is 26-adjacency.

We discuss the *digital n -cycle*, the n -point image $C_n = \{x_0, \dots, x_{n-1}\}$ in which each x_i is adjacent only to x_{i+1} and x_{i-1} , and subscripts are always read modulo n .

The literature also contains several adjacencies to exploit properties of Cartesian products of digital images. These include the following.

Definition 2.1 ([1]). Let (X, κ) and (Y, λ) be digital images. The *normal product adjacency* or *strong adjacency* on $X \times Y$, denoted $NP(\kappa, \lambda)$, is defined as follows. Given $x_0, x_1 \in X$, $y_0, y_1 \in Y$ such that

$$p_0 = (x_0, y_0) \neq (x_1, y_1) = p_1,$$

we have $p_0 \leftrightarrow_{NP(\kappa, \lambda)} p_1$ if and only if one of the following is valid:

- $x_0 \leftrightarrow_{\kappa} x_1$ and $y_0 = y_1$, or
- $x_0 = x_1$ and $y_0 \leftrightarrow_{\lambda} y_1$, or
- $x_0 \leftrightarrow_{\kappa} x_1$ and $y_0 \leftrightarrow_{\lambda} y_1$.

Theorem 2.2 ([9]). Let $X \subset \mathbb{Z}^m$, $Y \subset \mathbb{Z}^n$. Then

$$(X \times Y, NP(c_m, c_n)) = (X \times Y, c_{m+n}),$$

i.e., the c_{m+n} -adjacency on $X \times Y \subset \mathbb{Z}^{m+n}$ coincides with the normal product adjacency based on c_m and c_n .

Building on the normal product adjacency, we have the following.

Definition 2.3 ([5]). Given $u, v \in \mathbb{N}$, $1 \leq u \leq v$, and digital images (X_i, κ_i) , $1 \leq i \leq v$, let $X = \prod_{i=1}^v X_i$. The adjacency $NP_u(\kappa_1, \dots, \kappa_v)$ for X is defined as follows. Given $x_i, x'_i \in X_i$, let

$$p = (x_1, \dots, x_v) \neq (x'_1, \dots, x'_v) = q.$$

Then $p \leftrightarrow_{NP_u(\kappa_1, \dots, \kappa_v)} q$ if for at least 1 and at most u indices i we have $x_i \leftrightarrow_{\kappa_i} x'_i$ and for all other indices j we have $x_j = x'_j$.

Notice $NP(\kappa, \lambda) = NP_2(\kappa, \lambda)$ [5].

When (X, κ) is understood to be a digital image under discussion, we use the following notations. For $x \in X$,

$$N(x) = \{y \in X \mid y \leftrightarrow_{\kappa} x\},$$

$$N^*(x) = \{y \in X \mid y \leftrightarrow_{\kappa} x\} = N(x) \cup \{x\}.$$

2.2. Digitally continuous functions. We denote by id or id_X the identity map $\text{id}(x) = x$ for all $x \in X$.

Definition 2.4 ([14, 3]). Let (X, κ) and (Y, λ) be digital images. A function $f : X \rightarrow Y$ is (κ, λ) -continuous, or *digitally continuous* or just *continuous* when κ and λ are understood, if for every κ -connected subset X' of X , $f(X')$ is a λ -connected subset of Y . If $(X, \kappa) = (Y, \lambda)$, we say a function is κ -continuous to abbreviate “ (κ, κ) -continuous.”

Theorem 2.5 ([3]). *A function $f : X \rightarrow Y$ between digital images (X, κ) and (Y, λ) is (κ, λ) -continuous if and only if for every $x, y \in X$, if $x \leftrightarrow_{\kappa} y$ then $f(x) \leftrightarrow_{\lambda} f(y)$.*

Theorem 2.6 ([3]). *Let $f : (X, \kappa) \rightarrow (Y, \lambda)$ and $g : (Y, \lambda) \rightarrow (Z, \mu)$ be continuous functions between digital images. Then $g \circ f : (X, \kappa) \rightarrow (Z, \mu)$ is continuous.*

A path is a continuous function $r : [0, m]_{\mathbb{Z}} \rightarrow X$.

We use the following notation. For a digital image (X, κ) ,

$$C(X, \kappa) = \{f : X \rightarrow X \mid f \text{ is continuous}\}.$$

Definition 2.7 ([3]; see also [13]). Let X and Y be digital images. Let $f, g : X \rightarrow Y$ be (κ, κ') -continuous functions. Suppose there is a positive integer m and a function $h : X \times [0, m]_{\mathbb{Z}} \rightarrow Y$ such that

- for all $x \in X$, $h(x, 0) = f(x)$ and $h(x, m) = g(x)$;
- for all $x \in X$, the induced function $h_x : [0, m]_{\mathbb{Z}} \rightarrow Y$ defined by

$$h_x(t) = h(x, t) \text{ for all } t \in [0, m]_{\mathbb{Z}}$$

is (c_1, κ') -continuous. That is, $h_x(t)$ is a path in Y .

- for all $t \in [0, m]_{\mathbb{Z}}$, the induced function $h_t : X \rightarrow Y$ defined by

$$h_t(x) = h(x, t) \text{ for all } x \in X$$

is (κ, κ') -continuous.

Then h is a *digital (κ, κ') -homotopy between f and g* , and f and g are *digitally (κ, κ') -homotopic in Y* , denoted $f \simeq_{\kappa, \kappa'} g$ or $f \simeq g$ when κ and κ' are understood. If $(X, \kappa) = (Y, \kappa')$, we say f and g are κ -*homotopic* to abbreviate “ (κ, κ) -homotopic” and write $f \simeq_{\kappa} g$ to abbreviate “ $f \simeq_{\kappa, \kappa} g$ ”.

If there is a κ -homotopy between id_X and a constant map, we say X is κ -*contractible*, or just *contractible* when κ is understood.

Definition 2.8. Let $A \subseteq X$. A κ -continuous function $r : X \rightarrow A$ is a *retraction*, and A is a *retract of X* , if $r(a) = a$ for all $a \in A$. If such a map r satisfies $i \circ r \simeq_{\kappa} \text{id}_X$ where $i : A \rightarrow X$ is the inclusion map, then r is a κ -*deformation retraction* and A is a κ -*deformation retract of X* .

A topological space X has the *fixed point property (FPP)* if every continuous $f : X \rightarrow X$ has a fixed point. A similar definition has appeared in digital topology: a digital image (X, κ) has the *fixed point property (FPP)* if every κ -continuous $f : X \rightarrow X$ has a fixed point. However, this property turns out to be trivial, in the sense of the following.

Theorem 2.9 ([8]). *A digital image (X, κ) has the FPP if and only if $\#X = 1$.*

The proof of Theorem 2.9 was due to the establishment of the following.

Lemma 2.10 ([8]). *Let (X, κ) be a digital image, where $\#X > 1$. Let $x_0, x_1 \in X$ be such that $x_0 \leftrightarrow_{\kappa} x_1$. Then the function $f : X \rightarrow X$ given by $f(x_0) = x_1$ and $f(x) = x_0$ for $x \neq x_0$ is κ -continuous and has 0 fixed points.*




FIGURE 1. A rigid image in \mathbb{Z}^2 with 8-adjacency

A function $f : (X, \kappa) \rightarrow (Y, \lambda)$ is an *isomorphism* (called a *homeomorphism* in [2]) if f is a continuous bijection such that f^{-1} is continuous.

3. RIGIDITY

We will say a function $f : X \rightarrow Y$ is *rigid* when no continuous map is homotopic to f except f itself. This generalizes a definition in [11]. When the identity map $\text{id} : X \rightarrow X$ is rigid, we say X is rigid.

Many digital images are rigid, though it can be difficult to show directly that a given example is rigid. A computer search described in [15] has shown that no rigid images in \mathbb{Z}^2 with 4-adjacency exist having fewer than 13 points, and no rigid images in \mathbb{Z}^2 with 8-adjacency exist having fewer than 10 points. We will demonstrate some methods for showing that a given image is rigid. For example, the digital image in Figure 1 is rigid, as shown below in Example 3.11.

An immediate consequence of the definition of rigidity is the following.

Proposition 3.1. *Let (X, κ) be a rigid digital image such that $\#X > 1$. Then X is not κ -contractible.*

Rigidity of functions is preserved when composing with an isomorphism, as the following theorems demonstrate:

Theorem 3.2. *Let $f : X \rightarrow Y$ be rigid and $g : Y \rightarrow Z$ be an isomorphism. Then $g \circ f : X \rightarrow Z$ is rigid.*

Proof. Suppose otherwise. Then there is a homotopy $h : X \times [0, m]_{\mathbb{Z}} \rightarrow Z$ from $g \circ f$ to a map $G : X \rightarrow Z$ such that $g \circ f \neq G$. Then by Theorem 2.6, $g^{-1} \circ h : X \times [0, m]_{\mathbb{Z}} \rightarrow X$ is a homotopy from f to $g^{-1} \circ G$, and since g^{-1} is one-to-one, $f \neq g^{-1} \circ G$. This contradiction of the assumption that f is rigid completes the proof. \square

Theorem 3.3. *Let $f : X \rightarrow Y$ be rigid. Let $g : W \rightarrow X$ be an isomorphism. Then $f \circ g$ is rigid.*

Proof. Suppose otherwise. Then there is a homotopy $h : W \times [0, m]_{\mathbb{Z}} \rightarrow Y$ from $f \circ g$ to some $G : W \rightarrow Y$ such that $G \neq f \circ g$. Thus, for some $w \in W$, $G(w) \neq f \circ g(w)$. Now consider the function $h' : X \times [0, m]_{\mathbb{Z}} \rightarrow Y$ defined by

$h'(x, t) = h(g^{-1}(x), t)$. By Theorem 2.6, h' is a homotopy from $f \circ g \circ g^{-1} = f$ to $G \circ g^{-1}$. Since

$$f \circ g(w) \neq G(w) = (G \circ g^{-1})(g(w)),$$

the homotopic functions f and $G \circ g^{-1}$ differ at $g(w)$, contrary to the assumption that f is rigid. The assertion follows. \square

As an immediate corollary, we obtain:

Corollary 3.4. *If $f : X \rightarrow Y$ is an isomorphism and one of X and Y are rigid, then f is rigid.*

Proof. In the case where X is rigid, the identity map id_X is rigid. Then by Theorem 3.3 we have $f \circ \text{id}_X = f$ is rigid. In the case where Y is rigid, similarly by Theorem 3.2 we have $\text{id}_Y \circ f = f$ is rigid. \square

The corollary above can be stated equivalently as follows:

Corollary 3.5. *A digital image X is rigid if and only if every digital image Y that is isomorphic to X is rigid.*

It is easy to see that no digital image in \mathbb{Z} is rigid:

Proposition 3.6. *If $X \subset \mathbb{Z}$ is a connected digital image with c_1 adjacency and $\#X > 1$, then X is not rigid.*

Proof. A connected subset of \mathbb{Z} having more than one point takes one of the forms

$$[a, b]_{\mathbb{Z}}, \{z \in \mathbb{Z} \mid z \geq a\}, \{z \in \mathbb{Z} \mid z \leq b\}, \mathbb{Z}.$$

In all of these cases, it is easily seen that there is a deformation retraction of X to a proper subset of X . Therefore, X is not rigid. \square

We also show that a normal product of images is rigid if and only if all of its factors are rigid.

Theorem 3.7. *Let (X_i, κ_i) be digital images for each $1 \leq i \leq v$, and*

$$(X, \kappa) = \left(\prod_{i=1}^v X_i, NP_u(\kappa_1, \dots, \kappa_v) \right)$$

for some $u, 1 \leq u \leq v$. Then X is rigid if and only if X_i is rigid for each i .

Proof. First we assume X is rigid, and we will show that X_i is rigid for each i . For some i , let $h_i : X_i \times [0, m]_{\mathbb{Z}} \rightarrow X_i$ be a κ_i -homotopy from id_{X_i} to $f_i : X_i \rightarrow X_i$. Without loss of generality we may assume $m = 1$, and we will show that $h_i(x_i, 1) = x_i$, and thus $f_i = \text{id}_{X_i}$. The function $h : X \times [0, 1]_{\mathbb{Z}} \rightarrow X$ defined by

$$h(x_1, \dots, x_v, t) = (x_1, \dots, x_{i-1}, h_i(x_i, t), x_{i+1}, \dots, x_v),$$

is a homotopy. Since X is rigid we must have $h(x_1, \dots, x_v, 1) = \text{id}_X$, and this means $h_i(x_i, 1) = x_i$ as desired.

Now we prove the converse: assume X_i is rigid, and we will show X is rigid. Let $J_i : X_i \rightarrow X$ be the function

$$J_i(x) = (x_1, \dots, x_{i-1}, x, x_{i+1}, \dots, x_n).$$

Let $p_i : X \rightarrow X_i$ be the projection function,

$$p_i(x_1, \dots, x_n) = x_i.$$

Then J_i is (κ_i, κ) -continuous, where $\kappa = NP_u(\kappa_1, \dots, \kappa_n)$, and p_i is (κ, κ_i) -continuous [5, 6].

For the sake of a contradiction, suppose X is not rigid. Then there is a homotopy $h : (X, \kappa) \times [0, m]_{\mathbb{Z}} \rightarrow X$ between id_X and a function g such that for some $y = (y_1, \dots, y_n) \in X$, $g(y) \neq y$. Then for some index j , $p_j(y) \neq p_j(g(y))$. Then the function $h' : X_j \times [0, m]_{\mathbb{Z}} \rightarrow X_j$ defined by $h'(x, t) = p_j(h(J_j(x), t))$, is a homotopy from id_{X_j} to a function g_j , with

$$g_j(y_j) = p_j(h(J_j(y_j), m)) = p_j(h(y, m)) = p_j(g(y)) \neq p_j(y) = y_j,$$

contrary to the assumption that X_i is rigid. We conclude that X is rigid. \square

We have a similar result when X is a disjoint union of digital images. Let X be a digital image of the form $X = A \sqcup B$ where A and B are disjoint and no point of A is adjacent to any point of B . We say X is the disjoint union of A and B , and we write $X = A \sqcup B$.

Theorem 3.8. *Let $X = A \sqcup B$. Then X is rigid if and only if A and B are rigid.*

Proof. First we assume that X is rigid, and we will show that A is rigid. (It will follow from a similar argument that B is rigid.) Let $f : A \rightarrow A$ be any self-map homotopic to id_A , and we will show that $f = \text{id}_A$. Define $g : X \rightarrow X$ by

$$g(x) = \begin{cases} f(x) & \text{if } x \in A; \\ x & \text{if } x \in B. \end{cases}$$

Then g is continuous and homotopic to id_X , and since X is rigid we must have $g = \text{id}_X$, which means that $f = \text{id}_A$.

Now for the converse, assume that A and B are both rigid. Take some self-map $f : X \rightarrow X$ homotopic to id_X , and we will show that $f = \text{id}_X$. Since f is homotopic to the identity, we must have $f(A) \subseteq A$ and $f(B) \subseteq B$. This is because there will always be a path from any point x to $f(x)$ given by the homotopy from id_X to $f(x)$. Thus if $x \in A$ we must also have $f(x) \in A$ since there are no paths from points of A to points of B .

Since $f(A) \subseteq A$ and $f(B) \subseteq B$, there are well-defined restrictions $f_A : A \rightarrow A$ and $f_B : B \rightarrow B$, and the homotopy from id_X to f induces homotopies from id_A to f_A and id_B to f_B . Since A and B are rigid we must have $f_A = \text{id}_A$ and $f_B = \text{id}_B$, and thus $f = \text{id}_X$ as desired. \square

Since every digital image is a disjoint union of its connected components, we have:

Corollary 3.9. *A digital image X is rigid if and only if every connected component of X is rigid.*

Let X be some digital image of the form $X = A \cup B$, where $A \cap B$ is a single point x_0 , and no point of A is adjacent to any point of B except x_0 . We say $X = A \cup B$ is the *wedge of A and B* , denoted $X = A \vee B$, and x_0 is called the *wedge point* of $A \vee B$. We have the following.

Theorem 3.10. *If $X = A \vee B$ and A and B are rigid, then X is rigid.*

Proof. Let x_0 be the wedge point of $A \vee B$, and let A_0 and B_0 be the components of A and B that include x_0 . If $\#A_0 = 1$ or $\#B_0 = 1$, then the components of $A \vee B$ are in direct correspondence to the components of A and B and the result follows by Corollary 3.9. Thus we assume $\#A_0 > 1$ and $\#B_0 > 1$.

Let $h : A \vee B \times [0, m]_{\mathbb{Z}} \rightarrow A \vee B$ be a homotopy such that $h(x, 0) = x$ for all $x \in A \vee B$. Without loss of generality, $m = 1$. If the induced map h_1 is not id_X then there is a point $x' \in X$ such that $h_1(x') = h(x', 1) \neq x'$. Without loss of generality, $x' \in A$. Let $p_A : X \rightarrow A$ be the projection

$$p_A(x) = \begin{cases} x & \text{for } x \in A; \\ x_0 & \text{for } x \in B. \end{cases}$$

Since $p_A \circ h$ is a homotopy from id_A to $p_A \circ h_1$, and A is rigid, we have

$$(3.1) \quad p_A \circ h_1 = \text{id}_A.$$

Were $h_1(x') \in A$ then it would follow that

$$h_1(x') = p_A \circ h_1(x') = x',$$

contrary to our choice of x' . Therefore we have $h_1(x') \in B \setminus \{x_0\}$. But $x' \leftrightarrow h_1(x')$, so $x' = x_0$.

Since A_0 is connected and has more than 1 point, there exists $x_1 \in A$ such that $x_1 \leftrightarrow_{\kappa} x_0$. By the continuity of h_1 and choice of x_0 , we must therefore have $h_1(x_1) = x_0$, and therefore $p_A \circ h_1(x_1) = p_A(x_0) = x_0$. This contradicts statement (3.1), so the assumption that h_1 is not id_X is incorrect, and the assertion follows. \square

A *loop* is a continuous function $p : C_m \rightarrow X$.

The converse of Theorem 3.10 is not generally true. In [11] it was mentioned (without proof) that a wedge of two long cycles is in general rigid. We give a specific example:

Example 3.11. Let A and B be non-contractible simple closed curves. Then A and B are non-rigid [11]. However, $X = A \vee B$ is rigid. E.g., using $c_2 = 8$ -adjacency in \mathbb{Z}^2 , let $A =$

$$\{a_0 = (0, 0), a_1 = (1, -1), a_2 = (2, -1), a_3 = (3, 0), a_4 = (2, 1), a_5 = (1, 1)\}$$

and let $B =$

$$\{b_0 = a_0, b_1 = (-1, -1), b_2 = (-2, -1), b_3 = (-3, 0), b_4 = (-2, 1), b_5 = (-1, 1)\}.$$

By continuity, if there is a homotopy $h : X \times [0, m]_{\mathbb{Z}} \rightarrow X$ - without loss of generality, $m = 1$ - such that $h_0 = \text{id}_X$ and $h(x, 1) \neq x$, then h “pulls” [11] every point of A or of B and therefore “breaks” one of the loops of X , a contradiction since “breaking” a loop is a discontinuity. Thus no such homotopy exists. \square

4. HOMOTOPY FIXED POINT SPECTRUM

The paper [10] gave a brief treatment of homotopy-invariant fixed point theory, defining two quantities $M(f)$ and $X(f)$, respectively the minimum and maximum possible number of fixed points among all maps homotopic to f . When $f : X \rightarrow X$, clearly we will have:

$$0 \leq M(f) \leq X(f) \leq \#X.$$

We will see in the examples below that any one of these inequalities can be strict in some cases, or equality in some cases.

More generally, for some map $f : X \rightarrow X$, let $\text{Fix}(f)$ denote the set of fixed points of f . We consider the following set $S(f)$, which we call the *homotopy fixed point spectrum* of f :

$$S(f) = \{\# \text{Fix}(g) \mid g \simeq f\} \subseteq \{0, \dots, \#X\}.$$

An immediate consequence of Lemma 2.10:

Corollary 4.1. *Let (X, κ) be a connected digital image, where $\#X > 1$. Then $0 \in S(c)$, where $c \in C(X, \kappa)$ is a constant map.*

We can also consider the *fixed point spectrum* of X , defined as:

$$F(X) = \{\# \text{Fix}(f) \mid f : X \rightarrow X \text{ is continuous}\}$$

Remark 4.2. The following assertions are immediate consequences of the relevant definitions.

- If X is a digital image of only one point, then $F(X) = \{1\}$.
- If $f : X \rightarrow X$ is rigid, then $S(f) = \{\# \text{Fix}(f)\}$. If X is rigid, then $S(\text{id}) = \{\#X\}$.

Since every image X has a constant map and an identity map, we always have:

$$\{1, \#X\} \subseteq F(X).$$

The number of fixed points is always preserved by isomorphism:

Lemma 4.3. *Let X and Y be isomorphic digital images. Let $f : X \rightarrow X$ be continuous. Then there is a continuous $g : Y \rightarrow Y$ such that $\# \text{Fix}(f) = \# \text{Fix}(g)$.*

Proof. Let $G : X \rightarrow Y$ be an isomorphism. Let $A = \text{Fix}(f)$. Since G is one-to-one, $\#G(A) = \#A$. Let $g : Y \rightarrow Y$ be defined by $g = G \circ f \circ G^{-1}$. For $y_0 \in G(A)$, let $x_0 = G^{-1}(y_0)$. Then

$$g(y_0) = G \circ f \circ G^{-1}(y_0) = G \circ f(x_0) = G(x_0) = y_0.$$

Let $B = \text{Fix}(g)$. It follows that $G(A) \subseteq B$, so $\#A \leq \#B$.

Since G^{-1} is an isomorphism, it similarly follows that $\#B \leq \#A$. Thus, $\#\text{Fix}(f) = \#\text{Fix}(g)$. \square

As an immediate consequence, we have the following.

Corollary 4.4. *Let X and Y be isomorphic digital images. Then $F(X) = F(Y)$.*

There is a certain regularity to the fixed point spectrum for connected digital images. When X has only a single point, we have already remarked that $F(X) = \{1\}$. For images of more than 1 point, we will show that $F(X)$ always includes 0, 1, and $\#X$, and, provided the image is large enough, the set $F(X)$ also includes 2 and 3.

The following statements hold for connected images. We discuss the fixed point spectrum of disconnected images in terms of their connected components in Theorem 7.2 and its corollary. We begin with a simple lemma:

Lemma 4.5. *Let X be any connected digital image with $\#X > 1$. Let $x_0 \in X$, and let $0 \leq k \leq \#N^*(x_0)$. Then $k \in S(c) \subseteq F(X)$, where c is the map with image $\{x_0\}$.*

Proof. By Corollary 4.1, a constant map is homotopic to a map with no fixed points, so $0 \in S(c)$ as desired.

For $k > 0$, let $n = \#N^*(x_0)$ and write

$$N^*(x_0) = \{x_0, x_1, \dots, x_{n-1}\}.$$

Then define $f : X \rightarrow X$ by:

$$f(x) = \begin{cases} x & \text{if } x = x_i \text{ for some } i < k, \\ x_0 & \text{otherwise.} \end{cases}$$

Then f is continuous with $\text{Fix}(f) = \{x_0, \dots, x_{k-1}\}$ and thus $k \in F(X)$. Furthermore, f is homotopic to the constant map at x_0 , and so in fact $k \in S(c)$. \square

Theorem 4.6. *Let X be a connected digital image, and let $c : X \rightarrow X$ be any constant map. If $\#X \geq 2$ then*

$$\{0, 1, 2\} \subseteq S(c).$$

If $\#X \geq 3$, then

$$\{0, 1, 2, 3\} \subseteq S(c).$$

Proof. If $\#X = 2$, then X consists simply of two adjacent points. Thus $\#N^*(x) = 2$ for each $x \in X$, and so Lemma 4.5 implies that $\{0, 1, 2\} \subseteq S(c)$.

When $\#X \geq 3$, there must be some $x \in X$ with $\#N^*(x) \geq 3$. (Otherwise the image would consist only of disjoint pairs of adjacent points, which would not be connected.) Thus by Lemma 4.5 we have $\{0, 1, 2, 3\} \subseteq S(c)$. \square

Since we always have $\#X \in S(\text{id})$ and

$$S(c) \cup S(\text{id}) \subseteq F(X) \subseteq \{0, 1, \dots, \#X\},$$

the theorem above directly gives:

Corollary 4.7. *Let X be a connected digital image. If $\#X = 2$ then*

$$F(X) = \{0, 1, 2\}.$$

If $\#X > 2$, then

$$\{0, 1, 2, 3, \#X\} \subseteq F(X).$$

We have already seen that $\#X \in F(X)$ in all cases. There is an easy condition that determines whether or not $\#X - 1 \in F(X)$.

Lemma 4.8. *Let X be connected with $n = \#X > 1$. Then $n - 1 \in F(X)$ if and only if there are distinct points $x_1, x_2 \in X$ with $N(x_1) \subseteq N^*(x_2)$.*

Proof. Suppose there are points $x_1, x_2 \in X$, $x_1 \neq x_2$, such that $N(x_1) \subseteq N^*(x_2)$. Then the map

$$f(x_1) = x_2, \quad f(x) = x \text{ for all } x \neq x_1,$$

is a self-map on X with exactly $n - 1$ fixed points. That f is continuous is seen as follows. Suppose $x, x' \in X$ with $x \leftrightarrow x'$.

- If $x_1 \notin \{x, x'\}$, then

$$f(x) = x \leftrightarrow x' = f(x').$$

- If, say, $x = x_1$, then $x' \in N(x_1) \subseteq N^*(x_2)$, so

$$f(x') = x' \leftrightarrow x_2 = f(x_1).$$

Thus f is continuous, and we conclude $n - 1 \in F(X)$.

Now assume that $n - 1 \in F(X)$. Thus there is some continuous self-map f with exactly $n - 1$ fixed points. Let x_1 be the single point not fixed by f , and let $x_2 = f(x_1)$. Then let $x \in X$ with $x \leftrightarrow x_1$. Then

$$x = f(x) \leftrightarrow f(x_1) = x_2,$$

so $N(x_1) \subseteq N^*(x_2)$. □

Lemma 4.8 can be used to show that a large class of digital images will satisfy $n - 1 \notin F(X)$. For example when $X = C_n$ for $n > 4$, no $N(x_i)$ is contained in $N^*(x_j)$ for $j \neq i$. Thus we have:

Corollary 4.9. *Let $n > 4$. Then $n - 1 \notin F(C_n)$.*

In particular this means that $4 \notin F(C_5)$, so the result of Theorem 4.7 cannot in general be improved to state that $4 \in F(X)$ for all images of more than 4 points.

5. PULL INDICES

Let $\overline{\text{Fix}}(f)$ be the complement of the fixed point set, that is,

$$\overline{\text{Fix}}(f) = \{x \in X \mid f(x) \neq x\}.$$

When $f(x) \neq x$, we say f moves x .

Definition 5.1. Let (X, κ) be a digital image with $\#X > 1$ and let $x \in X$. The pull index of x , $P(x)$ or $P(x, X)$ or $P(x, X, \kappa)$, is

$$P(x) = \min\{\#\overline{\text{Fix}}(f) \mid f : X \rightarrow X \text{ is continuous and } f(x) \neq x\}.$$

When $f(x) \neq x$, the set $\overline{\text{Fix}}(f)$ always contains at least the point x , and so $P(x) \geq 1$ for any x that is moved by some f .

Example 5.2. Let $X = [1, 3]_{\mathbb{Z}}$ with c_1 -adjacency.

To compute $P(3)$, consider the function $f(x) = \min\{x, 2\}$. This is continuous, not the identity, and $\text{Fix}(f) = \{1, 2\}$, and thus $P(3) = 1$. Similarly we can show that $P(1) = 1$.

But we have $P(2) = 2$, since any continuous self-map f on X that moves 2 must also move at least one other point: if $f(2) = 1$ we must have $f(3) \in \{1, 2\}$, and if $f(2) = 3$ we must have $f(1) \in \{2, 3\}$.

Proposition 5.3. Let (X, κ) be a connected digital image with $n = \#X > 1$. Let $m \in \mathbb{N}$, $1 \leq m \leq n$. Suppose, for all $x \in X$, we have $P(x) \geq m$. Then

$$F(X) \cap \{i\}_{i=n-m+1}^{n-1} = \emptyset.$$

Proof. By hypothesis, $f \in C(X, \kappa) \setminus \{\text{id}_X\}$ implies f moves at least m points, hence $\#\text{Fix}(f) \leq n - m$. The assertion follows. \square

Theorem 5.4. Let (X, κ) be a connected digital image with $n = \#X > 1$. The following are equivalent.

- 1) $n - 1 \in F(X)$.
- 2) There are distinct $x_1, x_2 \in X$ such that $N(x_1) \subseteq N^*(x_2)$.
- 3) There exists $x \in X$ such that $P(x) = 1$.

Proof. 1) \Leftrightarrow 2) is shown in Lemma 4.8.

1) \Leftrightarrow 3): We have $n - 1 \in F(X) \Leftrightarrow$ there exists $f \in C(X)$ with exactly $n - 1$ fixed points, i.e., the only $x \in X$ not fixed by f has $P(x) = 1$. \square

The following generalizes 1) \Rightarrow 3) of Theorem 5.4.

Proposition 5.5. Let (X, κ) be a connected digital image with $n = \#X > 1$. Let $k \in [1, n-1]_{\mathbb{Z}}$. Then $k \in F(X)$ implies there exist distinct $x_1, \dots, x_{n-k} \in X$ such that $P(x_i) \leq n - k$.

Proof. $k \in F(X)$ implies there exists $f \in C(X)$ with exactly k fixed points, hence distinct $x_1, \dots, x_{n-k} \in X$ such that $x_i \notin \text{Fix}(f)$. Thus for each i , the members of $\text{Fix}(f)$ are not pulled by f and x_i . Thus $P(x_i) \leq n - k$. \square

6. RETRACTS

In this section, we study how retractions interact with fixed point spectra.

Theorem 6.1 ([2]). Let (X, κ) be a digital image and let $A \subseteq X$. Then A is a retract of X if and only if for every continuous $f : (A, \kappa) \rightarrow (Y, \lambda)$ there is an extension of f to a continuous $g : (X, \kappa) \rightarrow (Y, \lambda)$.

In the proof of Theorem 6.1, an extension of f is obtained by using $g = f \circ r$, where $r : X \rightarrow A$ is a retraction. We use this in the proof of the next assertion.

Theorem 6.2. *Let A be a retract of (X, κ) . Then $F(A) \subseteq F(X)$.*

Proof. Let $f : A \rightarrow A$ be κ -continuous. Let $r : X \rightarrow A$ be a κ -retraction. Let $i : A \rightarrow X$ be the inclusion function. By Theorem 2.6, $G = i \circ f \circ r : X \rightarrow X$ is continuous. Further, $G(x) = f(x)$ if and only if $x \in A$, so $\text{Fix}(G) = \text{Fix}(f)$. Since f was taken arbitrarily, the assertion follows. \square

Remark 6.3. We do not have an analog to Theorem 6.2 by replacing fixed point spectra by spectra of identity maps. E.g, in Example 3.11 we have $\{0, \#A\} \subseteq S(\text{id}_A)$, and A is a retract of X , but X is rigid, so $S(\text{id}_X) = \{\#X\}$. However, we have the following Corollaries 6.4 and 6.5.

Corollary 6.4. *Let A be a deformation retract of X . Then $S(\text{id}_A) \subseteq S(\text{id}_X) \subseteq F(X)$. In particular, $\#A \in S(\text{id}_X)$.*

Corollary 6.5. *Let $a, b \in \mathbb{Z}$, $a < b$. Then*

$$S(\text{id}_{[a,b]_{\mathbb{Z}}}, c_1) = F([a,b]_{\mathbb{Z}}, c_1) = \{0, 1, \dots, b - a + 1\}.$$

Proof. Since $a < b$ and $[a,b]_{\mathbb{Z}}$ is c_1 -contractible, it follows from Theorem 2.9 that $0 \in S(\text{id}_{[a,b]_{\mathbb{Z}}}, c_1)$. Since for each $d \in [a,b]_{\mathbb{Z}}$ there is a c_1 -deformation of $[a,b]_{\mathbb{Z}}$ to $[a,d]_{\mathbb{Z}}$, it follows from Corollary 6.4 that $\#[a,d]_{\mathbb{Z}} \in S(\text{id}_{[a,b]_{\mathbb{Z}}}, c_1)$. Thus,

$$F([a,b]_{\mathbb{Z}}, c_1) \subseteq \{i\}_{i=0}^{b-a+1} = S(\text{id}_{[a,b]_{\mathbb{Z}}}, c_1) \subseteq F([a,b]_{\mathbb{Z}}, c_1).$$

The assertion follows. \square

We can generalize this result about intervals to a two-dimensional box in \mathbb{Z}^2 .

Theorem 6.6. *Let $X = [1, a]_{\mathbb{Z}} \times [1, b]_{\mathbb{Z}}$, with adjacency $\kappa \in \{c_1, c_2\}$. Then*

$$S(\text{id}_X) = F(X) = \{0, 1, \dots, ab\}$$

Proof. All self-maps on $[1, a]_{\mathbb{Z}} \times [1, b]_{\mathbb{Z}}$ are homotopic to the identity, so it suffices only to show that $F(X) = \{0, 1, \dots, ab\}$. The proof is by induction on a . For $a = 1$, our image X is isomorphic to the one-dimensional image $[1, b]_{\mathbb{Z}}$. Thus by Theorem 6.5 we have

$$F(X) = \{0, 1, \dots, b\} = \{0, 1, \dots, ab\}$$

as desired.

For the inductive step, first note that $[1, a - 1]_{\mathbb{Z}} \times [1, b]_{\mathbb{Z}}$ is a retract of X (using either $\kappa = c_1$ or c_2). Thus by induction and Theorem 6.2 we have

$$\{0, 1, \dots, (a - 1)b\} \subseteq F(X).$$

It remains only to show that

$$\{(a - 1)b + 1, (a - 1)b + 2, \dots, ab\} \subseteq F(X).$$

We do this by exhibiting a family of self-maps of X having these numbers of fixed points.




FIGURE 2. The map f_t from Theorem 6.6, pictured in the case $t = 2$. All points are fixed except those with arrows indicating where they map to.

Let $t \in \{0, \dots, b - 1\}$, and define $f_t : X \rightarrow X$ as follows:

$$f_t(x, y) = \begin{cases} (x, y) & \text{if } x > 1 \text{ or } y > t, \\ (x + 1, y + 1) & \text{if } x = 1 \text{ and } y \leq t \end{cases}$$

See Figure 2 for a pictorial depiction of f_t . This f_t is well-defined and both c_1 - and c_2 -continuous for each $t \in \{0, \dots, b - 1\}$ and has $ab - t$ fixed points. Thus we have

$$\{ab, ab - 1, \dots, ab - (b - 1) = (a - 1)b + 1\} \subseteq F(X)$$

as desired. □

7. CARTESIAN PRODUCTS AND DISJOINT UNIONS

In the following, assume $A_i \subset \mathbb{N}$, $1 \leq i \leq v$. Define

$$\bigotimes_{i=1}^v A_i = \left\{ \prod_{i=1}^v a_i \mid a_i \in A_i \right\}$$

and

$$\bigoplus_{i=1}^v A_i = \left\{ \sum_{i=1}^v a_i \mid a_i \in A_i \right\}.$$

If $f_i : X_i \rightarrow Y_i$, let $\prod_{i=1}^v f_i : \prod_{i=1}^v X_i \rightarrow \prod_{i=1}^v Y_i$ be the product function defined by

$$\prod_{i=1}^v f_i(x_1, \dots, x_v) = (f_1(x_1), \dots, f_v(x_v)) \text{ for } x_i \in X_i.$$

Theorem 7.1. *Suppose (X_i, κ_i) is a digital image, $1 \leq i \leq v$. Let $X = \prod_{i=1}^v X_i$. Then $\bigotimes_{i=1}^v F(X_i, \kappa_i) \subseteq F(X, NP_v(\kappa_1, \dots, \kappa_v))$.*

Proof. Let $f_i : X_i \rightarrow X_i$ be κ_i -continuous. Let $X = \prod_{i=1}^v X_i$. Then the product function

$$f = \prod_{i=1}^v f_i(x_1, \dots, x_v) : X \rightarrow X$$

is $NP_v(\kappa_1, \dots, \kappa_v)$ -continuous [5]. If $A_i = \{y_{i,j}\}_{j=1}^{p_i}$ is the set of distinct fixed points of f_i , then each point $(y_{1,j_1}, \dots, y_{v,j_v})$, for $1 \leq j_i \leq p_i$, is a fixed point of f . The assertion follows. □

We note that the conclusion of Theorem 7.1 cannot in general be strengthened to say that $\bigotimes_{i=1}^v F(X_i) = F(X)$. For example, if $X = [1, 3]_{\mathbb{Z}} \times [1, 3]_{\mathbb{Z}}$, we have $F(X) = \{0, 1, \dots, 9\}$ by Theorem 6.6, but

$$F([1, 3]_{\mathbb{Z}}) \otimes F([1, 3]_{\mathbb{Z}}) = \{0, 1, 2, 3\} \otimes \{0, 1, 2, 3\} = \{0, 1, 2, 3, 4, 6, 9\}.$$

We do have a similar result, this time with equality, for a disjoint union of digital images.

Theorem 7.2. *Let $X = A \sqcup B$. If A and B both have at least 2 points, then*

$$F(X) = F(A) \oplus F(B).$$

Proof. First we show that $F(A) \oplus F(B) \subseteq F(X)$. Take some $k \in F(A) \oplus F(B)$, say $k = m + n$ with $m \in F(A)$ and $n \in F(B)$. That means there are two self-maps $f : A \rightarrow A$ and $g : B \rightarrow B$ with $\# \text{Fix}(f) = m$ and $\# \text{Fix}(g) = n$. Let $h : X \rightarrow X$ be defined by:

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B \end{cases}$$

Then

$$\# \text{Fix}(h) = \# \text{Fix}(f) + \# \text{Fix}(g) = m + n = k$$

and so $k \in F(X)$ as desired.

Next we show $F(X) \subseteq F(A) \oplus F(B)$. Take some $k \in F(X)$, so there is some self-map f with $\# \text{Fix}(f) = k$. Let $f_A : A \rightarrow X$ and $f_B : B \rightarrow X$ be the restrictions of f to A and B . Since $X = A \cup B$, we have

$$\text{Fix}(f) = \text{Fix}(f_A) \cup \text{Fix}(f_B),$$

and $\text{Fix}(f_A) = \text{Fix}(f) \cap A$ and $\text{Fix}(f_B) = \text{Fix}(f) \cap B$. Since A and B are disjoint, the union of the fixed point sets above is disjoint. Thus we have $k = \# \text{Fix}(f_A) + \# \text{Fix}(f_B)$.

Since continuous functions preserve connectedness, we must have $f_A(A) \subseteq A$ or $f_A(A) \subseteq B$. Similarly $f_B(B) \subseteq A$ or $f_B(B) \subseteq B$. We show that $k \in F(A) \oplus F(B)$ in several cases.

In the case where $f_A(A) \subseteq B$ and $f_B(B) \subseteq A$, there are no fixed points of f_A or f_B , and thus no fixed points of f . Thus $k = 0$, and it is true that $k \in F(A) \oplus F(B)$ since $0 \in F(A)$ and $0 \in F(B)$ by Theorem 4.6.

In the case where $f_A(A) \subseteq B$ and $f_B(B) \subseteq B$, there are no fixed points of f_A , and thus $\text{Fix}(f) = \text{Fix}(f_B)$. In this case in fact f_B is a self-map of B , and so

$$k = \# \text{Fix}(f) = 0 + \# \text{Fix}(f_B) \in F(A) \oplus F(B)$$

since $0 \in F(A)$ by Theorem 4.6 and $\# \text{Fix}(f_B) \in F(B)$ since f_B is a self-map on B . The case where $f_A(A) \subseteq A$ and $f_B(B) \subseteq A$ is similar.

The final case is when $f_A(A) \subseteq A$ and $f_B(B) \subseteq B$. In this case f_A is a self-map of A and f_B is a self-map of B . Since $\text{Fix}(f) = \text{Fix}(f_A) \cup \text{Fix}(f_B)$, the k fixed points of f must partition into m fixed points of f_A and n fixed points of f_B , where $m + n = k$. Thus $m \in F(A)$ and $n \in F(B)$, and so $k = m + n \in F(A) \oplus F(B)$. \square

The assumption above that A and B have at least 2 points is necessary. For example if A and B are each a single point, then $F(X) = \{0, 1, 2\}$ while $F(A) = F(B) = \{1\}$ and thus $F(A) \oplus F(B) = \{2\}$.

Since any digital image is a disjoint union of its connected components, we have:

Corollary 7.3. *Let X_1, \dots, X_k be the connected components of a digital image X , and assume that $\#X_i > 1$ for all i . Then we have:*

$$F(X) = \bigoplus_{i=1}^k F(X_i)$$

8. LOCATIONS OF FIXED POINTS

In many cases, the existence of two fixed points will imply that other fixed points must exist in certain locations. In some cases we will show that $\text{Fix}(f)$ must be connected. We do not have $\text{Fix}(f)$ connected in general, as shown by the following.

Example 8.1. Let $X = \{p_0 = (0, 0), p_1 = (1, 0), p_2 = (2, 0), p_3 = (1, 1)\}$. Let $f : X \rightarrow X$ be defined by

$$f(p_0) = p_0, \quad f(p_1) = p_3, \quad f(p_2) = p_2, \quad f(p_3) = p_1.$$

Then X is c_2 -connected, $f \in C(X, c_2)$, and $\text{Fix}(f) = \{p_0, p_2\}$ is c_2 -disconnected.

Lemma 8.2. *Let (X, κ) be a digital image and $f : X \rightarrow X$ be continuous. Suppose that $x, x' \in \text{Fix}(f)$ and that $y \in X$ lies on every path of minimal length between x and x' . Then $y \in \text{Fix}(f)$.*

Proof. Let k be the minimal length of a path from x to x' . First we show that y must occur at the same location along any minimal path from x to x' . That is, we show that there is some $i \in [0, k]_{\mathbb{Z}}$ with $p(i) = y$ for every minimal path p from x to x' . This we prove by contradiction: assume we have two minimal paths p and q with $p(i) = y = q(j)$ for some $j < i$. Then construct a new path r by traveling from x to y along q , and then from y to x' along p . Then this path r has length less than the length of p , contradicting the minimality of p .

Thus we have some $i \in [0, k]_{\mathbb{Z}}$ with $p(i) = y$ for every minimal path p from x to x' . Let p be some minimal path from x to x' , and since the endpoints of p are fixed, the path $f(p)$ is also a path from x to x' . Furthermore the length of $f(p)$ must be at most k , and thus must equal k since this is the minimal possible length of a path from x to x' .

Since both p and $f(p)$ are minimal paths from x to x' , we have $p(i) = f(p(i)) = y$, and thus $y = f(y)$ as desired. \square

A vertex v of a connected graph (X, κ) is an *articulation point* of X if $(X \setminus \{v\}, \kappa)$ is disconnected. We have the following immediate consequences of Lemma 8.2.

Corollary 8.3. *Let (X, κ) be a connected digital image. Let v be an articulation point of X . Suppose $f \in C(X, \kappa)$ has fixed points in distinct components of $X \setminus \{v\}$. Then v is a fixed point of f .*

Corollary 8.4. *Let (X, κ) be a digital image and $f \in C(X, \kappa)$. Suppose $x, x' \in \text{Fix}(f)$ are such that there is a unique shortest κ -path P in X from x to x' . Then $P \subseteq \text{Fix}(f)$.*

Proof. This follows immediately from Lemma 8.2. □

Corollary 8.5. *Let (X, κ) be a digital image that is a tree. Then $f \in C(X, \kappa)$ implies $\text{Fix}(f)$ is κ -connected.*

Proof. This follows from Corollary 8.4, since given x, x' in a tree X , there is a unique shortest path in X from x to x' . □

For a digital cycle, the fixed point set is typically connected. The only exception is in a very particular case, as we see below.

Theorem 8.6. *Let $f : C_n \rightarrow C_n$ be any continuous map. Then $\text{Fix}(f)$ is connected, or is a set of 2 nonadjacent points. The latter case occurs only when n is even and the two fixed points are at opposite positions in the cycle.*

Proof. If $\#\text{Fix}(f) \in \{0, 1\}$, then $\text{Fix}(f)$ is connected. When $\#\text{Fix}(f) > 1$, we show that if $x_i, x_j \in \text{Fix}(f)$ are two distinct fixed points, then either there is a path from x_i to x_j through other fixed points, or that no other points are fixed.

There are two canonical paths p and q from x_i to x_j : the two injective paths going in either “direction” around the cycle. Without loss of generality assume $|p| \geq |q|$. This means that $|q|$ is the shortest possible length of a path from x_i to x_j .

Consider the case in which $|p| > |q|$. In this case $|q|$ is the unique shortest path from x_i to x_j , and by Lemma 8.4, $q \subseteq \text{Fix}(f)$, and so x_i and x_j are connected by a path of fixed points as desired.

Now consider the case in which $|p| = |q|$. In this case again $|q|$ is the shortest possible length of a path from x_i to x_j , and p and q are the only two paths from x_i to x_j having this length. Then $f(q)$ is a path from x_i to x_j of length $|q|$, and so we must have either $f(q) = q$ or $f(q) = p$. In the former case, q is a path of fixed points connecting x_i and x_j as desired. In the latter case, $\text{Fix}(f) \cap q = \{x_i, x_j\}$.

Similarly considering the path $f(p)$, we must have either $f(p) = p$ (in which case p is a path of fixed points connecting x_i and x_j); or $f(p) = f(q)$, in which case $\text{Fix}(f) \cap p = \{x_i, x_j\}$.

Considering all cases, either a minimal-length path from x_i to x_j is contained in $\text{Fix}(f)$, or $\text{Fix}(f) = \{x_i, x_j\}$.

The second sentence of the theorem follows from our analysis of the various cases. The only case which gives 2 nonadjacent fixed points requires x_i and x_j to be opposite points on the cycle, which requires n to be even. □




FIGURE 3. A contractible image for which $F(X) \neq \{0, 1, \dots, \#X\}$.

9. REMARKS AND EXAMPLES

In classical topology $M(f)$ is the only interesting homotopy invariant count of the number of fixed points. $S(f)$ is not studied in classical topology, since in all typical cases (all continuous maps on polyhedra) we would have $S(f) = [M(f), \infty)_{\mathbb{Z}}$.

In classical topology the value of $M(f)$ is generally hard to compute. The Lefschetz number gives a very rough indication of homotopy invariant fixed point information, and the more sophisticated Nielsen number is a homotopy invariant lower bound for $M(f)$. See [12].

When X is contractible, all self-maps are homotopic, so $S(f) = F(X)$ for any self-map f . It is natural to suspect that when X is contractible with $\#X > 1$, we will always have $F(X) = \{0, 1, \dots, \#X\}$. This is false, however, as the following example shows:

Example 9.1. Let $X \subset \mathbb{Z}^3$ be the unit cube of 8 points with c_1 adjacency, shown in Figure 3. Then X is contractible, so $S(f) = F(X)$ for any self-map f . By projecting the cube into one of its faces, we see that X retracts to C_4 , and since $F(C_4) = \{0, 1, 2, 3, 4\}$, we have $\{0, 1, 2, 3, 4\} \subseteq F(X)$ by Theorem 6.2.

In fact there are also continuous maps having 5 or 6 fixed points: Let:

$$g(x_5) = x_0, \quad g(x_6) = x_3, \quad g(x_i) = x_i \text{ for } i \notin \{5, 6\}$$

Then g is continuous with 6 fixed points. Let:

$$h(x_5) = h(x_7) = x_0, \quad h(x_6) = x_3, \quad h(x_i) = x_i \text{ for } i \notin \{5, 6, 7\}$$

Then h is continuous with 5 fixed points. Since of course the identity map has 8 fixed points, we have so far shown that $\{0, 1, 2, 3, 4, 5, 6, 8\} \subseteq F(X)$.

In fact $7 \notin F(X)$. This follows from Lemma 4.8. We have shown that:

$$F(X) = \{0, 1, 2, 3, 4, 5, 6, 8\}.$$

The computation of $S(f)$ in general seems to be a difficult and interesting problem. Even in the case of self-maps on the cycle C_n , the results are interesting. First we show that in fact there are exactly 3 homotopy classes of self-maps on C_n : the identity map $\text{id}(x_i) = x_i$, the constant map $c(x_i) = x_0$, and the flip map $l(x_i) = x_{-i}$.

Theorem 9.2. *Given $f \in C(C_n)$, f is homotopic to one of: a constant map, the identity map, or the flip map.*

Proof. We have noted above that if $n \leq 4$ then C_n is contractible, so every $f \in C(C_n)$ is homotopic to a constant map. Thus, in the following, we assume $n > 4$.

We can compose f by some rotation r to obtain $g = r \circ f \simeq f$ such that $g(x_0) = x_0$. We will show that g is either the identity, the flip map, or homotopic to a constant map.

If g is not a surjection, then its continuity implies $g(C_n)$ is a connected proper subset of C_n , hence is contractible. Therefore, g is homotopic to a constant map.

If g is a surjection, then g is a bijection because the domain and codomain of g both have cardinality n . By continuity, $g(x_1) \leftrightarrow g(x_0) = x_0$. Therefore, either $g(x_1) = x_{n-1}$ or $g(x_1) = x_1$.

If $g(x_1) = x_{-1}$, then continuity and the fact that g is a bijection yield an easy induction showing that $g(x_i) = x_{-i}$, $0 \leq i < n$. Therefore, g is the flip map.

If $g(x_1) = x_1$, a similar argument shows that g is the identity. \square

In fact the proof of Theorem 9.2 demonstrates the following stronger statement. Let $r_d : C_n \rightarrow C_n$ be the rotation map $r_d(x_i) = x_{i+d}$. The following generalizes Theorem 3.4 of [4], which states that any map homotopic to the identity must be a rotation.

Theorem 9.3. *Let $f : C_n \rightarrow C_n$ be continuous. Then one of the following is true:*

- f is homotopic to a constant map
- f is homotopic to the identity, and $f = r_d$ for some d
- f is homotopic to the flip map l , and $f = r_d \circ l$ for some d

The proof of Theorem 9.2 also demonstrated that all non-isomorphisms on C_n must be nullhomotopic. Thus all isomorphisms on C_n fall into the second and third categories above, and in fact all maps in those two categories are isomorphisms. Thus we obtain:

Corollary 9.4. *Let $n > 4$, and $f : C_n \rightarrow C_n$ be an isomorphism with $f \simeq g$ for some g . Then g is an isomorphism.*

Now we are ready to compute the values of $S(f)$ for our three classes of self-maps on C_n .

Theorem 9.5. *We have $S(f) = \{1\}$ for every $f : C_1 \rightarrow C_1$.*

When $1 < n \leq 4$, we have $S(f) = \{0, \dots, n\}$ for any $f : C_n \rightarrow C_n$.




FIGURE 4. The map f from Theorem 9.5 pictured in the cases $n = 5$ and $n = 6$. All points are fixed except those with arrows indicating where they map to. The path $f(C_n)$ is in bold.

When $n > 4$, let c be any constant map, id be the identity map, and l be the flip map on C_n . We have:

$$\begin{aligned}
 S(\text{id}) &= \{0, n\} \\
 S(c) &= \{0, 1, \dots, \lfloor n/2 \rfloor + 1\} \\
 S(l) &= \begin{cases} \{0, 1\} & \text{if } n \text{ is odd} \\ \{0, 2\} & \text{if } n \text{ is even} \end{cases}
 \end{aligned}$$

Proof. When $n = 1$, our image is a single point, and the constant map (which is also the identity map) is the only continuous self-map. Thus $S(f) = F(X) = \{1\}$ for every $f : C_1 \rightarrow C_1$.

When $1 < n \leq 4$, again all maps are homotopic, and we have $S(f) = F(X) = \{0, \dots, n\}$ for any f by Theorem 4.7.

Now we consider C_n with $n > 4$, which is the interesting case.

By Theorem 9.3 the only maps homotopic to id are rotation maps r_d . Since $\#\text{Fix}(r_0) = n$ and $\#\text{Fix}(r_d) = 0$ for $d \neq 0$, we have

$$S(\text{id}) = \{0, n\}.$$

Now we consider the constant map $c(x_i) = x_0$. Let $f \in C(C_n)$ be defined as follows.

$$f(x_i) = \begin{cases} x_{-i} & \text{for } 0 \leq i \leq \lfloor n/2 \rfloor; \\ x_i & \text{for } \lfloor n/2 \rfloor < i < n. \end{cases}$$

This map “folds” the cycle onto a path that is “about half” of the cycle, with $\lfloor n/2 \rfloor + 1$ fixed points. See Figure 4. This can be taken as the first step of a homotopy, in which successive steps shrink the path and the number of fixed points by one per step, until a constant map is reached at the end of the homotopy. Thus $\{1, \dots, \lfloor n/2 \rfloor + 1\} \subseteq S(c)$, and of course $0 \in S(c)$ also by Theorem 4.6.

Thus we have shown there is a fixed path p between fixed points of f , x_i, x_j , of length at least $\lfloor n/2 \rfloor + 1$. We wish to show that in fact $S(c) = \{0, \dots, \lfloor n/2 \rfloor + 1\}$. We show this by contradiction: take some nullhomotopic f , assume that $k \in S(f)$ with $k > \lfloor n/2 \rfloor + 1$, and we will show in fact that




FIGURE 5. An image having a self-map with $X(f) = 0$

all points are fixed; this would be a contradiction since $f \neq \text{id}$. Choose any $x \in C_n \setminus p$. Then x lies on the unique shortest path in C_n from x_i to x_j . Then $x \in \text{Fix}(f)$ by Lemma 8.4; this gives our desired contradiction.

Finally we consider the flip map $l(x_i) = x_{-i}$. By Theorem 9.3, all maps homotopic to l have the form $f(x_i) = r_d \circ l(x_i) = x_{d-i}$. Such a map has a fixed point at x_i if and only if $d = 2i \pmod n$. When d is odd there are no solutions, and so $\#\text{Fix}(f) = 0$. When d is even, say $d = 2a$, and n is odd, there is one solution: $i = a$. When d is even and n is also even, there are two solutions: $i = a$ and $i = a + n/2$. Thus we have some maps with no fixed points, and when k is odd we have some with one fixed point, and when k is even we have some with two. We conclude:

$$S(l) = \begin{cases} \{0, 1\} & \text{if } n \text{ is odd} \\ \{0, 2\} & \text{if } n \text{ is even} \end{cases} \quad \square$$

By Theorem 9.2, any self-map on C_n is homotopic to the constant, identity, or flip. Thus by taking unions of the sets above, we have:

Corollary 9.6.

$$F(C_n) = \begin{cases} \{1\} & \text{if } n = 1, \\ \{0, \dots, n\} & \text{if } 1 < n \leq 4, \\ \{0, 1, \dots, \lfloor n/2 \rfloor + 1, n\} & \text{if } n > 4. \end{cases}$$

From the Corollary above we see that $F(C_5) = \{0, 1, 2, 3, 5\}$, and thus the formula of Corollary 4.7 is exact for $X = C_5$. This is the only example known to the authors in which this occurs.

Question 9.7. *Is there any digital image $X \neq C_5$ with $\#X > 4$ and $F(X) = \{0, 1, 2, 3, \#X\}$?*

We conclude this section with two interesting examples showing the wide variety of fixed point sets that can be exhibited for other digital images. Tools we use in our discussion include the following.

A path $r : [0, m]_{\mathbb{Z}} \rightarrow X$ that is an isomorphism onto $r([0, m]_{\mathbb{Z}})$ is a *simple path*. If a loop p is an isomorphism onto $p(C_m)$, p is a *simple loop*.

Definition 9.8 ([11]). A simple path or a simple loop in a digital image X has *no right angles* if no pair of consecutive edges of the path or loop belong to a loop of length 4 in X .




FIGURE 6. A lasso for the points $x = (0, 1)$ and $x' = (0, 0)$




FIGURE 7. An image with many different values for $S(f)$

Definition 9.9 ([11]). A *lasso* in X is a simple loop $p : C_m \rightarrow X$ and a path $r : [0, k]_{\mathbb{Z}} \rightarrow X$ such that $k > 0$, $m \geq 5$, $r(k) = p(x_0)$, and neither $p(x_1)$ nor $p(x_{m-1})$ is adjacent to $r(k-1)$.

The lasso *has no right angles* if neither p nor r has a right angle, and no right angle is formed where r meets p , i.e., the final edge of r and neither of the edges of p at $p(x_0)$ form 2 edges of a loop of length 4 in X .

Theorem 9.10 ([11]). *Let X be an image in which, for any two adjacent points $x \leftrightarrow x' \in X$, there is a lasso with no right angles having path $r : [0, k]_{\mathbb{Z}} \rightarrow X$ with $r(0) = x$ and $r(1) = x'$. Then X is rigid.*

Example 9.11. Let X be the digital image

$$X = ([0, 6]_{\mathbb{Z}} \times \{0, 2\}) \cup \{(0, 1), (2, 1), (4, 1), (6, 1)\}$$

(see Figure 5), with 4-adjacency.

By Theorem 9.10, this image is rigid. It is easy, though a bit tedious, to verify that the hypothesis of Theorem 9.10 is satisfied by X . For example, in Figure 6 we exhibit a lasso with no right angles for two adjacent points. It is easy to construct such lassos for any pair of adjacent points.

Since X is rigid, we have $S(\text{id}) = \{\#X\} = \{18\}$. Let $f : X \rightarrow X$ be the 180-degree rotation of X . Then f is an isomorphism, and so by Theorem 3.2, $f = f \circ \text{id}_X$ is rigid. Thus $S(f) = \{\#\text{Fix}(f)\} = \{0\}$. In particular this provides an example for the question posed in [10] if $X(f)$ could ever equal 0 for a connected image.

The following example demonstrates an image which has many different possible sets which can occur as $S(f)$ for various self-maps f .

Example 9.12. Let

$$X = ([0, 5]_{\mathbb{Z}} \times \{0, 2\}) \cup \{(0, 1), (2, 1), (5, 1)\}$$

(see Figure 7), with 4-adjacency. In this image we have several different homotopy classes of maps. We will derive sufficient information about S for some of these to compute $F(X)$.

By Theorem 9.10, X is rigid, so $S(\text{id}) = \{\#X\} = \{15\}$.

Let f be a vertical reflection. Then f is rigid by Corollary 3.4, and has 3 fixed points, so $S(f) = \{3\}$.

Let g be the function that maps the bottom horizontal bar onto the top one, and fixes all other points. Then g has 9 fixed points, and is homotopic to a constant map. We can retract the image of g down to a point one point at a time, and so $\{0, 1, \dots, 9\} \subseteq S(g)$.

Let h be the function which maps the left vertical bar into the middle vertical bar and fixes all other points. Then h has 12 fixed points. We can additionally map one or both of the next two points into the middle vertical bar to obtain maps homotopic to h with 11 or 10 fixed points. We can do these retractions followed by a rotation around the 10-cycle on the right to obtain a map homotopic to h with no fixed points. Thus $\{0, 10, 11, 12\} \subseteq S(h)$.

We have $F(X) \cap \{13, 14\} = \emptyset$ by Proposition 5.3, since for all $x \in X$ we see easily that $P(x) \geq 3$.

We therefore have $F(X) = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15\}$.

10. FURTHER REMARKS

We have studied several measures concerning the fixed point set of a continuous self-map on a digital image. We anticipate further research in this area.

ACKNOWLEDGEMENTS. *We are grateful for the suggestions of an anonymous reviewer, and some careful corrections by Muhammad Sirajo Abdullahi and Jamilu Abubakar.*

REFERENCES

- [1] C. Berge, *Graphs and Hypergraphs*, 2nd edition, North-Holland, Amsterdam, 1976.
- [2] L. Boxer, Digitally continuous functions, *Pattern Recognition Letters* 15 (1994), 833–839.
- [3] L. Boxer, A classical construction for the digital fundamental group, *Journal of Mathematical Imaging and Vision* 10 (1999), 51–62.
- [4] L. Boxer, Continuous maps on digital simple closed curves, *Applied Mathematics* 1 (2010), 377–386.
- [5] L. Boxer, Generalized normal product adjacency in digital topology, *Applied General Topology* 18, no. 2 (2017), 401–427.

- [6] L. Boxer, Alternate product adjacencies in digital topology, *Applied General Topology* 19, no. 1 (2018), 21–53.
- [7] L. Boxer, Fixed points and freezing sets in digital topology, *Proceedings, 2019 Interdisciplinary Colloquium in Topology and its Applications*, in Vigo, Spain; 55–61.
- [8] L. Boxer, O. Ege, I. Karaca, J. Lopez, and J. Louwsma, Digital fixed points, approximate fixed points, and universal functions, *Applied General Topology* 17, no. 2 (2016), 159–172.
- [9] L. Boxer and I. Karaca, Fundamental groups for digital products, *Advances and Applications in Mathematical Sciences* 11, no. 4 (2012), 161–180.
- [10] L. Boxer and P. C. Staecker, Remarks on fixed point assertions in digital topology, *Applied General Topology* 20, no. 1 (2019), 135–153.
- [11] J. Haarmann, M. P. Murphy, C. S. Peters and P. C. Staecker, Homotopy equivalence in finite digital images, *Journal of Mathematical Imaging and Vision* 53 (2015), 288–302.
- [12] B. Jiang, Lectures on Nielsen fixed point theory, *Contemporary Mathematics* 18 (1983).
- [13] E. Khalimsky, Motion, deformation, and homotopy in finite spaces, in *Proceedings IEEE Intl. Conf. on Systems, Man, and Cybernetics* (1987), 227–234.
- [14] A. Rosenfeld, ‘Continuous’ functions on digital pictures, *Pattern Recognition Letters* 4 (1986), 177–184.
- [15] P. C. Staecker, Some enumerations of binary digital images, arXiv:1502.06236, 2015.