

Document downloaded from:

<http://hdl.handle.net/10251/146000>

This paper must be cited as:

Essam Elsayed-Moham Okda, H.; El Sayed, S.; Ferreira, RCM.; Costa, SP.; Raposo, M.; Martínez-Máñez, R.; Sancenón Galarza, F. (12-2). 4-(4,5-Diphenyl-1H-imidazole-2-yl)-N,N-dimethylaniline-Cu(II) complex, a highly selective probe for glutathione sensing in water-acetonitrile mixtures. *Dyes and Pigments*. 159:45-48.  
<https://doi.org/10.1016/j.dyepig.2018.05.069>



The final publication is available at

<https://doi.org/10.1016/j.dyepig.2018.05.069>


Copyright Elsevier

Additional Information

# Accepted Manuscript

4-(4,5-Diphenyl-1*H*-imidazole-2-yl)-*N,N*-dimethylaniline-Cu(II) complex, a highly selective probe for glutathione sensing in water-acetonitrile mixtures

Hazem Essam Okda, Sameh El Sayed, Rosa C.M. Ferreira, Susana P.G. Costa, M.Manuela M. Raposo, Ramón Martínez-Máñez, Félix Sancenón



PII: S0143-7208(18)30949-5

DOI: [10.1016/j.dyepig.2018.05.069](https://doi.org/10.1016/j.dyepig.2018.05.069)

Reference: DYPI 6798

To appear in: *Dyes and Pigments*

Received Date: 26 April 2018

Revised Date: 28 May 2018

Accepted Date: 28 May 2018

Please cite this article as: Okda HE, El Sayed S, Ferreira RCM, Costa SPG, Raposo MMM, Martínez-Máñez Ramón, Sancenón Fé, 4-(4,5-Diphenyl-1*H*-imidazole-2-yl)-*N,N*-dimethylaniline-Cu(II) complex, a highly selective probe for glutathione sensing in water-acetonitrile mixtures, *Dyes and Pigments* (2018), doi: 10.1016/j.dyepig.2018.05.069.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **4-(4,5-Diphenyl-1*H*-imidazol-2-yl)-*N,N*-dimethylaniline-Cu(II)**  
2 **complex, a highly selective probe for glutathione sensing in**  
3 **water-acetonitrile mixtures**

4  
5 Hazem Essam Okda,<sup>a,b,c</sup> Sameh El Sayed,<sup>a,b,c</sup> Rosa C. M. Ferreira,<sup>d</sup> Susana P. G. Costa,<sup>d</sup> M.  
6 Manuela M. Raposo,<sup>d\*</sup> Ramón Martínez-Mañez,<sup>a,b,c\*</sup> and Félix Sancenón<sup>a,b,c</sup>

7  
8 <sup>a</sup> Instituto Interuniversitario de Investigación de Reconocimiento Molecular y Desarrollo  
9 Tecnológico (IDM), Universitat Politècnica de València, Universitat de València. Spain. E-mail:  
10 rmaez@qim.upv.es

11 <sup>b</sup> Departamento de Química, Universitat Politècnica de València. Camino de Vera s/n, 46022,  
12 València, Spain.

13 <sup>c</sup> CIBER de Bioingeniería, Biomateriales y Nanomedicina (CIBER-BBN).

14 <sup>d</sup> Centro de Química, Universidade do Minho, Campus de Gualtar, 4710-057, Braga, Portugal.  
15 E-mail: mfox@quimica.uminho.pt

16  
17 **Abstract**

18 The imidazole derivative 4-(4,5-diphenyl-1*H*-imidazol-2-yl)-*N,N*-dimethylaniline (probe **1**) formed  
19 a highly coloured and non-emissive 1:1 stoichiometry complex with Cu(II) in water-acetonitrile  
20 1:1 (v/v) solutions. Among all the amino acids (Lys, Val, Gln, Leu, His, Thr, Trp, Gly, Phe, Arg,  
21 Ile, Met, Ser, Ala, Pro, Tyr, Gly, Asn, Asp, Glu, Cys and Hcy) and tripeptides (GSH) tested only  
22 GSH induced the bleaching of the 1·Cu(II) solution together with a marked emission  
23 enhancement at 411 nm (excitation at 320 nm). These chromo-fluorogenic changes were  
24 ascribed to a selective GSH-induced demetallation of the 1·Cu(II) complex that resulted in a  
25 recovery of the spectroscopic features of probe **1**. In addition to the remarkable selectivity of  
26 1·Cu(II) complex toward GSH a competitive limit of detection as low as 2 µM was determined  
27 using fluorescence measurements.

28 **Keywords**

29 GSH; chromo-fluorogenic detection; Cu(II) complex; displacement assay

30 **1. Introduction**

31 Biothiols, such as glutathione (GSH), cysteine (Cys), and homocysteine (Hcy), are biomolecules  
32 containing thiol groups.[1] Cys and Hcy are components of many peptides that have a wide  
33 range of cellular biological functions. Besides, the three biothiols play important roles in the  
34 body's biochemical defence system because they are involved in reversible redox homeostasis  
35 processes which maintain the equilibrium of reduced free thiol and oxidized disulphide forms.[2]  
36 As the most abundant reductive biothiol (with concentrations in the millimolar range in living  
37 systems), GSH mediates many cellular functions such as maintenance of intracellular redox  
38 activities, xenobiotic metabolism, intracellular signal transduction and gene regulation.[3]  
39 Moreover, abnormal levels of biothiols affect the normal physiological and pathological functions  
40 and are related with a number of diseases such as cancer, AIDS, liver damage, Alzheimer,  
41 osteoporosis, inflammatory bowel diseases and cardiovascular diseases.[4-9] In this context,  
42 several studies have been devoted to the development of efficient methods for the detection of  
43 the concentration of GSH in physiological media. Several strategies such as mass  
44 spectrometry,[10] high-performance liquid chromatography,[11,12] enzymatic methods,[13]  
45 electrochemical assays,[14] surface-enhanced Raman scattering,[15-17] and combinatorial  
46 library-based sensors[18,19] have been described for the detection and quantification of GSH.  
47 However, these methods require expensive equipment, are time-consuming and the selectivity  
48 achieved is, in some cases, low.

49 Bearing in mind the above-mentioned facts, the development of probes able to display colour  
50 and/or fluorescence changes in water or mixed aqueous solutions in the presence of target bio-  
51 relevant thiols is a timely research area.[20] Within different approaches described for the  
52 preparation of chromo-fluorogenic sensors of biothiols, the use of displacement processes  
53 involving non-emissive fluorophore-Cu(II) complexes has attracted great attention in the last  
54 years.[21] In spite of the fact that most of the reported fluorophore-Cu(II) complexes allowed  
55 GSH detection in aqueous environments their response is in general unselective and Cys and  
56 Hcy also induced emission modulations.[21] Only three recently published examples, based on

57 the use of a coumarin derivative,[22] graphitic carbon nitride,[23] and a displacement assay with  
58 an iminophenol-Cu(II) complex,[24] allowed GSH selective detection in the presence of Cys and  
59 Hcy in aqueous environments.

60 Given our interest in the design of optical chemosensors for the detection of anions and cations  
61 of biological and environmental significance [25] herein we report the selective chromo-  
62 fluorogenic detection of GSH using a complex formed by 4-(4,5-diphenyl-1*H*-imidazol-2-yl)-*N,N*-  
63 dimethylaniline (**1** in Scheme 1) and Cu(II). The emission of probe **1** was selectively quenched  
64 in the presence of Cu(II) due to the formation of 1:1 stoichiometry complex. In the presence of  
65 GSH the emission of probe **1** was fully restored due to a demetallation reaction.

## 66 **2. Experimental section**

67 *Chemicals:* Commercially available reagents 4-(dimethylamino)benzaldehyde (**1a**), 1,2-  
68 diphenylethane-1,2-dione (**1b**), and ammonium acetate were purchased from Sigma-Aldrich  
69 and Acros and used as received. TLC analyses were carried out on 0.25 mm thick pre-coated  
70 silica plates (Merck Fertigplatten Kieselgel 60F<sub>254</sub>) and spots were visualized under UV light.  
71 Chromatography on silica gel was carried out on Merck Kieselgel (230-240 mesh).

72 *Materials and methods:* All melting points were measured on a Stuart SMP3 melting point  
73 apparatus. IR spectra were determined on a BOMEM MB 104 spectrophotometer using KBr  
74 discs. NMR spectra were obtained on a Bruker Avance III 400 at an operating frequency of 400  
75 MHz for <sup>1</sup>H and 100.6 MHz for <sup>13</sup>C using the solvent peak as internal reference at 25 °C. All  
76 chemical shifts are given in ppm using <sup>0</sup>H Me<sub>4</sub>Si = 0 ppm as reference. Assignments were  
77 supported by spin decoupling-double resonance and bi-dimensional heteronuclear correlation  
78 techniques. UV/visible titration profiles were carried out with JASCO V-650 spectrophotometer  
79 (Easton, MD, USA). Fluorescence measurements were recorded with a JASCO FP-8500  
80 spectrophotometer.

81 *Synthesis of probe 1:* 4-(Dimethylamino) benzaldehyde (**1a**) (0.15 g, 1 mmol), 1,2-  
82 diphenylethane-1,2-dione (**1b**) (0.33 g, 1 mmol) and NH<sub>4</sub>OAc (1.54 g, 20 mmol) were  
83 dissolved in glacial acetic acid (5 mL), followed by stirring and heating at reflux for 8 h.  
84 The reaction mixture was cooled to room temperature, ethyl acetate was added (15 mL)  
85 and the mixture was washed with water (3 x 10 mL). After, the organic phase was dried  
86 with anhydrous MgSO<sub>4</sub>, the solution was filtered and the solvent was evaporated to

87 dryness. The resulting crude product was purified by column chromatography (silica gel,  
88 DCM/MeOH (100:01)) and was obtained as yellow oil (11 mg, 10%).


89  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  = 3.04 (s, 6H,  $\text{NMe}_2$ ), 6.84 (dd,  $J$  = 6.8 and 2.0 Hz, 2H,  
90 H2 and H6 aniline), 7.27-7.32 (m, 2H,  $2\times\text{H}_4$  Ph), 7.34-7.39 (m, 4H,  $2\times\text{H}_2$  and H6 Ph),  
91 7.61 (d,  $J$  = 8.4 Hz, 4H,  $2\times\text{H}_3$  and H5 Ph), 8.03 (dd,  $J$  = 6.8 and 2.0 Hz, 2H, H3 and H5  
92 aniline) ppm.

93  $^{13}\text{C}$  NMR (100.6 MHz,  $\text{DMSO-}d_6$ ):  $\delta$  = 40.36, 112.83, 119.39, 127.31, 127.70, 128.00,  
94 128.20, 128.70, 129.10, 134.62, 147.71, 151.70 ppm.

95 IR (liquid film):  $\nu$  = 3420, 2926, 2856, 1692, 1646, 1615, 1549, 1509, 1495, 1446, 1405, 1363,  
96 1250, 1202, 1172, 1124, 1071, 1026, 1002, 966, 945, 822, 766, 696  $\text{cm}^{-1}$ .

### 97 3. Results

98 The synthesis of 4-(4,5-diphenyl-1*H*-imidazol-2-yl)-*N,N*-dimethylaniline (probe **1**) was published  
99 elsewhere (using different catalysts such as oxalic acid,  $\text{SnCl}_4\text{-SiO}_2$ ,  $\text{H}_3\text{BO}_3$ -ultrasounds).[26] In  
100 this study we used an one-step Radziszewski reaction between 4-dimethylamino benzaldehyde  
101 (**1a**) and 1,2-diphenylethane-1,2-dione (**1b**) in the presence of ammonium acetate in acidic  
102 media which directly yielded probe **1** (see Scheme 1).




110 **Scheme 1.** (a) Synthesis of probe **1** and (b) GSH recognition mechanism using **1**-Cu(II) complex.

111

112 Probe **1** was not fully water soluble and, for this reason, we carried out the spectroscopic  
113 characterization in water-acetonitrile mixtures. In this respect, water (pH 7.4)-acetonitrile 1:1

114 (v/v) solutions of probe **1** ( $5.0 \times 10^{-5} \text{ mol L}^{-1}$ ) presented an absorption band, of charge-transfer  
 115 nature (due to the presence of a donor *N,N*-dimethylaniline moiety and an electron-deficient  
 116 imidazole heterocycle as acceptor group), centred at ca. 320 nm. In a first step, UV-visible  
 117 changes of probe **1** solutions were studied in the presence of 10 eq. of selected metal cations  
 118 (i.e. Cu(II), Pb(II), Mg(II), Ge(II), Ca(II), Zn(II), Co(II), Ni(II), Ba(II), Cd(II), Hg(II), Fe(III), In(III),  
 119 As(III), Al(III), Cr(III), Ga(III), K(I), Li(I) and Na(I)). The obtained results are shown in Figure 1.  
 120 As could be seen, among all cations tested, only Cu(II) was able to induce a remarkable  
 121 appearance of a new red-shifted band centred at ca. 490 nm. These facts were reflected in a  
 122 marked colour change from colourless to reddish-brown (Figure 1). In more detail, addition of  
 123 increasing quantities of Cu(II) induced a progressive decrease of the band centred at 320 nm  
 124 with a growth in absorbance at 490 nm (see Supporting Information for the UV-visible titration  
 125 profile of probe **1** with Cu(II)). The appearance of a red-shifted band upon addition of Cu(II) is  
 126 tentatively attributed to an interaction of this cation with the acceptor part of the probe **1**, i.e the  
 127 imidazole ring.



134 **Figure 1.** UV-visible spectra of **1** in water (pH 7.4)-acetonitrile 1:1 (v/v) ( $5.0 \times 10^{-5} \text{ mol L}^{-1}$ ) alone and in the  
 135 presence of 10 eq. of selected metal cations. The inset shows the change in colour of **1** in the presence of  
 136 Cu(II).

138 Changes in the UV-visible bands (reflected in marked colour changes) of probe **1** upon addition  
 139 of Cu(II) were ascribed to the formation of a 1:1 stoichiometry complex as was assessed from  
 140 the Job's plot shown in Figure 2. From the UV-visible titration profile a logarithm of the stability  
 141 constant for the formation of the **1**·Cu(II) complex of  $5.0 \pm 0.1$  was determined.

142


143

144

145

146

147



148 **Figure 2.** Job's plot of **1** and Cu(II) in water (pH 7.4)-acetonitrile 1:1 (v/v). Total concentration of **1** and  
 149 Cu(II) of  $2.0 \times 10^{-5} \text{ mol L}^{-1}$ .

150

151 Furthermore, probe **1** was also emissive and, upon excitation of water (pH 7.4)-acetonitrile 1:1  
 152 (v/v) solution of the probe at 320 nm, a marked fluorescence at 445 nm appeared (see Figure  
 153 3). The emission behaviour of probe **1** in the presence of selected cations was also tested. As in  
 154 the UV-visible studies, the unique cation able to induce changes in the emission band of probe  
 155 **1** was Cu(II). As could be seen in Figure 3, addition of increasing amounts of Cu(II) to water (pH  
 156 7.4)-acetonitrile 1:1 (v/v) solution of probe **1** induced a progressive quenching of the 445 nm  
 157 emission together with a moderate blue shift of the band. From the emission titration profile  
 158 obtained, a limit of detection of  $3.2 \mu\text{M}$  for Cu(II) was determined.

159


160

161

162

163

164




165 **Figure 3.** Fluorescence titration profile of **1** in water (pH 7.4)-acetonitrile 1:1 (v/v) ( $5.0 \times 10^{-5} \text{ mol L}^{-1}$ ) upon  
 166 addition of increasing amounts of Cu(II) (from 0 to 10 eq.). Inset: plot of the emission intensity at 445 nm  
 167 vs Cu(II) concentration.

168



169 Taking into account the high affinity of Cu(II) toward thiol moieties [27] we envisioned that  
170 1·Cu(II) complex could be a promising ensemble for fluorescence “off-on” detection of certain  
171 biothiols via a Cu(II) displacement approach. For this purpose, water (pH 7.4)-acetonitrile 1:1  
172 (v/v) solution of 1·Cu(II) complex ( $6.2 \times 10^{-6}$  mol L<sup>-1</sup>) were prepared and the UV-absorption  
173 behaviour in the presence of GSH (2.0 eq.) and selected amino acids (2.0 eq. of Lys, Val, Gln,  
174 Leu, His, Thr, Trp, Gly, Phe, Arg, Ile, Met, Ser, Ala, Pro, Tyr, Gly, Asn, Asp, Glu, Cys and Hcy)  
175 was tested. The obtained results are shown in Figure 4.

176 As could be seen in Figure 4, only GSH was able to induce a complete disappearance of the  
177 490 nm band ascribed to the 1·Cu(II) complex, which was reflected in a marked colour change  
178 from reddish-brown to colourless. In contrast, none of the amino acids tested induced  
179 remarkable changes in the visible band. Addition of increasing quantities of GSH induced a  
180 progressive decrease of the absorbance at 490 nm (see Supporting Information). From the  
181 titration profile a limit of detection for GSH of 3  $\mu$ M was determined.




188 **Figure 4.** UV-visible changes of water (pH 7.4)-acetonitrile 1:1 (v/v) solution of 1·Cu(II) complex ( $6.2 \times 10^{-6}$   
189 mol L<sup>-1</sup>) in the presence of GSH (2.0 eq.) and selected amino acids (also 2.0 eq.).

190

191 Additionally, the fluorescence response of water (pH 7.4)-acetonitrile 1:1 (v/v) solution of  
192 1·Cu(II) complex in the presence of GSH and selected amino acids was also tested. Upon  
193 excitation at 320 nm, water (pH 7.4)-acetonitrile 1:1 (v/v) solution of 1·Cu(II) complex presented  
194 a weak emission band centred at 404 nm (see Figure 5). Addition of amino acids induced  
195 negligible changes in the emission profile (data not shown) whereas in the presence of  
196 increasing quantities of GSH a marked emission enhancement together with a moderate red

197 shift (to 411 nm) was observed (see Figure 5). From the emission titration profile, obtained upon  
198 addition of increasing quantities of GSH, a limit of detection of 2  $\mu\text{M}$  for GSH was calculated.  
199 The obtained limit of detection is similar to that found by Jiang et al. using a coumarin-Cu(II)  
200 complex (0.36  $\mu\text{M}$ ) [22] and by Kim and co-workers using and iminophenol-Cu(II) complex (5.86  
201  $\mu\text{M}$ ).[24] However, Yan and co-workers measured a limit of detection of 0.02  $\mu\text{M}$  for GSH using  
202 graphitic carbon nitride.[23]

203 The UV-visible and emission changes obtained when GSH was added to aqueous solutions of  
204 **1**-Cu(II) complex pointed to a demetallation process as mechanism of the optical response  
205 observed. GSH is able to displace Cu(II) from the **1**-Cu(II) complex restoring the UV-visible and  
206 emission spectra of probe **1**. The absence of optical response in the presence of thiol-  
207 containing amino acids Cys and Hcy and the remarkable selectivity of the **1**-Cu(II) complex  
208 toward GSH could be ascribed to a preferential coordination of the tripeptide with Cu(II). The  
209 structure of GSH presented several potential coordinating sites (amino, sulfhydryl and  
210 carboxylates) in a flexible backbone and could coordinate Cu(II) more effectively than Cys and  
211 Hcy which presented the same functional groups but in a more rigid skeleton.[28] This **1**-Cu(II)  
212 demetallation process regenerates the optical features of the free probe.



220 **Figure 5.** Fluorescence titration profile of **1**-Cu(II) complex in water (pH 7.4)-acetonitrile 1:1 (**v/v**) ( $6.2 \times 10^{-6}$   
221  $\text{mol L}^{-1}$ ) upon addition of increasing amounts of GSH (excitation at 320 nm).

222

223 Besides, the selectivity of **1**-Cu(II) for GSH detection in the presence of other competitive  
224 biothiols (such as Cys and Hcy) was also tested (see Supporting Information). For this purpose,

225 the emission intensity of **1**-Cu(II) (at 411 nm upon excitation at 320 nm) alone, in the presence  
226 of GSH (2.0 eq.) and with a mixture of GSH+Cys+Hcy (2.0 eq. of each biothiol) was measured.  
227 The emission intensity measured in the presence of GSH and with the three biothiols are nearly  
228 the same. This fact pointed to a selective response of **1**-Cu(II) toward GSH (this biothiol is the  
229 only able to demetallate **1**-Cu(II)) and opens the possible use of this complex for the detection  
230 of this tripeptide in real samples.

#### 231 **4. Conclusions**

232 In summary, we described herein the use of probe **1** complexed with Cu(II) as selective and  
233 sensitive chromo-fluorogenic sensor for GSH. Probe **1** forms a coloured and weakly-emissive  
234 complex with Cu(II) in water (pH 7.4)-acetonitrile 1:1 (v/v) solution. Moreover, **1**-Cu(II) complex  
235 exhibits unique selectivity and sensitivity for GSH detection in aqueous environments. Addition  
236 of GSH to water (pH 7.4)-acetonitrile 1:1 (v/v) solutions of **1**-Cu(II) complex induced a marked  
237 bleaching of the colour and the appearance of an intense emission band. The optical changes  
238 were ascribed to a GSH-induced demetallation of **1**-Cu(II) complex which regenerated the free  
239 probe. The response to GSH was quite selective because other biothiols tested (Cys and Hcy)  
240 were unable to induce any colour or emission changes. Besides, the system presented a  
241 competitive limit of detection for GSH (2  $\mu$ M using emission measurements). Moreover, the  
242 **1**-Cu(II) complex is one of the few examples of chromo-fluorogenic probes which selectively  
243 detect GSH in the presence of Cys and Hcy in aqueous environments.

#### 244 **Acknowledgments**

245 We thank the Spanish Government (MAT2015-64139-C4-1-R) and Generalitat Valenciana  
246 (PROMETEOII/2014/047). H. E. O. thanks Generalitat Valenciana for his Grisolia fellowship.  
247 Thanks are also due to Fundação para a Ciência e Tecnologia (Portugal) for financial support to  
248 the Portuguese NMR network (PTNMR, Bruker Avance III 400-Univ. Minho), FCT and FEDER  
249 (European Fund for Regional Development)-COMPETE-QREN-EU for financial support to the  
250 research centre CQ/UM [PEst-C/QUI/UI0686/2013 (FCOMP-01-0124-FEDER-037302)], and a  
251 post-doctoral grant to R.M.F. Batista (SFRH/BPD/79333/2011).

#### 252 **References**

253 [1] Wood ZA, Schroder E, Harris JR, Poole LB, Trends Biochem. Sci. 2003; 28: 32-40.

- 254 [2] Dey SK, Roy S, Bull. Environ. Contam. Toxicol. 2010; 84: 385-389.
- 255 [3] (a) Sies H, Free Radic. Biol. Med. 1999; 27: 916-921; (b) Lushchak VI, J. Amino Acids  
256 2012; 736837.
- 257 [4] Biswas SK, Rahman I, Mol. Asp. Med. 2009; 30: 60-76.
- 258 [5] Jay FH, Hongqiao Z, Alessandra A, Mol. Asp. Med. 2009; 30: 1-12.
- 259 [6] William MJ, Wilson-Delfosse AL, Mieyal JJ, Nutrients 2012; 4: 1399-1440.
- 260 [7] Townsend DM, Tew KD, Tapiero H, Biomed. Pharmacother. 2003; 57: 145-155.
- 261 [8] Hilderbrand SA, Weissleder R, Curr. Opin. Chem. Biol. 2010; 14: 71-79.
- 262 [9] James ML, Gambhir SS, Physiol. Rev. 2012; 92: 897-965.
- 263 [10] Huang YF, Chang HT, Anal. Chem. 2007; 79: 4852-4859.
- 264 [11] Chwatko G, Elzbieta K, Kubalczyk P, Borowczyk K, Rokieli MW, Glowacki R, Anal.  
265 Methods 2014; 6: 8039-8044.
- 266 [12] Moore T, Anthony L, Niemi AK, Kwan T, Cusmano-Ozog K, Gregory ME, Cowan MT,  
267 J. Chromatogr. B 2013; 929: 51-55.
- 268 [13] Timur S, Odaci D, Dincer A, Zihnioglu F, Telefoncu A, Talanta 2008; 74: 1492-1497.
- 269 [14] Cavanillas S, Serrano N, Diaz-Cruz JM, Arino C, Esteban M, Electroanalysis  
270 2014; 26: 581-587.
- 271 [15] Arindam S, Nikhil JR, Anal. Chem. 2013; 85: 9221-9228.
- 272 [16] Lv M, Gu H, Yuan X, Gao J, Cai T, J. Mol. Struct. 2012; 1029: 75-80.
- 273 [17] Ouyang L, Zhu L, Jiang J, Tang H, Anal. Chim. Acta 2014; 816: 41-49.
- 274 [18] A. Buryak, A. Severin, J. Comb. Chem. 2006; 8: 540-543.
- 275 [19] Ahn YH, Lee JS, Chang YT, J. Am. Chem. Soc. 2007; 129: 4510-4511.
- 276 [20] (a) Liu Y, Hu Y, Lee S, Lee D, Yoon J, Bull. Kor. Chem. Soc. 2016; 37: 1661-1678; (b)  
277 Yin CX, Xiong KM, Huo FJ, Salamanca JC, Strongin RM, Angew. Chem. Int. Ed. 2017;  
278 56: 13188-13198; (c) Jung HS, Chen X, Kim JS, Yoon Y, Chem. Soc. Rev. 2013; 42:  
279 6019-6031.
- 280 [21] (a) Hu Y, Heo CH, Kim G, Jun EJ, Yin J, Kim HM, Yoon J, Anal. Chem. 2015; 87:  
281 3308-3313; (b) Shi YG, Yao JH, Duan YL, Mi QL, Chen JH, Xu QQ, Gou GZ, Zhou Y,  
282 Zhang JF, Bioorg. Med. Chem. 2013; 23: 2538-2542; (c) Jung HS, Han JH, Habata Y,  
283 Kang C, Kim JS, Chem. Commun. 2011; 47: 5142-5144; (d) You QH, Lee AWM, Chan

- 284 WH, Zhu XM, Leung KCF, Chem. Commun. 2014; 50: 6207-6210; (e) Zhao CC, Chen  
285 Y, Zhang HY, Zhou BJ, Lv XJ, Fu WF, J. Photochem. Photobiol. A: Chem. 2014; 282:  
286 41-46; (f) Fu ZH, Yan LB, Zhang X, Zhu FF, Han XL, Fang J, Wang YW, Peng Y, Org.  
287 Biomol. Chem. 2017; 15: 4115-4121; (g) Lee KS, Park J, Park HJ, Chung YK, Park SB,  
288 Kim HJ, Shin IS, Hong JI, Sens. Actuators B Chem. 2016; 237: 256-261.
- 289 [22] Qiang HZ, Li SL, Ying GY, Jiang Y, Sens. Actuators B Chem. 2015; 212: 220-224.
- 290 [23] Yang C, Wang X, Liu H, Ge S, Yu J, Yan M, New J. Chem. 2017; 41: 3374-3379.
- 291 [24] You GR, Jang HJ, Jo TG, Kim C, RSC Adv. 2016; 6: 74400-74408.
- 292 [25] See for example: (a) El Sayed S, Licchelli M, Martínez-Máñez R, Sancenón F, Chem.  
293 As. J. 2017; 12: 2670-2674; (b) Ferreira RCM, Costa SPG, Gonçalves H, Raposo MMM,  
294 New J. Chem. 2017; 41: 12866-12878; (c) Marín-Hernández C, Santos-Figueroa LE, El  
295 Sayed S, Pardo T, Raposo MMM, Batista RMF, Costa SPG, Sancenón F, Martínez-  
296 Máñez R, Dyes Pigm. 2015; 122: 50-58; (d) Marín-Hernández C, Santos-Figueroa LE,  
297 Moragues ME, Raposo MMM, Batista RMF, Costa SPG, Pardo T, Martínez-Máñez R,  
298 Sancenón F, J. Org. Chem. 2014; 79: 10752-10761.
- 299 [26] See for example: (a) Maleki A, Paydar R, RSC Adv. 2015; 5: 33177-33184; (b) Kokare  
300 ND, Sangshetti JN, Shinde DB, Synthesis 2007; 2829-2834; (c) Shelke KF, Sapkal SB,  
301 Sonar SS, Madje BR, Shingate BB, Shingare MS, Bull. Kor. Chem. Soc. 2009; 30: 1057-  
302 1060; (d) Samai S, Nandi GC, Singh P, Singh MS, Tetrahedron 2009; 65: 10155-10161.
- 303 [27] Mandal S, Das G, Singh R, Shukla R, Bharadwaj PK, Coord. Chem. Rev. 1997; 160:  
304 191-235.
- 305 [28] Shi Y, Pan Y, Zhang H, Zhang Z, Li MJ, Yi C, Yang M, Biosensors Bioelectron. 2014;  
306 56: 39-45.

**Highlights**

- Detection of GSH in aqueous environments
- GSH-induced demetallation of Cu(II) complex yielded colour and emission changes
- Selective detection of GSH in the presence of cysteine and homocysteine

ACCEPTED MANUSCRIPT