

Document downloaded from:

<http://hdl.handle.net/10251/161701>

This paper must be cited as:

Jornet, M.; Calatayud, J.; Le Maître, OP.; Cortés, J. (2020). Variance reduction methods and multilevel Monte Carlo strategy for the density estimation of random second order linear differential equations solutions. *International Journal for Uncertainty Quantification*. 10(5):467-497. <https://doi.org/10.1615/Int.J.UncertaintyQuantification.2020032659>



The final publication is available at

<https://doi.org/10.1615/Int.J.UncertaintyQuantification.2020032659>

Copyright Begell House

Additional Information

VARIANCE REDUCTION METHODS AND MULTILEVEL MONTE CARLO STRATEGY FOR ESTIMATING DENSITIES OF SOLUTIONS TO RANDOM SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS

Marc Jornet,^{1,} Julia Calatayud,¹ Olivier P. Le Maître,² &
Juan Carlos Cortés¹*

¹*Instituto de Matemática Multidisciplinar, Universitat Politècnica de València, Camí de Vera s/n, 46022, Valencia, Spain*

²*CMAP, CNRS, INRIA, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France.*

*Address all correspondence to: Marc Jornet, Instituto de Matemática Multidisciplinar, Universitat Politècnica de València, Camí de Vera s/n, 46022, Valencia, Spain, E-mail: marjorsa@doctor.upv.es

Original Manuscript Submitted: mm/dd/yyyy; Final Draft Received: mm/dd/yyyy

This paper concerns the estimation of the density function of the solution to a random non-autonomous second-order linear differential equation with analytic data processes. In a recent contribution, we proposed to express the density function as an expectation, and we used a standard Monte Carlo algorithm to approximate the expectation. Although the algorithms worked satisfactorily for most test-problems, some numerical challenges emerged for others, due to large statistical errors. In these situations, the convergence of the Monte Carlo simulation slows down severely, and noisy features plague the estimates. In this paper, we focus on computational aspects and propose several variance reduction methods to remedy these issues and speed up the convergence. First, we introduce a path-wise selection of the approximating processes which aims at controlling the variance of the estimator. Second, we propose a hybrid method, combining Monte Carlo and deterministic quadrature rules, to estimate the expectation. Third, we exploit the series expansions of the solutions to design a multilevel Monte Carlo estimator. The proposed methods are implemented and tested on several numerical examples to highlight the theoretical discussions and demonstrate the significant improvements achieved.

KEY WORDS: *random linear differential equation; probability density function; standard and multilevel Monte Carlo simulation; analysis of algorithms*

1. INTRODUCTION AND PRELIMINARIES

This paper concerns the analysis of stochastic processes $X(t)$ solution to general random non-autonomous second order linear differential equations:

$$\begin{cases} \ddot{X}(t) + A(t)\dot{X}(t) + B(t)X(t) = 0, \\ X(t_0) = Y_0, \dot{X}(t_0) = Y_1. \end{cases} \quad (1)$$

The time-dependent coefficients $A(t)$ and $B(t)$ are stochastic processes and the initial conditions Y_0 and Y_1 are random variables. These random quantities are defined in an underlying complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where Ω is the sample space formed by outcomes $\omega \in \Omega$, $\mathcal{F} \subseteq 2^\Omega$ is the σ -algebra of events, and \mathbb{P} is the probability measure. Problem (1) belongs to a broader class of equations referred to as random differential equations [1–3]. Random differential equations are those in which random effects are manifested through input coefficients, initial condition and/or forcing term, in the form of random variables or regular stochastic processes with any type of probability distribution; the solution stochastic process is differentiable (not to be confused with stochastic differential equations of Itô type driven by irregular processes, whose solutions exhibit nondifferentiable sample paths; see [4, pp. 96–98] for the distinction between random and stochastic differential equations). Random differential equations are generally studied with two distinct approaches, based on random calculus and sample paths. For the necessary theory on random calculus, specifically mean square calculus with the norm $\|Z\|_2 = \mathbb{E}[Z^2]^{1/2}$ for random variables Z in the Lebesgue space $L^2(\Omega)$, we refer the reader to [1].

In this paper, we assume that $A(t)$ and $B(t)$ are analytic stochastic processes in the mean square sense [1, p. 99], so they can be expanded as

$$A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n, \quad B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n, \quad t \in (t_0 - r, t_0 + r), \quad (2)$$

where $A_0, A_1, \dots, B_0, B_1, \dots$ are second order random variables, which may or may not be statistically independent, given by $A_i = A^{(i)}(t_0)/i!$ and $B_i = B^{(i)}(t_0)/i!$, with the i -th derivative considered in the mean square sense. The Taylor series of $A(t)$ and $B(t)$ in (2) present mean square convergence. It is interesting to observe that the analytic property in the mean sense of a stochastic process is determined by the ordinary analytic property of its correlation function [1, Th. 4.4.3]. The regularity of $A(t)$ and $B(t)$ does not allow them to be Brownian motions or white noise processes, thus the distinct treatment of our problem compared to Itô calculus.

1 The linear problem (1) with analytic data processes has raised interest in the literature on random differential
 2 equations. The study was initialized in particular equations from Mathematical Physics, such as Airy, Hermite and
 3 Legendre differential equations [5–7]. The main goal of these contributions was to construct the mean square analytic
 4 solution

$$X(t) = \sum_{n=0}^{\infty} X_n(t - t_0)^n \quad (3)$$

5 on $(t_0 - r, t_0 + r)$ via the Fröbenius method, thus obtaining approximations of the expectation and the variance of
 6 $X(t)$, $\mathbb{E}[X(t)]$ and $\mathbb{V}[X(t)]$. Other techniques to study particular cases of (1) are PC expansions [8,9], homotopy
 7 method [10], variational iteration [11], Adomian decomposition [12], and differential transform method [13,14].

8 The general problem (1) was successfully solved in [15,16] using the Fröbenius method. It was proved that, if
 9 the power series from (2) converge in $L^\infty(\Omega)$ (with the essential supremum norm $\|\cdot\|_\infty$) and if $Y_0, Y_1 \in L^2(\Omega)$, then
 10 there is a unique mean square solution of the form (3). This result is extendable to a more general measure norm,
 11 in the Lebesgue space $(L^p(\Omega), \|\cdot\|_p)$, $1 \leq p \leq \infty$: if $Y_0, Y_1 \in L^p(\Omega)$, then the series (3) converges in $L^p(\Omega)$ and
 12 solves (1) in the $L^p(\Omega)$ sense. The convergence is exponentially fast, although not uniformly in t ; it deteriorates when
 13 t or the norms of the input coefficients grow.

14 Throughout the paper, we assume that the power series from (2) converge in $L^\infty(\Omega)$ (with $A_0, A_1, \dots, B_0, B_1, \dots$
 15 bounded) and that $Y_0, Y_1 \in L^2(\Omega)$, so that the conditions stated above for the existence of mean square stochastic
 16 solution are fulfilled.

17 The computation of the probability density function of $X(t)$ is a more ambitious goal for quantifying uncer-
 18 tainty. When $X(t)$ is absolutely continuous, meaning that its probability law is absolutely continuous with respect to
 19 the Lebesgue measure, then there exists a non-negative Borel measurable function $f_{X(t)}(x) = \frac{d(\mathbb{P} \circ X(t)^{-1})(x)}{dx}$ char-
 20 acterized by $\mathbb{P}[X(t) \in \mathcal{C}] = \int_{\mathcal{C}} f_{X(t)}(x) dx$ ($\mathcal{C} \subseteq \mathbb{R}$ is any Borel set). Confidence intervals and statistical moments
 21 of the response $X(t)$ can be computed through the probability density function.

22 In random differential equation problems, the probability density function is usually computed through trans-
 23 formation of random variables [17–23]. This approach was used in [24] for the autonomous version of (1), when
 24 $A(t) = A$ and $B(t) = B$ are time-independent random variables. We extended the computation of the density
 25 function to the general problem (1) in [25].

26 The procedure presented in [25] to derive the probability density function $f_{X(t)}(x)$ is briefly summarized. Let
 27 $\{S_0(t), S_1(t)\}$ be the fundamental set of solutions to (1) satisfying the deterministic initial conditions $S_0(t_0) = 1$,
 28 $\dot{S}_0(t_0) = 0$, $S_1(t_0) = 0$ and $\dot{S}_1(t_0) = 1$. Both $S_0(t)$ and $S_1(t)$ are given by infinite random power series, converging

1 in $L^\infty(\Omega)$. The solution $X(t)$ is expressed as $X(t) = Y_0 S_0(t) + Y_1 S_1(t)$. If $S_0(t) \neq 0$ almost surely and Y_0 has a
2 density f_{Y_0} , we obtain

$$f_{X(t)}(x) = \mathbb{E}[Z_0(x, t)], \quad (4)$$

3 where

$$Z_0(x, t) = f_{Y_0} \left(\frac{x - Y_1 S_1(t)}{S_0(t)} \right) \frac{1}{|S_0(t)|}. \quad (5)$$

4 Similarly, if $S_1(t) \neq 0$ almost surely and Y_1 possesses a density f_{Y_1} , then

$$f_{X(t)}(x) = \mathbb{E}[Z_1(x, t)], \quad (6)$$

5 where

$$Z_1(x, t) = f_{Y_1} \left(\frac{x - Y_0 S_0(t)}{S_1(t)} \right) \frac{1}{|S_1(t)|}. \quad (7)$$

6 When a simple closed-form expression for $S_0(t)$ and $S_1(t)$ is not available, truncating the power series $S_0(t)$ and $S_1(t)$
7 to the $N \geq 0$ first terms is necessary. We denote $S_0^N(t)$ and $S_1^N(t)$, the partial sums which converge respectively to
8 $S_0(t)$ and $S_1(t)$, in $L^\infty(\Omega)$, as $N \rightarrow \infty$. The truncated series of $X(t)$ is given by $X^N(t) = Y_0 S_0^N(t) + Y_1 S_1^N(t)$,
9 which converges to $X(t)$ as $N \rightarrow \infty$ in the mean square sense. The probability density function of $X^N(t)$ is,
10 whenever $S_0^N(t) \neq 0$ almost surely or $S_1^N(t) \neq 0$ almost surely,

$$f_{X^N(t)}(x) = \mathbb{E}[Z_0^N(x, t)] \quad (8)$$

11 or

$$f_{X^N(t)}(x) = \mathbb{E}[Z_1^N(x, t)], \quad (9)$$

12 where

$$Z_0^N(x, t) = f_{Y_0} \left(\frac{x - Y_1 S_1^N(t)}{S_0^N(t)} \right) \frac{1}{|S_0^N(t)|}, \quad Z_1^N(x, t) = f_{Y_1} \left(\frac{x - Y_0 S_0^N(t)}{S_1^N(t)} \right) \frac{1}{|S_1^N(t)|}. \quad (10)$$

13 Several theoretical results from [25] justify that $f_{X^N(t)}(x)$ tends to the target density function $f_{X(t)}(x)$ as $N \rightarrow \infty$ in
14 a neighborhood of t_0 . The convergence is pointwise and in $L^p(\mathbb{R})$, for $1 \leq p < \infty$. Convergence in terms of the total
15 variation and the Hellinger distances also holds. The hypotheses set to establish the convergence are mainly related
16 to the continuity of f_{Y_0} or f_{Y_1} . If f_{Y_0} or f_{Y_1} are Hölder continuous, then exponential convergence holds, albeit not
17 uniform in t and x .

1 In [25], the expectation from (8) and (9) is approximated using Monte Carlo (MC) simulation, via an explicit
 2 algorithm. Either (8) or (9) are selected, and realizations of the involved random variables are generated to compute the
 3 sample average. This introduces an error due to sampling, apart from the bias error $\theta_N(x, t) = f_{X(t)}(x) - f_{X^N(t)}(x)$.
 4 The convergence rate of the MC procedure towards $f_{X^N(t)}(x)$ depends on the finite number of realizations and on
 5 the variances $\mathbb{V}[Z_0^N(x, t)]$, $\mathbb{V}[Z_1^N(x, t)]$. If these variances are large or infinite, the convergence rate of the MC
 6 simulation deteriorates severely and noisy estimates of $f_{X^N(t)}(x)$ are produced, thus invalidating the results. This
 7 phenomenon was observed in the numerical experiments from [25, Example 5.3]. It is highly related to having small
 8 denominators $|S_0^N(t)|$ and $|S_1^N(t)|$ at certain realizable paths, as this may produce higher dispersion of $1/|S_0^N(t)|$ and
 9 $1/|S_1^N(t)|$.

10 The main goal of this paper is to improve the algorithm from [25] via several variance reduction methods to
 11 achieve good convergence of the MC simulation and prevent noisy features. All the proposed methods are imple-
 12 mented and tested using the software Mathematica[®], version 11.2 [26].

13 In Section 2, we propose a path-wise selection of the initial condition to avoid the smallest denominator, $|S_0^N(t)|$
 14 or $|S_1^N(t)|$, used in (4)–(6). In this manner, we control the variance of the sampling error, and we improve the quality
 15 of the numerical results significantly. An important assumption to apply this selection is that the two initial conditions
 16 have probability density functions.

17 In Section 3, we improve further the algorithm of Section 2 with a deterministic quadrature to approximate the
 18 expectation with respect to Y_0 – Y_1 . It allows for a reduction of variance when Y_0 and Y_1 are responsible for most of
 19 the estimator variability (this can be checked using Sobol indices). A detailed comparison with the methodology of
 20 Section 2, in terms of variance reduction, density approximation, and error versus complexity, is carried out.

21 In Section 4, the MC approach from Section 2 is extended considering a multilevel strategy. Originally introduced
 22 in [27,28], the idea of multilevel MC is to accelerate the convergence of the MC simulation by decomposing the
 23 target expectation into the sum of expectations of increments (through a telescopic sum identity) whose variances
 24 decay rapidly. Multilevel MC simulation is an improved sampling strategy that balances the bias and sampling errors,
 25 to obtain an overall reduction of the computational complexity compared to the standard MC algorithm. Multilevel
 26 sampling strategies have been employed for the approximation of statistics in stochastic differential equations and
 27 random partial differential equations, where the bias classically arises from a mesh discretization [29–31]. However,
 28 to our knowledge, the application to the density estimation of random differential equations solution is completely
 29 original.

30 Finally, Section 5 draws the main conclusions of the paper and discusses potential avenues for future research on

1 algorithmic improvements, linear random differential equations, and density approximations.

2 The proofs of the mathematical results (propositions and lemmas) are postponed to the Appendix.

3 2. VARIANCE REDUCTION BY PATH-WISE SELECTION OF THE INITIAL CONDITION USED IN THE 4 DENSITY EXPRESSION

5 In [25], the density function $f_{X^N(t)}(x)$ is approximated using a standard MC sampling strategy, by initially select-
6 ing either $Z_0^N(x, t)$ or $Z_1^N(x, t)$ (see (10)), and only one of these processes. The convergence rate of the sample
7 average towards $\mathbb{E}[Z_0^N(x, t)]$ or $\mathbb{E}[Z_1^N(x, t)]$ depends on $\mathbb{V}[Z_0^N(x, t)]$ or $\mathbb{V}[Z_1^N(x, t)]$, respectively. If $\mathbb{V}[Z_0^N(x, t)]$
8 and $\mathbb{V}[Z_1^N(x, t)]$ are large or infinity, the convergence rate deteriorates severely and noisy features may plague the
9 statistical approximation to $f_{X^N(t)}(x)$. This issue was observed in [25, Example 5.3].

10 In this section we propose a variance reduction method by combining the choices of $Z_0^N(x, t)$ and $Z_1^N(x, t)$.
11 Consider one realization of $S_0^N(t)$ and $S_1^N(t)$. These realizations are computed recursively in the computer from
12 realizations of $A_0, \dots, A_{N-2}, B_0, \dots, B_{N-2}$, see [25]. If $|S_0^N(t)| \geq |S_1^N(t)|$ then we pick $Z_0^N(x, t)$, and vice versa
13 otherwise. In such a way, we avoid the smallest denominator in the expressions of $Z_0^N(x, t)$ and $Z_1^N(x, t)$.

14 We assume that Y_0, Y_1 and $(A_0, A_1, \dots, B_0, B_1, \dots)$ are independent. We also suppose that Y_0 and Y_1 are ab-
15 solutely continuous random variables (otherwise this method is no longer applicable). Given the events $G_0(t) =$
16 $\{|S_0(t)| \geq |S_1(t)|\}$, $G_1(t) = \{|S_1(t)| > |S_0(t)|\}$, $G_0^N(t) = \{|S_0^N(t)| \geq |S_1^N(t)|\}$ and $G_1^N(t) = \{|S_1^N(t)| >$
17 $|S_0^N(t)|\}$, we define the random processes

$$Z(x, t) = Z_0(x, t)1_{G_0(t)} + Z_1(x, t)1_{G_1(t)}, \quad (11)$$

18

$$Z^N(x, t) = Z_0^N(x, t)1_{G_0^N(t)} + Z_1^N(x, t)1_{G_1^N(t)}. \quad (12)$$

19 These processes correspond to path-wise selecting f_{Y_0} or f_{Y_1} according to $|S_0(t)| \geq |S_1(t)|$ or $|S_1(t)| > |S_0(t)|$
20 (in (11)), and $|S_0^N(t)| \geq |S_1^N(t)|$ or $|S_1^N(t)| > |S_0^N(t)|$ (in (12)). By path-wise selection, we refer to generate real-
21 izations of $S_0^N(t)$ and $S_1^N(t)$ (the paths/trajectories of the stochastic processes) and then select $Z_0^N(x, t)$ or $Z_1^N(x, t)$
22 accordingly. The density functions of $X^N(t)$ and $X(t)$ are then expressed as follows.

23 **Proposition 1.** *The following relationships hold:*

$$f_{X(t)}(x) = \mathbb{E}[Z(x, t)], \quad f_{X^N(t)}(x) = \mathbb{E}[Z^N(x, t)],$$

1 where $Z(x, t)$ and $Z^N(x, t)$ are defined in (11) and (12).

2 The expectation $f_{X^N(t)}(x) = \mathbb{E}[Z^N(x, t)]$ may be approximated via a standard MC sampling strategy. A key
3 feature here is the uniform boundedness of $\mathbb{V}[Z^N(x, t)]$ with N , as shown in Proposition 3.

4 **Lemma 2.** *Given any $T > t_0$, there exists a constant $\alpha > 0$ such that*

$$\max\{|S_0(t)|, |S_1(t)|\} \geq 2\alpha \quad (13)$$

5 holds almost surely, for all $t \in [t_0, T]$. In particular, there exists an integer $N_{T, \alpha} \geq 0$ such that

$$\max\{|S_0^N(t)|, |S_1^N(t)|\} \geq \alpha \quad (14)$$

6 holds almost surely, for all $t \in [t_0, T]$ and $N \geq N_{T, \alpha}$.

7 **Proposition 3.** *Suppose that f_{Y_0} and f_{Y_1} are bounded density functions. Given any $T > t_0$, the $L^2(\Omega)$ norms of
8 $Z(x, t)$ and $Z^N(x, t)$ are controlled as follows:*

$$\|Z(x, t)\|_2 \leq \frac{\max\{\|f_{Y_0}\|_\infty, \|f_{Y_1}\|_\infty\}}{2\alpha}, \quad \|Z^N(x, t)\|_2 \leq \frac{\max\{\|f_{Y_0}\|_\infty, \|f_{Y_1}\|_\infty\}}{\alpha},$$

9 for $x \in \mathbb{R}$, $t \in [t_0, T]$ and $N \geq N_{T, \alpha}$, where α is the constant from Lemma 2.

10 As a consequence, the variance here is well controlled, in contrast with [25]. The estimate of $f_{X^N(t)}(x) =$
11 $\mathbb{E}[Z^N(x, t)]$ using MC simulation does not present convergence problems.

12 **Remark 1.** *Lemma 2 is not only important from a numerical point of view, but also for the following theoretical fact.
13 Our paper [25] justified the pointwise convergence of $\{f_{X^N(t)}(x)\}_{N=0}^\infty$ towards $f_{X(t)}(x)$ in a neighborhood of t_0 .
14 The neighborhood was constructed in such a way that $S_0(t)$ and/or $S_1(t)$ are greater than a positive constant almost
15 surely (see [25, Remarks 3.5, 3.10]), so that the denominators from (10) are controlled. Lemma 2 shows that*

$$\frac{1}{|S_0(t)|} \mathbf{1}_{G_0(t)} + \frac{1}{|S_1(t)|} \mathbf{1}_{G_1(t)} = \frac{1}{\max\{|S_0(t)|, |S_1(t)|\}}$$

16 is upper bounded by a finite constant almost surely, for every t in the domain of definition of (1). Hence, the theoretical
17 results from [25] justifying the convergence of $f_{X^N(t)}(x) = \mathbb{E}[Z^N(x, t)]$ to $f_{X(t)}(x) = \mathbb{E}[Z(x, t)]$ as $N \rightarrow \infty$ do
18 hold for every t where $X(t)$ is well-defined, not just on a neighborhood of t_0 . Notice that this statement requires both

1 Y_0 and Y_1 to be absolutely continuous, and Y_0, Y_1 and $(A_0, A_1, \dots, B_0, B_1, \dots)$ to be independent. These are the
2 hypotheses under which the methodology from this section is supported.

3 Given a time variable t , a point x in the density domain of $X^N(t)$, a truncation order N and a number M , the
4 whole procedure described in this section is structured as follows:

5 **PROCEDURE 1.**

6 **Part 1** Generate M realizations of $S_0^N(t)$ and $S_1^N(t)$.

7 **Part 2** For each realization, if $|S_0^N(t)| \geq |S_1^N(t)|$, set a realization of $Y_1 \sim f_{Y_1}$ to obtain the realization of $Z^N(x, t)$
8 given by $Z_0^N(x, t)$ (see (10), (12)). Otherwise set a realization of $Y_0 \sim f_{Y_0}$ to obtain the realization of $Z^N(x, t)$
9 given by $Z_1^N(x, t)$.

10 **Part 3** Compute the sample mean of the M realizations of $Z^N(x, t)$. This value estimates the expectation $f_{X^N(t)}(x) =$
11 $\mathbb{E}[Z^N(x, t)]$ with MC simulation. We denote the estimate (output function) by $f_X^{N,M}(x, t)$, which tends to
12 $f_{X^N(t)}(x)$ as $M \rightarrow \infty$.

13 The complexity (or cost, or work, which is defined in this paper as the total number of arithmetic operations) of
14 Procedure 1 is $\mathcal{O}(MN^2)$, in general. Recall that each realization of $S_0^N(t)$ and $S_1^N(t)$ is computed recursively with
15 N and requires $\mathcal{O}(N^2)$ operations [25].

16 As discussed in [25], when $A(t)$ and $B(t)$ are random polynomials instead of infinite random series, the cost
17 per realization of $S_0^N(t)$ and $S_1^N(t)$ is reduced to $\mathcal{O}(N)$ operations. Then the global complexity of Procedure 1 is
18 $\mathcal{O}(MN)$ only.

19 Obviously, in the case where $S_0(t)$ and $S_1(t)$ are known in closed-form expression (this occurs for simple prob-
20 lems (1), such as the autonomous case), Procedure 1 can be run with $N = \infty$, that is, by computing realizations of
21 $S_0(t)$ and $S_1(t)$ directly. The complexity of the algorithm is $\mathcal{O}(M)$ in such a case. The error is statistical and only
22 due to the finite sampling.

23 Procedure 1 may be run symbolically on the variable x , thus having a symbolic expression for the output
24 $f_X^{N,M}(x, t)$ in the end. The algorithm from [25] was run symbolically. However, symbolic computations do not
25 allow a large number of simulations M . Thus, in cases where M needs to be large enough to achieve convergence,
26 we recommend to discretize x in a domain of interest for the target density.

27 In the following two examples, we illustrate and assess the theoretical discussion from this section. The main
28 goal is to highlight the improvement of Procedure 1 compared with [25], in terms of variance reduction of the MC

1 estimator and accuracy of the density estimation. The first example, Example 1, focuses on a simple autonomous
 2 problem with exact solution, so that $N = \infty$. The second example, Example 2, deals with a more complex problem,
 3 where $A(t)$ and $B(t)$ are one-degree random polynomials and the series solution is truncated to a finite number N of
 4 terms.

5 **Example 1.** We start by considering a very simple problem (1):

$$\begin{cases} \ddot{X}(t) + B^2 X(t) = 0, & t \in \mathbb{R}, \\ X(t_0 = 0) = Y_0, \quad \dot{X}(t_0 = 0) = Y_1. \end{cases} \quad (15)$$

6 Despite its simplicity, (15) is a useful example to illustrate the theoretical discussion from this section.

7 The following probability distributions are considered: $B \sim \text{Uniform}(1, 2.5)$, $Y_0, Y_1 \sim \text{Normal}(1, \sigma = 0.1)$,
 8 all of them independent. As B is bounded and the initial conditions Y_0 and Y_1 have absolute moments of any order,
 9 there exists an analytic solution $X(t)$ in the $L^p(\Omega)$ sense, for $1 \leq p < \infty$. The fundamental system is explicitly
 10 known: $S_0(t) = \cos(Bt)$, $S_1(t) = \sin(Bt)/B$, $t \in \mathbb{R}$. Thus, Procedure 1 can be applied with $N = \infty$, by taking
 11 realizations from $S_0(t)$ and $S_1(t)$ directly. We consider time $t = 10$, and the goal is to approximate the density
 12 function $f_{X(t=10)}(x)$ using MC methods.

13 In Figure 1, left panel, we estimate $\mathbb{E}[Z_0(x, t = 10)]$ and $\mathbb{E}[Z_1(x, t = 10)]$ with the MC procedure from [25], by
 14 generating $M = 100,000$ realizations of the random inputs. These expectations correspond to $f_{X(t=10)}(x)$. Observe
 15 that noisy features plague the estimates of the density function. In Figure 1, right panel, we plot estimates of the vari-
 16 ances $\mathbb{V}[Z_0(x, t = 10)]$ and $\mathbb{V}[Z_1(x, t = 10)]$. Their large values slow down the convergence of the MC simulation,
 17 thus explaining the noisy behavior of the estimates in the left plot. In fact, at the points x where the variance is larger,
 18 more noise is perceived in the estimate of the expectation.

19 In Figure 2, left panel, we approximate $f_{X(t=10)}(x)$ via $\mathbb{E}[Z(x, t = 10)]$, by using the proposed Procedure 1
 20 with $M = 100,000$ realizations. Observe that now the estimate is smooth, thus resolving the convergence challenge.
 21 In Figure 2, right panel, we depict an estimate for $\mathbb{V}[Z(x, t = 10)]$. This variance is shown to be small compared
 22 to $\mathbb{V}[Z_0(x, t = 10)]$ and $\mathbb{V}[Z_1(x, t = 10)]$ in Figure 1, thus allowing for faster convergence of the MC simulation.
 23 Hence, Procedure 1 improves [25] significantly.

24 It is interesting to observe the different behaviors of $Z_0(x, 10)$, $Z_1(x, 10)$ and $Z(x, 10) = Z_0(x, 10)1_{G_0(10)} +$
 25 $Z_1(x, 10)1_{G_1(10)}$ as real functions of B , Y_0 and Y_1 . Figure 3 depicts the 3D graphs of the real functions $Z_0(0, 10)$,
 26 $Z_1(0, 10)$, $Z_0(0, 10)1_{G_0(10)}$ and $Z_1(0, 10)1_{G_1(10)}$ (obviously, any other point x different from 0 can also be taken). The




FIG. 1: Left: approximations of $f_{X(t=10)}(x)$ using estimates $\hat{\mathbb{E}}[Z_0(x, 10)]$ and $\hat{\mathbb{E}}[Z_1(x, 10)]$. Right: estimates $\hat{V}[Z_0(x, 10)]$ and $\hat{V}[Z_1(x, 10)]$. This figure corresponds to Example 1.




FIG. 2: Left: approximation of $f_{X(t=10)}(x)$ using estimate $\hat{\mathbb{E}}[Z(x, 10)]$. Right: estimate $\hat{V}[Z(x, 10)]$. This figure corresponds to Example 1.

- 1 functions $Z_0(0, 10)$ and $Z_1(0, 10)$ have singularities (because of vanishing denominators $|\cos(10B)|$ and $|\sin(10B)/B|$)
 2 and are strongly peaked. On the contrary, although $Z_0(0, 10)1_{G_0(10)}$ and $Z_1(0, 10)1_{G_1(10)}$ possess discontinuity points,
 3 their peaks are not that sharp. As we are getting rid of small denominators, the sharp peaks are avoided.




FIG. 3: 3D plots of the integrands $Z_0(0, 10)$, $Z_1(0, 10)$ (their vertical axes have been restricted to $[0, 50]$), $Z_0(0, 10)1_{G_0(10)}$ and $Z_1(0, 10)1_{G_1(10)}$. This figure corresponds to Example 1.

- 4 **Example 2.** We consider a more complex example now, based on degree one polynomial coefficients. Let

$$\begin{cases} \ddot{X}(t) + (A_0 + A_1 t)\dot{X}(t) + (B_0 + B_1 t)X(t) = 0, & t \in \mathbb{R}, \\ X(t_0 = 0) = Y_0, \quad \dot{X}(t_0 = 0) = Y_1, \end{cases} \quad (16)$$

- 5 where $A_0 = 4$, $A_1 \sim \text{Uniform}(0, 1)$, $B_0 \sim \text{Gamma}(2, 2)|_{[0, 4]}$, $B_1 \sim \text{Bernoulli}(0.35)$ and $Y_0, Y_1 \sim \text{Normal}(2, 1)$, all
 6 of them independent. Since the random coefficients A_0, A_1, B_0 and B_1 are bounded and the initial conditions Y_0 and

1 Y_1 have finite absolute moments, there is an analytic solution $X(t)$ in the $L^p(\Omega)$ sense, $1 \leq p < \infty$.

2 According to [25], the probability density function $f_{X(t)}(x)$ can be approximated pointwise using $f_{X^N(t)}(x)$
 3 (with exponential convergence with N because f_{Y_0} and f_{Y_1} are Lipschitz continuous on \mathbb{R}). We focus on the order of
 4 truncation $N = 10$ to highlight the improvement of Procedure 1 compared with [25]. We estimate $f_{X^{N=10}(t=1.5)}(x)$
 5 with both approaches.

6 In Figure 4, left panel, we estimate $\mathbb{E}[Z_0^{N=10}(x, t = 1.5)]$ and $\mathbb{E}[Z_1^{N=10}(x, t = 1.5)]$ with the algorithm described
 7 in [25]. With $M = 20,000$ realizations, we observe noise in both estimates. This phenomenon is accurately explained
 8 by the plots of $\mathbb{V}[Z_0^{N=10}(x, t = 1.5)]$ and $\mathbb{V}[Z_1^{N=10}(x, t = 1.5)]$ in the right panel. It is very interesting to observe
 9 that the points x where the variances are larger correspond to higher noise in the left plot. Indeed, large variance
 10 affects the rate of convergence of the MC simulation severely.




FIG. 4: Left: approximations of $f_{X^{N=10}(t=1.5)}(x)$ using estimates $\hat{\mathbb{E}}[Z_0^{N=10}(x, 1.5)]$ and $\hat{\mathbb{E}}[Z_1^{N=10}(x, 1.5)]$. Right: estimates $\hat{\mathbb{V}}[Z_0^{N=10}(x, 1.5)]$ and $\hat{\mathbb{V}}[Z_1^{N=10}(x, 1.5)]$. This figure corresponds to Example 2.

11 Procedure 1 has been designed to fix this issue. Figure 5, left panel, shows the smooth estimate of $f_{X^{N=10}(t=1.5)}(x)$
 12 with $\mathbb{E}[Z^{N=10}(x, t = 1.5)]$ and $M = 20,000$ realizations. This good convergence is due to the small variance
 13 $\mathbb{V}[Z^{N=10}(x, t = 1.5)]$ compared to $\mathbb{V}[Z_0^{N=10}(x, t = 1.5)]$ and $\mathbb{V}[Z_1^{N=10}(x, t = 1.5)]$, see the right panel of Figure 5.
 14 Hence, Procedure 1 shows significantly superior to a mere MC approach as in [25].

15 **3. VARIANCE REDUCTION BY USING QUADRATURE RULE FOR THE EXPECTATION WITH**
 16 **RESPECT TO THE INITIAL CONDITIONS**

17 By Proposition 1, the probability density function of the truncated series $X^N(t)$ is expressed as $f_{X^N(t)}(x) =$
 18 $\mathbb{E}[Z^N(x, t)]$, where $Z^N(x, t)$ was defined in (12). The randomness in $Z^N(x, t)$ comes from the initial conditions,
 19 Y_0 and Y_1 , and from the truncated series from the fundamental set, $S_0^N(t)$ and $S_1^N(t)$. An MC sampling procedure




FIG. 5: Left: approximation of $f_{X^{N=10}(1.5)}(x)$ using estimate $\hat{\mathbb{E}}[Z^{N=10}(x, 1.5)]$. Right: estimate $\hat{\mathbb{V}}[Z^{N=10}(x, 1.5)]$. This figure corresponds to Example 2.

1 generates M independent realizations from these random quantities and approximates $f_{X^{N(t)}}(x)$ using the sample
 2 average, see Procedure 1.

3 The convergence rate of the sample mean towards the true expectation depends heavily on the variance of
 4 $Z^N(x, t)$ (which is uniformly bounded on N by Proposition 3). In this section, we present a method for variance
 5 reduction. Each time we generate a realization of $S_0^N(t)$ and $S_1^N(t)$ and check whether $|S_0^N(t)| \geq |S_1^N(t)|$ or
 6 $|S_1^N(t)| > |S_0^N(t)|$, now we do not generate a realization of $Y_0 - Y_1$, but compute the expectation with respect to
 7 $Y_0 - Y_1$ with a numerical integration method, instead.

8 We write $f_{X^{N(t)}}(x)$ as $f_{X^{N(t)}}(x) = \mathbb{E}[\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]]$, by the law of total expectation. By an MC
 9 sampling procedure, we approximate the outer expectation as

$$f_{X^{N(t)}}(x) \approx \frac{1}{M} \sum_{i=1}^M \mathbb{E}[Z^N(x, t)|S_0^N(t) = s_{0,i}^N, S_1^N(t) = s_{1,i}^N], \quad (17)$$

10 where $\{(s_{0,i}^N, s_{1,i}^N)\}_{i=1}^M$ are M independent realizations of $(S_0^N(t), S_1^N(t))$.

The variance of the sampling error from the approximation (17) is always smaller than in the standard MC
 approach from Procedure 1. Indeed, by the law of total variance,

$$\begin{aligned} \mathbb{V}[\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]] &= \mathbb{V}[Z^N(x, t)] - \mathbb{E}[\mathbb{V}[Z^N(x, t)|S_0^N(t), S_1^N(t)]] \\ &\leq \mathbb{V}[Z^N(x, t)]. \end{aligned} \quad (18)$$

11 Equality holds only when $\mathbb{V}[Z^N(x, t)|S_0^N(t), S_1^N(t)] = 0$. Essentially, using (17) we gain more reduction in the

1 sampling error when Y_0 - Y_1 produce high variability in $Z^N(x, t)$. The extreme case occurs when $A(t)$ and $B(t)$
 2 are deterministic functions, as in such a case a deterministic integration rule for Y_0 - Y_1 makes the method sampling
 3 error-free.

4 The influence of the variability of Y_0 , Y_1 , $S_0^N(t)$ and $S_1^N(t)$ on $Z^N(x, t)$ (variance-based sensitivity analysis)
 5 can be assessed using Sobol indices [32,33]. The random process $Z^N = Z^N(x, t)$ has the following orthogonal
 6 decomposition:

$$Z^N = \mathbb{E}[Z^N] + \tilde{Z}_{S_0^N(t), S_1^N(t)}^N + \tilde{Z}_{Y_0, Y_1}^N + \tilde{Z}_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N,$$

7 where

$$\tilde{Z}_{S_0^N(t), S_1^N(t)}^N = Z_{S_0^N(t), S_1^N(t)}^N - \mathbb{E}[Z_{S_0^N(t), S_1^N(t)}^N], \quad Z_{S_0^N(t), S_1^N(t)}^N = \mathbb{E}[Z^N | S_0^N(t), S_1^N(t)],$$

8

$$\tilde{Z}_{Y_0, Y_1}^N = Z_{Y_0, Y_1}^N - \mathbb{E}[Z_{Y_0, Y_1}^N], \quad Z_{Y_0, Y_1}^N = \mathbb{E}[Z^N | Y_0, Y_1],$$

9

$$\tilde{Z}_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N = Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N - \mathbb{E}[Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N],$$

10

$$Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N = Z^N - \tilde{Z}_{S_0^N(t), S_1^N(t)}^N - \tilde{Z}_{Y_0, Y_1}^N.$$

11 Applying squared $L^2(\Omega)$ norms, we derive the ANOVA-Hoeffding decomposition of the variance of $Z^N(x, t)$:

$$\mathbb{V}[Z^N] = \mathbb{V}[Z_{S_0^N(t), S_1^N(t)}^N] + \mathbb{V}[Z_{Y_0, Y_1}^N] + \mathbb{V}[Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N],$$

12 where $\mathbb{V}[Z_{S_0^N(t), S_1^N(t)}^N]$, $\mathbb{V}[Z_{Y_0, Y_1}^N]$ and $\mathbb{V}[Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N]$ measure the contribution to $Z^N(x, t)$ of $(S_0^N(t), S_1^N(t))$
 13 alone, (Y_0, Y_1) alone, and the interactions between them, respectively. The standardization of the variance decompo-
 14 sition gives rise to the Sobol indices:

$$\mathbb{S}_{S_0^N(t), S_1^N(t)}^N = \frac{\mathbb{V}[Z_{S_0^N(t), S_1^N(t)}^N]}{\mathbb{V}[Z^N]}, \quad \mathbb{S}_{Y_0, Y_1}^N = \frac{\mathbb{V}[Z_{Y_0, Y_1}^N]}{\mathbb{V}[Z^N]}, \quad (19)$$

15

$$\mathbb{S}_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N = \frac{\mathbb{V}[Z_{S_0^N(t), S_1^N(t), Y_0, Y_1}^N]}{\mathbb{V}[Z^N]}, \quad (20)$$

16 which are scaled in $[0, 1]$. When \mathbb{S}_{Y_0, Y_1}^N is near 1, we expect more variance reduction in the MC estimate (17).

Fix the index of the realizations $i = 1, \dots, M$. If $|s_{0,i}^N| \geq |s_{1,i}^N|$, then

$$\begin{aligned} \mathbb{E} [Z^N(x, t) | S_0^N(t) = s_{0,i}^N, S_1^N(t) = s_{1,i}^N] &= \mathbb{E} [Z_0^N(x, t) | S_0^N(t) = s_{0,i}^N, S_1^N(t) = s_{1,i}^N] \\ &= \frac{1}{|s_{0,i}^N|} \int_{\mathcal{S}_{Y_0}} f_{Y_0} \left(\frac{x - y_1 s_{1,i}^N}{s_{0,i}^N} \right) f_{Y_1}(y_1) dy_1, \end{aligned} \quad (21)$$

where \mathcal{S}_{Y_1} denotes the support of the random variable Y_1 . If $|s_{1,i}^N| > |s_{0,i}^N|$, then

$$\begin{aligned} \mathbb{E} [Z^N(x, t) | S_0^N(t) = s_{0,i}^N, S_1^N(t) = s_{1,i}^N] &= \mathbb{E} [Z_1^N(x, t) | S_0^N(t) = s_{0,i}^N, S_1^N(t) = s_{1,i}^N] \\ &= \frac{1}{|s_{1,i}^N|} \int_{\mathcal{S}_{Y_0}} f_{Y_1} \left(\frac{x - y_0 s_{0,i}^N}{s_{1,i}^N} \right) f_{Y_0}(y_0) dy_0, \end{aligned} \quad (22)$$

1 being \mathcal{S}_{Y_0} the support of Y_0 . These integrals (21), (22), can be approximated using numerical integration schemes.
 2 For instance, Gaussian quadrature rules [9, pp. 39–41] are appropriate since the density functions $f_{Y_1}(y_1)$ and $f_{Y_0}(y_0)$
 3 act as integration weights. In the case of (21) (the case of (22) is analogous), one considers orthogonal polynomials
 4 $\{P_j(y_1)\}_{j=0}^\infty$ (the degree of P_j is j) with respect to the inner product of $L^2(\mathcal{S}_{Y_1})$ with weight function $f_{Y_1}(y_1)$. For
 5 example, if Y_1 has a Normal or Uniform distribution, then $\{P_j(y_1)\}_{j=0}^\infty$ corresponds to the family of Hermite and
 6 Legendre polynomials, respectively. In other cases where Y_1 does not possess a standard probability distribution from
 7 the Askey scheme, $\{P_j(y_1)\}_{j=0}^\infty$ may be constructed through a Gram-Schmidt orthogonalization procedure from the
 8 canonical basis $\{1, y_1, y_1^2, y_1^3, \dots\}$. The integral (21) is approximated as follows:

$$\frac{1}{|s_{0,i}^N|} \int_{\mathcal{S}_{Y_1}} f_{Y_0} \left(\frac{x - y_1 s_{1,i}^N}{s_{0,i}^N} \right) f_{Y_1}(y_1) dy_1 \approx \frac{1}{|s_{0,i}^N|} \sum_{j=1}^Q w_{1,j}^{(Q)} f_{Y_0} \left(\frac{x - y_{1,j}^{(Q)} s_{1,i}^N}{s_{0,i}^N} \right). \quad (23)$$

9 Here, Q is the finite degree of the quadrature, the nodes $\{y_{1,j}\}_{j=1}^Q$ are the zeros of $P_Q(y_1)$, and $\{w_{1,j}^{(Q)}\}_{j=1}^Q$ are the
 10 weights so that the integration rule is exact for polynomials of degree less than or equal to $2Q - 1$

11 The nodes $\{y_{0,j}\}_{j=1}^Q$, $\{y_{1,j}\}_{j=1}^Q$, and the weights $\{w_{0,j}^{(Q)}\}_{j=1}^Q$, $\{w_{1,j}^{(Q)}\}_{j=1}^Q$, only depend on f_{Y_0} and f_{Y_1} , therefore
 12 they are independent of the realization $(s_{0,i}^N, s_{1,i}^N)$, the number of simulations M and the truncation order N . Hence,
 13 the nodes and the weights should always be computed once for all at the beginning.

14 As more regularity we demand on f_{Y_0} and f_{Y_1} , faster convergence the quadrature rule will achieve with Q . The
 15 extreme case appears when f_{Y_0} and f_{Y_1} are analytic real functions: the quadrature rule converges exponentially with
 16 Q and therefore the bias error $\theta_N(x, t) = f_{X(t)}(x) - f_{X^N(t)}(x)$ is not seriously affected by the deterministic error
 17 arisen from the finite degree Q . Nonetheless, numerical problems in the quadrature approximations may arise when

1 the integrands are strongly peaked. In this case, the degree Q must be taken larger, otherwise the bias error may
 2 increase dramatically.

3 The procedure is described in the following steps:

4 **PROCEDURE 2.**

5 **Part 1** Generate M realizations of $S_0^N(t)$ and $S_1^N(t)$.

6 **Part 2** For each realization, if $|S_0^N(t)| \geq |S_1^N(t)|$, consider the realization $\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]$ approximated
 7 by the quadrature $\sum_{j=1}^Q w_{1,j}^{(Q)} Z_0^N(x, t)|_{Y_1=y_{1,j}^{(Q)}}$ (see (10), (12), (21), (23)). Otherwise consider the realization
 8 $\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]$ approximated by $\sum_{j=1}^Q w_{0,j}^{(Q)} Z_1^N(x, t)|_{Y_0=y_{0,j}^{(Q)}}$.

9 **Part 3** Compute the sample mean of the M realizations of $\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]$. This value estimates the expec-
 10 tation $f_{X^N(t)}(x) = \mathbb{E}[Z^N(x, t)] = \mathbb{E}[\mathbb{E}[Z^N(x, t)|S_0^N(t), S_1^N(t)]]$ with MC simulation. We denote the estimate
 11 (output function) by $f_X^{N,M,Q}(x, t)$.

12 The complexity of Procedure 2 is $\mathcal{O}(M(N^2 + Q))$, in general. Recall that one realization of $S_0^N(t)$ and $S_1^N(t)$ re-
 13 quires $\mathcal{O}(N^2)$ operations when $A(t)$ and $B(t)$ are given by infinite random expansions. The quadrature rule demands
 14 $\mathcal{O}(Q)$ operations.

15 When $A(t)$ and $B(t)$ are random polynomials, one realization of $S_0^N(t)$ and $S_1^N(t)$ requires $\mathcal{O}(N)$ operations
 16 only. The complexity of the whole algorithm thus becomes $\mathcal{O}(M(N + Q))$.

17 In the case of simple problems (1) in which $S_0(t)$ and $S_1(t)$ are exactly known, Procedure 2 is run with $N = \infty$,
 18 that is, by computing exact realizations of $S_0(t)$ and $S_1(t)$. The error in this case comes from the MC simulation and
 19 the finite degree of the quadrature rule. The total complexity is given by $\mathcal{O}(MQ)$.

20 We finish this section with numerical experiments concerning the methodology and Procedure 2. In the following
 21 three examples, we will compare Procedure 1 and Procedure 2 in terms of variance reduction, density approximation
 22 and error against complexity. The first two examples, Example 3 and Example 4, deal with problems having closed-
 23 form expressions for the fundamental sets ($N = \infty$). In Example 3, the initial conditions Y_0 and Y_1 have large
 24 dispersion compared to $S_0(t)$ and $S_1(t)$ (this is checked using Sobol indices), so a significant variance reduction of
 25 the MC estimator is achieved by employing quadrature rules with respect to f_{Y_0} and f_{Y_1} . On the contrary, Example 4
 26 presents two initial conditions with small dispersion compared to $S_0(t)$ and $S_1(t)$, hence the crude MC approach from
 27 Procedure 1 is more efficient. The third example, Example 5, deals with orders of truncation $N < \infty$, which implies
 28 that computing the fundamental set $\{S_0(t), S_1(t)\}$ of the problem takes certain computational burden. This entails

1 that, albeit the initial conditions have small dispersion compared to $S_0(t)$ and $S_1(t)$, both Procedure 1 and Procedure 2
 2 present similar efficiency. After the three examples, we will conclude the section with specific recommendations
 3 regarding the use of Procedure 1 or Procedure 2.

4 **Example 3.** Consider (15) with input data $B \sim \text{Uniform}(1, 1.01)$ and $Y_0, Y_1 \sim \text{Normal}(1, \sigma = 2)$, all of them
 5 independent. As the fundamental set $\{S_0(t), S_1(t)\}$ is known with closed-form expressions, $S_0(t) = \cos(Bt)$ and
 6 $S_1(t) = \sin(Bt)/B$, we take $N = \infty$, by sampling from $S_0(t)$ and $S_1(t)$ directly.

7 Here the randomness mainly comes from the initial conditions. This is checked using the Sobol indices from (19)
 8 and (20). As proposed in [34,35], the Sobol indices are computable using MC simulation with the formulas

$$\mathbb{V} [Z_{S_0(t), S_1(t)}] \approx \frac{1}{M} \sum_{i=1}^M Z(s_i^I, y_i^I) Z(s_i^I, y_i^{II}) - \frac{1}{M} \sum_{i=1}^M Z(s_i^I, y_i^I) Z(s_i^{II}, y_i^{II}), \quad (24)$$

9

$$\mathbb{V} [Z_{Y_0, Y_1}] \approx \frac{1}{M} \sum_{i=1}^M Z(s_i^I, y_i^I) Z(s_i^{II}, y_i^I) - \frac{1}{M} \sum_{i=1}^M Z(s_i^I, y_i^I) Z(s_i^{II}, y_i^{II}), \quad (25)$$

10 where $\{s_i^I\}_{i=1}^M \subseteq \mathbb{R}^2$ and $\{s_i^{II}\}_{i=1}^M \subseteq \mathbb{R}^2$ denote two sets of joint realizations of $(S_0(t), S_1(t))$, and $\{y_i^I\}_{i=1}^M \subseteq \mathbb{R}^2$
 11 and $\{y_i^{II}\}_{i=1}^M \subseteq \mathbb{R}^2$ denote two sets of joint realizations of (Y_0, Y_1) , being all of them independent. The notation used
 12 here, for instance $Z(s_i^I, y_i^I)$, means evaluating $Z(x, t)$ (given by (11)) at those specific realizations. In Figure 6, we
 13 report the estimates of the Sobol indices for $t = 10$. We observe that $\hat{\mathbb{S}}_{Y_0, Y_1}$ is very close to 1, therefore indicating
 14 that the variability of $Z(x, t)$ is dominated by the initial conditions. Thus we expect Procedure 2 to be superior to
 15 Procedure 1 in terms of variance reduction.




FIG. 6: Estimated Sobol indices of the random inputs for $Z(x, t = 10)$. This figure corresponds to Example 3.

16 In this example, the Gaussian quadrature rule is based on Hermite polynomials. We work at time $t = 10$, with

1 degree of quadrature $Q = 30$. Essentially, the quadrature rule approximation is error-free. In Figure 7 we plot esti-
 2 mates $\hat{V}[Z(x, 10)]$ and $\hat{V}[\mathbb{E}[Z(x, 10)|S_0(10), S_1(10)]]$. Observe that the magnitude of $\hat{V}[\mathbb{E}[Z(x, 10)|S_0(10), S_1(10)]]$
 3 is smaller than $\hat{V}[Z(x, 10)]$, due to the law of total variance (18). Anyway, none of the variances becomes large at any
 4 point x , therefore good estimates of $f_{X(t=10)}(x)$ are obtained with any of the two algorithms.




FIG. 7: Estimates of $\hat{V}[\mathbb{E}[Z(x, 10)|S_0(10), S_1(10)]]$ and $\hat{V}[Z(x, 10)]$. This figure corresponds to Example 3.

5 In Figure 8 we depict the estimates for $f_{X(t=10)}(x)$ with Procedure 1 (legend “Density MC”) and Procedure 2
 6 (legend “Density MCQ”). Very similar smooth estimates are obtained, in fact they overlap, with $Q = 30$ and $M =$
 7 10,000 realizations. We also compare the two estimates with the exact density function (legend “Exact density”).
 8 The exact density function is computable using a cubature rule for the double integral (4) (cubature rule constructed
 9 using a tensor product approach based on univariate Gauss-Hermite and Gauss-Legendre quadrature rules), because
 10 the integrand is smooth with no peaks (see Figure 9), thus posing no difficulties. The second panel of the figure shows
 11 the plot in logarithmic scale, to better appreciate the agreement at the tails.




FIG. 8: Estimates of $f_{X(t=10)}(x)$ with MC simulation, MC simulation plus quadrature rule (MCQ), and exact density function. The right panel shows the plot in logarithmic scale. This figure corresponds to Example 3.




FIG. 9: 3D plots of the integrand $Z_0(x, t = 10)$ at different points x . This figure corresponds to Example 3.

1 A further analysis is performed by studying the relation between the error and the complexity of Procedure 1
 2 and Procedure 2. In Figure 10 we show plots of mean relative error versus complexity in log-log scale for Proce-
 3 dure 1 (legend “MC”) and Procedure 2 (legend “MCQ”), at different points x . Here we consider the complexity of
 4 Procedure 1 as M , while the complexity of Procedure 2 as MQ , being $Q = 30$ (recall that M denotes the number
 5 of realizations). The relative error is defined as the absolute error divided by the value of the exact density function.
 6 We run both algorithms up to complexity $\text{floor}(1.4^{44})$, and we keep the relative errors for complexities $\text{floor}(1.4^i)$,
 7 $i = 14, 15, \dots, 44$ (a geometric progression makes equidistant complexities in log-scale). This gives a discretized
 8 realizable path of relative errors. We repeat this demanding process $n = 20$ times with parallel computing to obtain
 9 an estimate of the mean relative error (so that the analysis performed does not depend upon a specific realizable
 10 relative error). A 95% confidence interval of Gaussian-type for the mean relative error is constructed (this is justi-
 11 fied by the central limit theorem), as mean plus/minus 1.96 times the standard error, where the standard error is the
 12 standard deviation of the original sample of length n divided by \sqrt{n} *. The relative error descends with slope 1/2
 13 approximately, because the error of the MC simulation goes down with M at rate $1/\sqrt{M}$. The mean relative error of
 14 Procedure 2 is shifted down the mean relative error of Procedure 1, due to the variance reduction of the estimators,
 15 thus indicating higher efficiency. Similar plots have been obtained for the points $x \in \{-1, 3\}$. Indeed, the effect of the
 16 initial conditions on the variability of Z is homogeneous in x , according to the Sobol indices from Figure 6, therefore

*Here n might not be that large to apply the central limit theorem. Other types of confidence intervals may be considered, for instance those constructed via bootstrapping [36] with b samples with replacement from the original sample of length n . The percentile confidence interval has as left endpoint the empirical 0.025 percentile, and as right endpoint the empirical 0.975 percentile [36, p. 170], [37, p. 203]. The pivotal confidence interval generated by bootstrapping has left endpoint given by twice the mean minus the empirical 0.975 percentile, and right endpoint given by twice the mean minus the empirical 0.025 percentile [37, p. 194]. Finally, another Gaussian-type confidence interval may consider the mean plus/minus 1.96 times the bootstrap standard deviation of the mean. In all our examples, we checked that these confidence intervals are coinciding, thus showing robustness of our estimates. As n increases, better estimates for the mean are obtained.

1 a similar performance of the algorithm is expected for the different x 's.




FIG. 10: Plots in log-log scale of mean relative error versus complexity, at $t = 10$. The dashed lines represent 95% confidence intervals. This figure corresponds to Example 3.

2 **Example 4.** We consider the same setting as Example 1: (15) with $B \sim \text{Uniform}(1, 2.5)$ and $Y_0, Y_1 \sim \text{Normal}(1, \sigma =$
 3 $0.1)$, all of them independent.

4 In this case, the variability of Z is mainly due to B . This is deduced from Figure 11, where we report the Sobol
 5 indices (19) and (20) for $t = 10$, using the MC estimates (24) and (25). The variance decomposition of Z is dominated
 6 by the effect of the fundamental set and the interactions, being the estimated Sobol index corresponding to the initial
 7 conditions very close to 0. Therefore, we do not expect much variance reduction from the quadrature rule.




FIG. 11: Estimated Sobol indices of the random inputs for $Z(x, t = 10)$. This figure corresponds to Example 4.

8 We work at $t = 10$, with degree of quadrature $Q = 30$ and Hermite polynomials. In Figure 12, left panel,
 9 we plot the variances of the statistical estimators from Procedure 1 and Procedure 2. We observe that the variances
 10 are greater than in Example 3. Also, in contrast with Example 3, the variance is not much reduced now with the
 11 quadrature rule. This was already expected when analyzing the effect of the inputs on the variability of Z using Sobol
 12 indices. Figure 12, right panel, depicts the estimated density function $f_{X(t=10)}(x)$ with Procedure 1 (legend “MC”)
 13 and Procedure 2 (legend “MCQ”), by setting $M = 40,000$ realizations and $Q = 30$. Both algorithms show similar

1 estimates.




FIG. 12: Left: estimates of $\hat{\mathbb{V}}[\mathbb{E}[Z(x, 10)|S_0(10), S_1(10)]]$ and $\hat{\mathbb{V}}[Z(x, 10)]$. Right: estimates of $f_{X(t=10)}(x)$ with MC simulation, and MC simulation plus quadrature rule (MCQ). This figure corresponds to Example 4.

2 Figure 13 reports the log-log relationship between the relative error and the complexity at $t = 10$, for different
 3 points x . The setting is analogous to Example 3. We consider the complexities of Procedure 1 and Procedure 2 as M
 4 and MQ , respectively, where $Q = 30$ and M is the number of realizations. The “absolute error” is calculated with
 5 respect to the density function obtained from Procedure 1 with 100,000,000 realizations (this is considered as our
 6 “exact density function”). We define the relative error as such absolute error divided by the value of the “exact density
 7 function”. We run both algorithms up to complexity $\text{floor}(1.4^{44})$, while keeping the relative errors for complexities
 8 $\text{floor}(1.4^i)$, $i = 14, 15, \dots, 44$. The procedure is repeated $n = 20$ times to determine an estimate for the mean relative
 9 error. A 95% Gaussian confidence interval for the mean relative error is constructed. See Example 3 for further details.
 10 We observe that the relative errors decay approximately with slope $1/2$, due to the rate of convergence proportional
 11 to $1/\sqrt{M}$ of the MC simulation. In this example, the decay rate with slope $1/2$ is true for both algorithms as long
 12 as the sampling error of the “exact density function” does not become dominant. In the figure, we observe that the
 13 estimated relative error of Procedure 1 tends to be constant sometime after 10^6 complexity, as it becomes of similar
 14 order of magnitude to the sampling error of the “exact density function”. Procedure 1 shows superior to Procedure 2
 15 in this example: in the plots, the depicted relation between the error and the complexity of Procedure 2 is shifted to
 16 the right, because of the multiplication by Q of the number of realizations M in the complexity. As the effect of the
 17 initial conditions on the variability of Z is small, the potentiality of Procedure 2 is lessened.

18 In the two Examples 3, 4 presented until now, we studied two cases. When $Y_0 - Y_1$ produce high variability on
 19 $\mathbb{V}[Z(x, t)]$, then Procedure 2 clearly wins Procedure 1 in terms of error versus complexity. On the contrary, when
 20 the dispersion of $Y_0 - Y_1$ is small compared to $S_0(t)$ and $S_1(t)$, Procedure 2 does no longer beat Procedure 1. But in




FIG. 13: Plots in log-log scale of mean relative error versus complexity, at $t = 10$. The dashed lines represent 95% confidence intervals. This figure corresponds to Example 4.

1 general, when the cost of obtaining realizations of $S_0(t)$ and $S_1(t)$ is larger (because we are working with truncated
 2 power series in N , for instance), the cost due to the MC procedure alone is comparable with the whole complexity.
 3 In such a case, Procedure 1 never defeats Procedure 2 by a big difference. This is the case studied in the following
 4 Example 5.

5 **Example 5.** Consider (16) with inputs $A_0 = 4$, $A_1 \sim \text{Uniform}(0, 1)$, $B_0 \sim \text{Triangular}(-1, 0)$, $B_1 \sim \text{Bernoulli}(0.35)$
 6 and $Y_0, Y_1 \sim \text{Normal}(1, \sigma = 0.1)$, all of them independent. The random coefficients A_0, A_1, B_0 and B_1 are bounded
 7 and the initial conditions Y_0 and Y_1 have finite absolute moments, therefore there is an analytic solution $X(t)$ in the
 8 $L^p(\Omega)$ sense, $1 \leq p < \infty$.

9 In this example, the fundamental set cannot be written with simple closed-form formulas. One needs to perform
 10 a dimensionality reduction of the problem by truncating the random power series at certain order N . We work at time
 11 $t = 1.5$ with order of truncation $N = 15$. The goal is to compare the approximations to the density $f_{X^{N=15}(t=1.5)}(x)$
 12 offered by Procedure 1 and Procedure 2.

13 In Figure 14 we show the Sobol indices (19) and (20) for $t = 1.5$, computed with the MC estimates (24) and (25).
 14 The Sobol indices demonstrate that the effect of $Y_0 - Y_1$ on the variability of $Z^{N=15}(x, t = 1.5)$ is very small.




FIG. 14: Estimated Sobol indices of the random inputs for $Z^{N=15}(x, t = 1.5)$. This figure corresponds to Example 5.

1 Hence, the variance reduction due to the quadrature rule is not very significant. This is confirmed by Figure 15,
 2 left panel, where we depict the variances of the statistical estimators. We use quadrature degree $Q = 30$, so that
 3 the expectation with respect to $Y_0 - Y_1$ is practically error-free. In the right plot, we estimate the density function
 4 $f_{X^{N=15}(t=1.5)}(x)$ using Procedure 1 and Procedure 2, in logarithmic scale. We generated $M = 10,000$ realizations.




FIG. 15: Left: estimates of $\hat{V}[\mathbb{E}[Z^{15}(x, 1.5)|S_0(1.5), S_1(1.5)]]$ and $\hat{V}[Z^{15}(x, 1.5)]$. Right: estimates of $f_{X^{15}(1.5)}(x)$ (the legend MCQ refers to MC plus quadrature rule), in logarithmic scale. This figure corresponds to Example 5.

5 An analogous analysis to Example 4 of relative error versus complexity is carried out here. The complexity of
 6 Procedure 1 is MN , whereas the complexity of Procedure 2 is $M(N+Q)$. Recall that M is the number of realizations,
 7 N is the truncation order for the power series, and Q is the quadrature degree. We work at the point $x = 1.4$, which
 8 belongs to the density domain (any other point is equally valid). We consider as our “exact density function” the output
 9 of Procedure 1 with $M = 10,000,000$ realizations, i.e., complexity $MN = 150,000,000$. We run both algorithms for
 10 complexities $\text{floor}(1.4^{37})$, and keep the relative errors for complexities $\text{floor}(1.4^i)$, $i = 14, 15, \dots, 37$. The procedure
 11 is repeated $n = 20$ times (in parallel) to obtain the mean relative error. A 95% Gaussian confidence interval for
 12 the mean relative error has been constructed. Procedure 2 is run with $Q = 10$ and $Q = 30$ degrees of quadrature.
 13 Figure 16 shows the results. Notice that, although $Y_0 - Y_1$ have small dispersion compared with the fundamental set,
 14 both Algorithms show similar performances. Indeed, as each realization of the truncated fundamental set requires
 15 now more expense (cost N), Procedure 1 cannot defeat Procedure 2 by a big difference. The relative errors decrease
 16 with slope $1/2$, but for $Q = 10$ we perceive constant error for complexities greater than 10^5 . This is because the bias
 17 error caused by the quadrature rule of degree $Q = 10$ has been attained, in contrast to the case $Q = 30$. This reflects
 18 the importance of selecting a sufficiently large quadrature degree, especially when f_{Y_0} and f_{Y_1} are peaked by small
 19 dispersion of Y_0 and Y_1 .

20 The conclusion of this section is summarized as follows:




FIG. 16: Plots in log-log scale of mean relative error versus complexity, at $t = 1.5$ and $N = 15$. The dashed lines represent 95% confidence intervals. This figure corresponds to Example 5.

1 **Case 1** When the dispersion of $Y_0 - Y_1$ is large compared with $S_0(t)$ and $S_1(t)$ (this is checked with Sobol indices),
 2 then Procedure 2 should be used. See Example 3.

3 **Case 2** When the variability of $Y_0 - Y_1$ is small compared with $S_0(t)$ and $S_1(t)$ (Sobol indices), and each realization of
 4 $S_0(t)$ and $S_1(t)$ takes certain computational burden (for instance, because we are dealing with truncated power
 5 series), then Procedure 1 and Procedure 2 are similar in terms of error versus complexity. See Example 5.

6 **Case 3** Finally, if the variability of $Y_0 - Y_1$ is small compared with $S_0(t)$ and $S_1(t)$ (Sobol indices), and each realiza-
 7 tion of $S_0(t)$ and $S_1(t)$ is immediately computed (when $S_0(t)$ and $S_1(t)$ are known with simple closed-form
 8 expressions), then Procedure 1 beats Procedure 2. See Example 4.

9 In any of the three cases, Q should be taken large enough to avoid a quadrature error greater than the sampling error.

10 **4. MULTILEVEL MC SIMULATION**

11 The idea of multilevel MC [27,28] is to speed up the convergence of the MC simulation by decomposing the target
 12 expectation into the sum of expectations of increments (through a telescopic sum identity), whose variances decay
 13 rapidly. In our case, we consider the following increments:

$$\Delta Z_0^l(x, t) = Z_0^{l+1}(x, t) - Z_0^l(x, t), \quad \Delta Z_1^l(x, t) = Z_1^{l+1}(x, t) - Z_1^l(x, t), \quad (26)$$

for each level $l \geq 0$, where each term was defined in (10). If we start from the level $l_0 \geq 0$, these increments yield the following telescopic sum identities, for $i \in \{1, 2\}$:

$$f_{X(t)}(x) = \mathbb{E} \left[Z_i^{l_0}(x, t) \right] + \sum_{l=l_0}^{\infty} \mathbb{E} \left[\Delta Z_i^l(x, t) \right] \approx f_{X^N(t)}(x) = \mathbb{E} \left[Z_i^{l_0}(x, t) \right] + \sum_{l=l_0}^{N-1} \mathbb{E} \left[\Delta Z_i^l(x, t) \right].$$

1 Each one of these expectations is approximated with MC simulation. The rate of convergence of the MC simulation
 2 depends on the number of realizations and $\mathbb{V}[Z_0^{l_0}(x, t)]$, $\mathbb{V}[\Delta Z_0^{l_0}(x, t)]$, and $\mathbb{V}[Z_1^{l_0}(x, t)]$, $\mathbb{V}[\Delta Z_1^{l_0}(x, t)]$. If $Z_0^l(x, t)$ or
 3 $Z_1^l(x, t)$ tend to $Z^l(x, t)$ in the mean square sense, then the variances of the increments, $\mathbb{V}[\Delta Z_0^l(x, t)]$, $\mathbb{V}[\Delta Z_1^l(x, t)]$,
 4 converge to 0, which makes the convergence of the MC procedure faster. The first level l_0 is usually chosen satisfying
 5 that, for $l \geq l_0$, the variances of the increments per level l are decaying monotonically and rapidly.

6 However, as we are selecting either f_{Y_0} or f_{Y_1} (as in [25]), this may lead to numerical instabilities due to small
 7 denominators and large or infinite variance, which causes noisy features and slows down the convergence of the MC
 8 procedure. This may hide the potential of multilevel MC.

9 To sort out this issue, we path-wise select $\Delta Z_0^l(x, t)$ or $\Delta Z_1^l(x, t)$, by discarding the increment with the lowest
 10 denominator. The minimum among the denominators is considered as a time-dependent random variable

$$V^l(t) = \min \{ |S_0^{l+1}(t)|, |S_0^l(t)|, |S_1^{l+1}(t)|, |S_1^l(t)| \}. \quad (27)$$

11 We pick the index with no smallest denominator through the random variable

$$U^l(t) = \{1 - i : \text{the minimum of } V^l(t) \text{ is attained at } i \in \{0, 1\}\}. \quad (28)$$

12 This random variable gives rise to two events, $H_0^l(t) = \{U^l(t) = 0\}$ and $H_1^l(t) = \{U^l(t) = 1\}$. Let $Z_\Delta^l(x, t) =$
 13 $\Delta Z_0^l(x, t)1_{H_0^l(t)} + \Delta Z_1^l(x, t)1_{H_1^l(t)}$. This random process models the selection of $\Delta Z_0^l(x, t)$ and $\Delta Z_1^l(x, t)$, according
 14 to whether their expressions do not have the smallest denominator.

15 **Proposition 4.** *We have the following sum identities for $f_{X(t)}(x)$ and $f_{X^N(t)}(x)$:*

$$f_{X(t)}(x) = \mathbb{E} \left[Z^{l_0}(x, t) \right] + \sum_{l=l_0}^{\infty} \mathbb{E} \left[Z_\Delta^l(x, t) \right], \quad f_{X^N(t)}(x) = \mathbb{E} \left[Z^{l_0}(x, t) \right] + \sum_{l=l_0}^{N-1} \mathbb{E} \left[Z_\Delta^l(x, t) \right].$$

16 With the objective of approximating the target density function $f_{X(t)}(x)$ pointwise, we truncate the telescopic
 17 sum identity at increment $N - 1$, which causes the bias error $\theta_N(x, t) = f_{X(t)}(x) - f_{X^N(t)}(x)$: $f_{X(t)}(x) \approx$

1 $f_{X^N(t)}(x) = \mathbb{E}[Z^{l_0}(x, t)] + \sum_{l=l_0}^{N-1} \mathbb{E}[Z_\Delta^l(x, t)]$. The expectations $\mathbb{E}[Z^{l_0}(x, t)]$, $\mathbb{E}[Z_\Delta^l(x, t)]$, $l = l_0, \dots, N-1$, are
 2 computed independently with MC simulation, whose rate of convergence depends on the finite number of realizations
 3 and on $\mathbb{V}[Z^{l_0}(x, t)]$, $\mathbb{V}[Z_\Delta^l(x, t)]$, $l = l_0, \dots, N-1$.

4 We have $\|Z_\Delta^l(x, t)\|_2 \leq \|\Delta Z_0^l(x, t)\|_2 + \|\Delta Z_1^l(x, t)\|_2$. Thus, if both $\Delta Z_0^l(x, t)$ and $\Delta Z_1^l(x, t)$ have finite vari-
 5 ances, then so does $Z_\Delta^l(x, t)$. On the contrary, if $\Delta Z_0^l(x, t)$ or $\Delta Z_1^l(x, t)$ have infinite variances, then $Z_\Delta^l(x, t)$ may
 6 have finite variance, instead. In any of these cases, we cannot get worse in terms of infinite variance.

7 We hope to have the variances $\mathbb{V}[Z^{l_0}(x, t)]$ and $\mathbb{V}[Z_\Delta^l(x, t)]$, $l = l_0, \dots, N-1$, finite, to exploit the decay of
 8 $\mathbb{V}[Z_\Delta^l(x, t)]$ to 0 with l . The first level l_0 is chosen such that $\mathbb{V}[Z_\Delta^l(x, t)]$ is decreasing with $l \geq l_0$, monotonically
 9 and fast. The best case is when $\log \mathbb{V}[Z_\Delta^l(x, t)]$ is decreasing linearly with $l \geq l_0$, as exponential decay holds for the
 10 variances of the increments.

11 Let $M = (N_{l_0}; M_{l_0}, \dots, M_{N-1})$ be the number of simulations performed: N_{l_0} for $Z^{l_0}(x, t)$, M_l for $Z_\Delta^l(x, t)$,
 12 $l = l_0, \dots, N-1$. In principle, the number of simulations may depend on (x, t) , although we drop this possible
 13 dependence along this discussion.

14 Let $f_X^{N,M}(x, t)$ be the final approximation to $f_{X(t)}(x)$:

$$f_X^{N,M}(x, t) = \frac{1}{N_{l_0}} \sum_{i=1}^{N_{l_0}} Z^{l_0}(x, t, \omega_{i,l_0}) + \sum_{l=l_0}^{N-1} \frac{1}{M_l} \sum_{i=1}^{M_l} Z_\Delta^l(x, t, \omega_{i,l,\Delta}), \quad (29)$$

where ω with the appropriate subindices denote outcomes. The mean square error in the approximation to $f_{X^N(t)}(x) =$
 $\mathbb{E}[f_X^{N,M}(x, t)]$ is given by the mean square norm of $\mathcal{E}_{N,M}(x, t) = f_{X^N(t)}(x) - f_X^{N,M}(x, t)$, which is

$$\left\| f_{X^N(t)}(x) - f_X^{N,M}(x, t) \right\|_2^2 = \mathbb{V}[\mathcal{E}_{N,M}(x, t)] = \mathbb{V}[f_X^{N,M}(x, t)] = \frac{\mathbb{V}[Z^{l_0}(x, t)]}{N_{l_0}} + \sum_{l=l_0}^{N-1} \frac{\mathbb{V}[Z_\Delta^l(x, t)]}{M_l}. \quad (30)$$

15 Notice that the first equality holds because $\mathbb{E}[\mathcal{E}_{N,M}(x, t)] = 0$, so the squared norm is equal to the variance. The
 16 second equality uses the fact that $f_{X^N(t)}(x)$ is not random. Finally, the third equality is deduced by applying the
 17 variance in (29). The total cost is

$$W_{N,M} = N_{l_0} c_{l_0} + \sum_{l=l_0}^{N-1} M_l C_l, \quad (31)$$

18 where c_{l_0} denotes the number of operations per realization of $Z^{l_0}(x, t)$, and C_l denotes the number of operations per
 19 realization of $Z_\Delta^l(x, t)$, $l = l_0, \dots, N-1$. In our case, we consider $c_l = C_l = l^2$ when dealing with $A(t)$ and $B(t)$
 20 as random infinite expansions, and $c_l = C_l = l$ when $A(t)$ and $B(t)$ are random polynomials.

1 **Proposition 5.** *Given a fixed variance (30) $\epsilon_{\text{sampling}}^2$, the cost (31) is minimized by selecting*

$$N_{l_0}(x, t) = \epsilon_{\text{sampling}}^{-2} \left(\sqrt{\mathbb{V}[Z^{l_0}(x, t)] c_{l_0}} + \sum_{l=l_0}^{N-1} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} \right) \sqrt{\frac{\mathbb{V}[Z^{l_0}(x, t)]}{c_{l_0}}}, \quad (32)$$

2

$$M_l(x, t) = \epsilon_{\text{sampling}}^{-2} \left(\sqrt{\mathbb{V}[Z^{l_0}(x, t)] c_{l_0}} + \sum_{l=l_0}^{N-1} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} \right) \sqrt{\frac{\mathbb{V}[Z_{\Delta}^l(x, t)]}{C_l}}, \quad (33)$$

3 for $l = l_0, \dots, N - 1$. In such a case, the total cost (31) is given by

$$W_{N,M} = \epsilon_{\text{sampling}}^{-2} \left(\sqrt{\mathbb{V}[Z^{l_0}(x, t)] c_{l_0}} + \sum_{l=l_0}^{N-1} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} \right)^2. \quad (34)$$

4 This proposition is a simple application of Lagrange multipliers.

5 For a truncation order N , and for a sampling error variance $\epsilon_{\text{sampling}}^2$ by running $\epsilon_{\text{sampling}}^{-2} \mathbb{V}[Z^N(x, t)]$ simulations,
 6 the cost of Procedure 1 is $\epsilon_{\text{sampling}}^{-2} \mathbb{V}[Z^N(x, t)] c_N$. For fixed N , both approaches (standard MC in Procedure 1 and
 7 multilevel MC) yield a cost $\mathcal{O}(\epsilon_{\text{sampling}}^{-2})$. However, there is a significant difference when N grows. The complexity
 8 $c_N = N$ (if $A(t)$ and $B(t)$ are random polynomials) or $c_N = N^2$ (if $A(t)$ and $B(t)$ are infinite random series) per
 9 realization of $Z^N(x, t)$ grows to infinity. Then the cost of Procedure 1 tends to infinity as $N \rightarrow \infty$, linearly (if $c_N =$
 10 N) or quadratically (if $c_N = N^2$). If we choose $N = \mathcal{O}(\log \epsilon_{\text{sampling}}^{-1})$ to ensure a bias error less than $\epsilon_{\text{sampling}}$ (under
 11 exponential convergence of $f_{X^N(t)}(x)$ with N), then the complexity of Procedure 1 is $\mathcal{O}(\epsilon_{\text{sampling}}^{-2} \log^2 \epsilon_{\text{sampling}})$, when
 12 $c_N = N^2$, and $\mathcal{O}(\epsilon_{\text{sampling}}^{-2} \log \epsilon_{\text{sampling}}^{-1})$, when $c_N = N$. However, in the multilevel approach the cost (34) remains
 13 $\mathcal{O}(\epsilon_{\text{sampling}}^{-2})$ despite having $N \rightarrow \infty$, as $\sqrt{\mathbb{V}[Z^{l_0}(x, t)] c_{l_0}} + \sum_{l=l_0}^{\infty} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} < \infty$ in general. For instance,
 14 when $\mathbb{V}[Z_{\Delta}^l(x, t)]$ decreases exponentially with l . Hence, from this theoretical analysis, multilevel MC is superior to
 15 standard MC, at least when N grows.

16 In practice, multilevel MC is more efficient than standard MC by choosing the initial level l_0 and truncation
 17 $N > l_0$ appropriately, which can be done heuristically as explained later.

18 In numerical applications, the main computational burden of multilevel MC is due to the number of samples N_{l_0}
 19 of $Z^{l_0}(x, t)$. For the last levels near N , which are the most complex ones, we run only few samples. In general, from
 20 a certain level l , only 1 realization of $Z_{\Delta}^l(x, t)$ is required. From a certain order of truncation N , the complexity of
 21 multilevel MC will be constant; whereas the complexity of the standard MC strategy will be growing up to infinity
 22 with N , linearly or quadratically. In our setting, multilevel MC turns out to be useful when high accuracy is required.

23 Given l_0 , $N > l_0$ and $\epsilon_{\text{sampling}}$, the cost of multilevel MC is approximately $\epsilon_{\text{sampling}}^{-2} \mathbb{V}[Z^{l_0}(x, t)] c_{l_0}$, because the

1 sequence $\mathbb{V}[Z_{\Delta}^l(x, t)]C_l$ decreases rapidly with l . The expense of standard MC is $\epsilon_{\text{sampling}}^{-2} \mathbb{V}[Z^N(x, t)]c_N$. Assuming
 2 that $\mathbb{V}[Z^{l_0}(x, t)] \approx \mathbb{V}[Z^N(x, t)]$, which is quite realistic if l_0 and $N > l_0$ are not too small, the cost of multilevel MC
 3 is reduced by factor c_{l_0}/c_N , approximately. The largest reduction is thus obtained for problems (1) with infinite series
 4 inputs, since in such a case c_N grows quadratically with N .

5 The complete multilevel MC approach is presented in what follows. Fixed the time t and the maximum level N ,
 6 the whole procedure is divided into different parts:

7 **PROCEDURE 3.**

8 **Part 1** We determine $\mathbb{V}[Z_{\Delta}^l(x, t)]$ for $0 \leq l \leq N - 1$. They are estimated à la MC, as Procedure 1, but not using an
 9 excessive amount $m_{\mathbb{V}}$ of realizations. The main loop over the levels l can be carried out in parallel.

10 **Part 2** We estimate $\mathbb{V}[Z^l(x, t)]$ for $1 \leq l \leq N$ (with parallel computing). Then we set the ratios between the
 11 complexities of Procedure 1,

$$W_N^{MC}(t) = \epsilon_{\text{sampling}}^{-2} c_N \max_x \mathbb{V}[Z^N(x, t)], \quad (35)$$

12 and of the multilevel MC strategy with initial level $l_1 = 1, \dots, N - 1$,

$$W_{l_1, N}^{multi}(t) = \epsilon_{\text{sampling}}^{-2} \max_x \left(\sqrt{\mathbb{V}[Z^{l_1}(x, t)] c_{l_1}} + \sum_{l=l_1}^{N-1} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} \right)^2. \quad (36)$$

13 These estimates are done via standard MC simulation with not too many realizations $m_{\mathbb{V}}$. The ratio of the
 14 complexities is given by

$$R_{l_1, N}(t) = \frac{W_N^{MC}(t)}{W_{l_1, N}^{multi}(t)} = \frac{\max_x \mathbb{V}[Z^N(x, t)]c_N}{\max_x \left(\sqrt{\mathbb{V}[Z^{l_1}(x, t)] c_{l_1}} + \sum_{l=l_1}^{N-1} \sqrt{\mathbb{V}[Z_{\Delta}^l(x, t)] C_l} \right)^2}. \quad (37)$$

15 Notice that this ratio is independent of any sampling error $\epsilon_{\text{sampling}}^2$. In order not to have x -dependent complex-
 16 ities, we take the maximum on x lying in the density domain of interest. If $R_{l_1, N}(t) > 1$, then multilevel MC
 17 starting from level l_1 is more efficient than standard MC.

18 We define the first level $l_0 = l_0(N, t)$ as the maximizer of

$$\max_{l_1: 1 \leq l_1 < N} R_{l_1, N}(t) > 1. \quad (38)$$

19 This way we are achieving the maximum optimality of multilevel MC. In general, the variances $\mathbb{V}[Z_{\Delta}^l(x, t)]$

1 *start to decrease monotonically and fast with l before the maximizer l_0 .*

2 **Part 3** *Finally, we perform the whole multilevel MC procedure. Given a squared norm of the sampling error $\epsilon_{\text{sampling}}^2$*
 3 *(see (30)), and given the first level l_0 obtained in Part 2, we obtain the optimal $N_{l_0}(x, t)$ and $M_{l_0}(x, t)$ for*
 4 *$l = l_0, \dots, N - 1$, using (32) and (33) from the constrained minimization problem. We set $N_{l_0}(t)$ and $M_{l_0}(t)$ as*
 5 *the ceiling of the maximum on x lying in the density domain of interest. Thus, the number of simulations is an*
 6 *integer independent of x .*

7 *Once the number of simulations is fixed, we estimate $\mathbb{E}[Z^{l_0}(x, t)]$ and $\mathbb{E}[Z_{\Delta}^l(x, t)]$, $l = l_0, \dots, N - 1$, using*
 8 *standard MC simulation. The output discretized function $f_X^{N, M}(x, t)$ is (29), which estimates $f_{X^{N(t)}}(x)$ with*
 9 *root mean square error $\|f_{X^{N(t)}}(x) - f_X^{N, M}(x, t)\|_2 \leq \epsilon_{\text{sampling}}$, for all x .*

10 In general, the squared norm of the sampling error $\epsilon_{\text{sampling}}^2$ and the order of truncation N are chosen as follows.
 11 At the beginning, one can estimate the bias error of the density approximation: given a very large order of truncation
 12 $N_{\infty} \gg 1$, and taking into account that $Z^{N_{\infty}}(x, t) \approx Z(x, t)$, the bias error at order $1 \leq N \ll N_{\infty}$, $|f_{X^{N(t)}}(x) -$
 13 $f_{X^{N_{\infty}(t)}}(x)|$, is estimated as

$$\left| \widehat{\mathbb{E}} [Z^{N_{\infty}}(x, t) - Z^N(x, t)] \right|. \quad (39)$$

14 The expectation is estimated using standard MC simulation. Suppose we fix a target global error δ . We split the target
 15 error as the sum of the bias error and the sampling error: $\delta = \epsilon_{\text{bias}} + \epsilon_{\text{sampling}}$. We take N such that (39) is less than
 16 ϵ_{bias} for all x . Then we apply the multilevel MC procedure with squared norm of the sampling error $\epsilon_{\text{sampling}}^2$. Keep in
 17 mind that the error δ cannot be as small as we want, since the sampling error $\epsilon_{\text{sampling}}$ must be achievable.

18 To apply the multilevel MC procedure, we need N such that the maximum (38) is greater than 1. This holds for
 19 large N for sure, according to the theoretical discussion. Thus, in practice, we can choose ϵ_{bias} small to make N large,
 20 and then run the multilevel MC algorithm with $\epsilon_{\text{sampling}} = \delta - \epsilon_{\text{bias}}$. In general, the decomposition of δ into ϵ_{bias} and
 21 $\epsilon_{\text{sampling}}$ is made heuristically, depending on the size of N for which (38) is greater than 1.

22 We showcase the multilevel MC method in numerical experiments. The first example deals with (1) having
 23 infinite series expansions in the input data; we observe that (38) is greater than 1 for $N \geq 1$, therefore multilevel MC
 24 is always superior to standard MC. In the second example, (38) is greater than 1 only for large N , so to take advantage
 25 of multilevel MC the error δ is chosen small, so that N is large enough.

26 These examples highlight what we believe is a general fact. Given N , multilevel MC outperforms crude MC
 27 simulation when $R_{l_0, N}(t) > 1$ (see (37) and (38)). But for a certain N , it may be possible that $R_{l_1, N}(t) \leq 1$ for
 28 $1 \leq l_1 < N$ (this is the case of Example 7 for $N < 36$). For such an N and the corresponding bias error, MC is more

1 efficient than the multilevel approach. When $A(t)$ and $B(t)$ are given by infinite random terms (Example 6), then c_l ,
 2 defined as the cost of one realization of $Z^l(x, t)$, grows as l^2 , so expression (38) is more inclined to be greater than 1
 3 for small N and higher tolerances (because the numerator of (37) grows quadratically). But when $A(t)$ and $B(t)$ are
 4 random polynomials (Example 7), then $c_l = l$, so a larger N and lower tolerances are required to ensure that (38) is
 5 greater than 1 (assuming that computational resources are available to achieve such an N).

6 **Example 6.** Consider (1) with $t_0 = 0$, $Y_0, Y_1 \sim \text{Normal}(2, 3)$, $A(t) = \sum_{n=0}^{\infty} \frac{U_n}{n!} t^n$ and $B(t) = t$, where U_0, U_1, \dots
 7 are independent random variables with distribution $\text{Beta}(11, 15)$. The series defining $A(t)$ converges in $L^\infty(\Omega)$ for
 8 all $t \in \mathbb{R}$. There is a unique analytic stochastic process $X(t)$ that solves (1) in the $L^p(\Omega)$ -sense, $1 \leq p < \infty$.

9 We estimate $\mathbb{V}[Z_\Delta^l(x, t)]$ for $N \leq 20$, using $m_V = 10,000$ realizations. With costs $c_l = C_l = l^2$, we estimate
 10 the ratios of complexities (37) for $l_1 \in [1, 19]$, $N \in [l_1 + 1, 20]$. In Figure 17 we present the ratios. Observe that they
 11 are greater than 1, therefore indicating higher efficiency of multilevel MC compared to crude MC for any level of
 12 tolerance. For each N , we keep the level l_0 maximizing the ratio, see (38). For $N \geq 7$, the same maximizer $l_0 = 6$ is
 13 obtained.




FIG. 17: Ratios of complexities between standard MC and multilevel MC (see (37)). The points indicate the maximum ratio with l_1 per truncation N . This figure corresponds to Example 6.

14 Figure 18 reports the ratios for $N > l_0 = 6$. The ratios grow roughly quadratically with N , with growth constant
 15 $1/l_0^2$. This is because the complexity of standard MC increases quadratically with N , while the cost of multilevel
 16 MC becomes constant from a certain level N . Small oscillations may be observed because of the finite number of
 17 simulations m_V used to estimate the variances.

18 Once we have the parameters of the multilevel MC strategy, we aim at approximating pointwise the probability
 19 density function of $X(t)$ at $t = 1.5$, $f_{X(t=1.5)}(x)$. Figure 19 reports in log-scale the estimated bias errors (39) with N




FIG. 18: Ratios of complexities between standard MC and multilevel MC (see (37)), for $N > l_0 = 6$. This figure corresponds to Example 6.

1 (the maximum in x of (39) has been taken). We fix $N_\infty = 100$ and consider $N \in [1, 20]$. The estimates are computed
 2 with standard MC simulation, using 20,000 realizations. Exponential decrease of the bias error is perceived, since
 3 it presents a concave and decreasing shape in log-scale, disregarding the oscillations due to the finite number of
 4 realizations. Further, the exponential decay of the bias error was rigorously proved in [25, Remark 3.8].




FIG. 19: Estimated bias errors with N in log-scale. This figure corresponds to Example 6.

5 Suppose that we wish to approximate $f_{X(t=1.5)}(x)$ uniformly with theoretical root mean square error $\delta > 0$. Let
 6 $\epsilon = \epsilon_{\text{bias}} = \epsilon_{\text{sampling}} = \delta/2$. We pick the first $N = N(\epsilon)$ for which the estimated bias error is less than ϵ . Multilevel
 7 MC can be applied with $l_0 = 6$, that $N(\epsilon)$, and sampling error $\epsilon_{\text{sampling}}^2 = \epsilon^2$. In Figure 20, we illustrate the relation
 8 between ϵ and the complexities of multilevel and standard MC, with maximum level $N = N(\epsilon)$. The following
 9 approximate relations hold:

$$W_{N(\epsilon)}^{\text{MC}} \propto \epsilon^{-2} N(\epsilon)^2, \quad W_{l_0, N(\epsilon)}^{\text{multi}} \propto \epsilon^{-2} l_0^2 \quad (40)$$

1 (the symbol \propto denotes proportionality to a constant). The cost of multilevel MC is reduced by factor $l_0^2/N(\epsilon)^2$. In
 2 log-log scale, the complexity of multilevel MC increases linearly with respect to ϵ^{-1} , with slope 2. On the contrary,
 3 for standard MC, such relation is shifted up with $2 \log N(\epsilon)$. The sequence $N(\epsilon)$ increases very slowly with $\epsilon \rightarrow 0$,
 4 because of the fast decrease of the bias error towards 0. But as $\epsilon \rightarrow 0$, multilevel MC becomes more and more
 5 efficient compared with standard MC.




FIG. 20: Sampling error versus complexity (see (35)–(36)), in log-log scale. The right plot is a zoom. This figure corresponds to Example 6.

6 Let us approximate $f_{X(t=1.5)}(x)$ uniformly with theoretical root mean square error $\delta = 0.001$. Let $\epsilon = \epsilon_{\text{bias}} =$
 7 $\epsilon_{\text{sampling}} = \delta/2 = 0.0005$. We pick the first N for which the estimated bias error is less than ϵ : $N = N(\epsilon) = 9$.
 8 Then we apply the multilevel MC procedure with sampling error ϵ^2 . The optimal number of simulations for $N = 9$
 9 is the following: $M = (N_6; M_6, M_7, M_8) = (19507; 1787, 861, 395)$. As expected, most of the cost is localized
 10 at level 6, while for the increments the cost goes down. The complexity of multilevel MC is 6.1×10^5 , which is
 11 smaller than that of standard MC, 1.2×10^6 . In Figure 21, left panel, we plot the estimate $\hat{f}_{X^{N=9}(t=1.5)}(x)$. It satisfies
 12 $\|f_{X(1.5)}(x) - \hat{f}_{X^9(1.5)}(x)\|_2 \leq \delta$, for all $x \in \mathbb{R}$. In the right plot, we show the estimated densities for $N = 8, 9$ and
 13 10, with root mean square norm of the sampling error $\epsilon = 0.0005$. Observe that the densities overlap, thus indicating
 14 the expected pointwise convergence.

15 **Example 7.** We study problem (16) with one degree polynomial coefficients. The probability distributions for the
 16 random inputs are $A_0 = 4$, $A_1 \sim \text{Uniform}(-3, 1)$, $B_0 \sim \text{Gamma}(2, 2)|_{[0,4]}$, $B_1 \sim \text{Bernoulli}(0.35)$ and $Y_0, Y_1 \sim$
 17 $\text{Normal}(2, \sigma = 3)$, all them independent. Due to the boundedness of A_0, A_1, B_0 and B_1 , this problem has a unique
 18 analytic stochastic solution $X(t)$, $t \in \mathbb{R}$, in the sense of the $L^p(\Omega)$ random calculus, for $1 \leq p < \infty$.

19 The objective is to approximate the probability density function $f_{X(t)}(x)$ using the multilevel MC approach, by
 20 estimating $f_{X^N(t)}(x)$. The theoretical results guarantee that $\{f_{X^N(t)}(x)\}_{N \geq 0}$ converges to $f_{X(t)}(x)$ pointwise and in




FIG. 21: Left plot: estimate $\hat{f}_{X^{N=9}(t=1.5)}(x)$. Right plot: estimates $\hat{f}_{X^N(t=1.5)}(x)$, for $N = 8, 9, 10$. This figure corresponds to Example 6.

1 $L^p(\mathbb{R})$, $1 \leq p < \infty$, as $N \rightarrow \infty$. We work at time $t = 1.5$.

2 We estimate $\mathbb{V}[Z_{\Delta}^l(x, t)]$ for N large (till $N = 70$), using parallel computing and $m_{\mathbb{V}} = 50,000$ realizations.
 3 The number of realizations $m_{\mathbb{V}}$ is actually excessively large, but we want to be quite exact in this example to show
 4 accurate complexities.

5 With costs $c_l = C_l = l$, we obtain the ratios of complexities between standard MC and multilevel MC, see (37).
 6 In Figure 22, we vary the starting level $l_1 \in [1, 69]$ and the maximum level $N \in [l_1 + 1, 70]$ and plot the corresponding
 7 ratios. Observe that the maximums (38) are greater than 1 (indicating that multilevel MC has higher efficiency) for
 8 $N \geq 36$ only, in contrast to Example 6. This means that for $N < 36$ and not sufficiently low tolerances crude MC
 9 simulation is more efficient. For each N , we keep l_0 maximizing the ratio, see (38). For $N = 36$, the maximizer is
 10 $l_0 = 35$; for $N = 37$ and 38 , the maximizer is $l_0 = 36$. For all $N \geq 39$, it is obtained the same $l_0 = 38$ as the
 11 maximizer.

12 In Figure 23 we report the ratios for $N > l_0 = 38$. The ratios are greater than 1 and, as N increases, they
 13 tend to infinity at linear rate approximately, although with small slope $\approx 1/l_0$. The complexity of standard MC grows
 14 linearly to infinity with N , while the complexity of multilevel MC becomes constant from a certain N . The oscillating
 15 behavior of the depicted ratios is due to the sampling error driven by the previous MC estimates for the variances.
 16 When $m_{\mathbb{V}}$ gets larger, the oscillations are mitigated and more exact results are obtained.

17 Let us approximate $f_{X(t=1.5)}(x)$ uniformly with root mean square error $\delta = 0.0001$. Figure 24 shows the esti-
 18 mated bias errors (39), with $N_{\infty} = 200$ and $20,000$ MC simulations (the maximum in x of (39) has been taken). We
 19 denote the bias errors as ϵ_{bias}^N (we make the dependence on N explicit). The bias error needs quite large truncation
 20 order N to start decreasing (up to $N = 25$ the bias is 0.1 approximately), and for $N \geq 35$ it drops abruptly. The




FIG. 22: Ratios of complexities between standard MC and multilevel MC (see (37)). The points indicate the maximum ratio with l_1 per truncation N . The transparent horizontal surface represents the ratio 1. This figure corresponds to Example 7.




FIG. 23: Ratios of complexities between standard MC and multilevel MC (see (37)), for $N > l_0 = 38$. This figure corresponds to Example 7.

1 bias error is less than δ only for $N \geq 37$, therefore we are in the region where multilevel MC is more efficient than
 2 standard MC. Notice that, in this example, we have fixed δ small on purpose to take advantage of the multilevel MC
 3 strategy. In the previous Example 6, we did not worry about δ nor its decomposition because multilevel MC showed
 4 more efficient than standard MC from the very first truncation order.




FIG. 24: Estimated bias errors with N in log-scale. This figure corresponds to Example 7.

5 In Figure 25 we plot, with $N \geq 37$, the complexities of the multilevel and the standard MC approaches with
 6 $\epsilon_{\text{sampling}} = \delta - \epsilon_{\text{bias}}^N$ ($\delta = 0.0001$), using (35)–(36). We observe that the complexity of multilevel MC is smaller, as
 7 expected. With formulas, we have

$$W_N^{\text{MC}}(\epsilon_{\text{sampling}}) \propto (\delta - \epsilon_{\text{bias}}^N)^{-2} N, \quad W_{l_0, N}^{\text{multi}}(\epsilon_{\text{sampling}}) \propto (\delta - \epsilon_{\text{bias}}^N)^{-2} l_0.$$

8 When $N \approx l_0$, we have $\delta - \epsilon_{\text{bias}}^N \approx 0$, so the complexities are large. When N is large, then $\delta - \epsilon_{\text{bias}} \approx \delta$, so
 9 $W_N^{\text{MC}}(\epsilon_{\text{sampling}}) \propto N \nearrow \infty$ and $W_{l_0, N}^{\text{multi}}(\epsilon_{\text{sampling}}) \searrow \text{constant}$.

10 Given $\delta = 0.0001$, in the case of standard MC the minimum complexity is attained when $N = 43$ (bias $\epsilon_{\text{bias}}^{N=43} =$
 11 5×10^{-7}), this being 2×10^8 .

12 For $N = 50 > l_0 = 38$, we run the multilevel MC procedure with sampling error $\epsilon_{\text{sampling}}^2 = (\delta - \epsilon_{\text{bias}}^{N=50})^2$. We
 13 choose $N = 50$ because the corresponding cost of multilevel MC is uniformly cheaper than standard MC applied
 14 with any order of truncation. Also, the complexity of multilevel MC becomes constant for $N \geq 50$. For $N = 50$, the
 15 optimal number of simulations is $M = (N_{38}; M_{38}, \dots, M_{50}) = (4527935; 12447, 4977, 2624, 1507, 475, 292, 126,$
 16 $45, 22, 9, 4, 2)$, with a complexity that is 1.7×10^8 . Figure 26, left panel, plots the graph of the estimated density
 17 function $\hat{f}_{X^{N=50}(t=1.5)}(x)$, which approximates the true density $f_{X(t=1.5)}(x)$ with root mean square error less than δ ,
 18 for all $x \in \mathbb{R}$. The right plot depicts the density estimates for $N = 49, 50$ and 51 , showing the expected overlapping.




FIG. 25: Complexities of the multilevel and the standard MC approaches with $\epsilon_{\text{sampling}} = \delta - \epsilon_{\text{bias}}^N$, $\delta = 0.0001$ fixed, using (35)–(36). This figure corresponds to Example 7.




FIG. 26: Left plot: estimate $\hat{f}_{X^{N=50}(t=1.5)}(x)$. Right plot: estimates $\hat{f}_{X^N(t=1.5)}(x)$, for $N = 49, 50, 51$. This figure corresponds to Example 7.

1 **Remark 2.** *Our application of the multilevel MC strategy is based on $L^\infty(\mathbb{R})$ norms. First, we acted pointwise in x*
 2 *in the constrained minimization problem from Proposition 5, and we took the $L^\infty(\mathbb{R})$ norm of the optimal number of*
 3 *simulations. Then the target density function was approximated uniformly in x . Second, the ratio (37) of complexities*
 4 *considered the $L^\infty(\mathbb{R})$ norms of the standard and the multilevel MC approaches.*

5 *Other norms can be considered. For instance, we could minimize the cost (31) given a fixed $L^1(\mathbb{R})$ norm of the*
 6 *sampling error variance (30): $\int_{\mathbb{R}} \|f_{X^N(t)}(x) - f_X^{N,M}(x, t)\|_2^2 dx = \epsilon_{\text{sampling}}^2$. In this case, we substitute each variance*
 7 *in x by its integral on \mathbb{R} . Then the approach becomes independent of x from the beginning.*

8 *When the cost is minimized pointwise in x , another possibility consists in considering the ratio of the $L^1(\mathbb{R})$*
 9 *norms of the complexities:*

$$W_N^{MC, \|\cdot\|_1}(t) = \epsilon_{\text{sampling}}^{-2} c_N \int_{\mathbb{R}} \mathbb{V}[Z^N(x, t)] dx, \quad (41)$$

$$W_{l_1, N}^{multi, \|\cdot\|_1}(t) = \epsilon_{\text{sampling}}^{-2} \int_{\mathbb{R}} \left(\sqrt{\mathbb{V}[Z^{l_1}(x, t)] c_{l_1}} + \sum_{l=l_1}^{N-1} \sqrt{\mathbb{V}[Z_\Delta^l(x, t)] C_l} \right)^2 dx, \quad (42)$$

$$R_{l_1, N}^{\|\cdot\|_1}(t) = \frac{W_N^{MC, \|\cdot\|_1}(t)}{W_{l_1, N}^{multi, \|\cdot\|_1}(t)} \quad (43)$$

12 *(the ratio is independent of $\epsilon_{\text{sampling}}$). In the two examples from this section, the conclusions derived are analogous*
 13 *considering $L^1(\mathbb{R})$ norms, thus showing robustness.*

14 *We present results regarding Example 6. Using $m_{\nabla} = 10,000$ realizations and costs $c_l = C_l = l^2$, we estimate*
 15 *the ratios of complexities (43) for $l_1 \in [1, 19]$ and $N \in [l_1 + 1, 20]$. Figure 27, left panel, shows the ratios. These*
 16 *ratios show a similar pattern to Figure 17. For each N , we keep the level l_0 maximizing the ratio (43). For $N \geq 7$,*
 17 *the same maximizer $l_0 = 6$ is obtained, as in Example 6. The right panel of Figure 27 compares the two ratios (37)*
 18 *and (43) for $N \geq l_0 = 6$. The two ratios are very similar, only showing a small discrepancy in part due to the MC*
 19 *procedure with m_{∇} realizations. The fact that the two ratios are practically coinciding is justified theoretically by the*
 20 *approximation $R_{l_1, N}^{\|\cdot\|_1}(t) \approx N^2/l_1^2$, which also held for (37).*

21 *Figure 28, left panel, analyzes theoretical error against complexity (see (41)–(42)). It corresponds to Figure 20*
 22 *by considering $L^1(\mathbb{R})$ norms, instead. When changing the norm, the “line” corresponding to error versus complexity*
 23 *in log-log scale is shifted up or down but has the same slope. This is because changing the norm only modifies the*
 24 *proportionality constants in (40). This is illustrated in Figure 28, right panel, where the two “lines” corresponding*
 25 *to multilevel MC are parallel, and also for standard MC.*




FIG. 27: Left panel: ratios of the $L^1(\mathbb{R})$ norms of the complexities between standard and multilevel MC (see (43)), where the points indicate the maximum ratio with l_1 per truncation N . Right panel: ratios of complexities between standard and multilevel MC with $L^\infty(\mathbb{R})$ norms (see (37)) and $L^1(\mathbb{R})$ norms (see (43)), for $N > l_0 = 6$. This figure corresponds to Remark 2.




FIG. 28: Left: sampling error $\epsilon = \epsilon_{\text{sampling}}$ versus $L^1(\mathbb{R})$ norm of the complexity (see (41)–(42)), in log-log scale. Right: sampling error $\epsilon = \epsilon_{\text{sampling}}$ versus the $L^1(\mathbb{R})$ (see (41)–(42)) and $L^\infty(\mathbb{R})$ (see (35)–(36)) norms of the complexity, in log-log scale. This figure corresponds to Remark 2.

5. CONCLUSIONS

This work focused on the improvement of the computational methods and algorithms from the recent contribution [25]. In [25], the authors studied the random second order linear differential equation with analytic data processes and random initial conditions, and constructed approximations to the probability density function of the stochastic solution based on dimensionality reduction and standard MC sampling strategy. The main goal in this paper is to speed up the convergence of the standard MC algorithm used in [25] by controlling the variance of the statistical estimators, thus avoiding noisy features in the density estimates.

The path-wise selection of the initial condition used in the density expression leads to a stable algorithm with good convergence. It was proved theoretically that the statistical estimators constructed with this method always possess finite variance. The implementation of the algorithm was illustrated through simple autonomous and non-autonomous linear differential equations.

Improvements are achieved by computing the expectation with respect to the initial conditions via numerical integration methods. By considering the densities of the initial conditions as integration weights, we focused on Gaussian quadrature rules. The orthogonal polynomials were considered with respect to standard probability distributions. Several conclusions are reached from this approach. When the initial conditions are comparable with the input coefficients of the equation in terms of random variability, the quadrature rule for integration allows significant variance reduction of the statistical estimators. Even when the initial conditions have small dispersion compared to the input coefficients, if the cost of each realization of the fundamental set is quite expensive, the quadrature rule yields similar expense to a plain MC strategy. The quadrature degree has to be selected sufficiently large to assure error-free approximations of the integrals.

A multilevel version of the MC algorithm allows for reduction of the complexity, by linear or quadratic factor. The multilevel MC approach turns out to be useful when high accuracy in the density approximation is required. The starting level was selected on the fly to maximize the ratio of complexities between the standard and the multilevel MC approach. Once the initial level is selected, and given a sampling error variance, we obtained the optimal number of simulations per level to minimize the complexity. The numerical results illustrated the methodology step-by-step in order to show the improvement with regard to the standard version of the MC simulation algorithm.

Several fundamental extensions and algorithmic improvements may be conceived, especially concerning the multilevel MC methodology. Firstly, the three parts in which we divided the multilevel procedure (variance of the increments, selection of the initial level by maximizing the ratio, and approximation of the density) could be combined in an automated algorithm, which would obtain the bias error, the sampling error, the variances, the initial level and the

1 optimal number of simulations on the fly. Secondly, the multilevel MC approach may be combined with a numerical
2 integration scheme for the expectation with respect to the initial conditions. Finally, our multilevel strategy relied
3 on considering consecutive levels, while other strategies could utilize different sequences of levels, such as linear or
4 geometric ones.

5 An interesting area to explore is the extension of the presently methodology to linear systems of second order
6 random differential equations. In this case, the aim is to approximate the joint probability density function of the
7 vector-valued stochastic solution. With the exception of simple autonomous linear systems whose density function can
8 be computed through transformations of random variables/vectors and direct integration, no analysis has been carried
9 out for non-autonomous systems. The application to non-autonomous systems would offer promising advances in the
10 active field of random differential equations.

11 **ACKNOWLEDGEMENTS**

12 This work is supported by Spanish “Ministerio de Economía y Competitividad” grant MTM2017–89664–P. M. Jor-
13 net acknowledges the doctorate scholarship granted by PAID, as well as “Ayudas movilidad de estudiantes de doc-
14 torado de la Universitat Politècnica de València para estancias en 2019”, for financing his research stay at CMAP.
15 J. Calatayud acknowledges “Fundació Ferran Sunyer i Balaguer”, “Institut d’Estudis Catalans” and the award from
16 “Borses Ferran Sunyer i Balaguer 2019” for funding her research stay at CMAP. All authors are grateful to Inria
17 (Centre de Saclay, DeFi Team), which hosted M. Jornet and J. Calatayud during their research stays at École Poly-
18 technique. The authors thank the reviewers for the valuable comments and suggestions, which have greatly enriched
19 the quality of the paper.

20 **CONFLICT OF INTEREST STATEMENT**

21 The authors declare that there is no conflict of interests regarding the publication of this article.

22 **APPENDIX: PROOFS**

23 **Proofs for Section 2.**

24 **Lemma 6.** *Let U be an absolutely continuous random variable, (Z_1, Z_2) be a random vector such that $Z_1 \neq 0$ almost*
25 *surely, and $G \in \mathcal{F}$ be an event of non-zero probability. Suppose that U and $(Z_1, Z_2, 1_G)$ are independent. Then the*

1 *probability law of $Z_1U + Z_2$ conditioned to G is absolutely continuous, with density function*

$$f_{Z_1U+Z_2}(z|G) = \mathbb{E} \left[f_U \left(\frac{z - Z_2}{Z_1} \right) \frac{1}{|Z_1|} \middle| G \right].$$

2 *Proof.* On the measurable space (Ω, \mathcal{F}) , define the conditional probability measure \mathbb{P}_G as $\mathbb{P}_G[E] = \mathbb{P}[E|G] :=$
3 $\mathbb{P}[E \cap G]/\mathbb{P}[G]$, for $E \in \mathcal{F}$.

4 Since U and G are independent, the probability law of U with respect to \mathbb{P}_G coincides with \mathbb{P} . Then the law of
5 U under \mathbb{P}_G is absolutely continuous, and $f_U(u|G) = f_U(u)$.

6 On the other hand, notice that U and (Z_1, Z_2) are independent under \mathbb{P}_G .

7 Finally, \mathbb{P}_G is absolutely continuous with respect to \mathbb{P} , $\mathbb{P}_G \ll \mathbb{P}$, therefore $Z_1 \neq 0$ \mathbb{P}_G -almost surely.

8 Thus, on the probability space $(\Omega, \mathcal{F}, \mathbb{P}_G)$ we are in conditions of applying [25, Lemma 3.2]:

$$f_{Z_1U+Z_2}(z|G) = \mathbb{E} \left[f_U \left(\frac{z - Z_2}{Z_1} \middle| G \right) \frac{1}{|Z_1|} \middle| G \right] = \mathbb{E} \left[f_U \left(\frac{z - Z_2}{Z_1} \right) \frac{1}{|Z_1|} \middle| G \right].$$

9

□

10 *Proof of Proposition 1.* We focus on the equality $f_{X(t)}(x) = \mathbb{E}[Z(x, t)]$, as the other one concerning $f_{X^N(t)}(x)$ is
11 analogous.

12 If either $\mathbb{P}[G_0(t)] = 0$ or $\mathbb{P}[G_1(t)] = 0$, then $Z(x, t) = Z_1(x, t)$ almost surely or $Z(x, t) = Z_0(x, t)$ almost
13 surely, respectively, so we are done.

14 Let us then assume that both $G_0(t)$ and $G_1(t)$ have non-zero probability. By the law of total probability, $f_{X(t)}(x) =$
15 $f_{X(t)}(x|G_0(t))\mathbb{P}[G_0(t)] + f_{X(t)}(x|G_1(t))\mathbb{P}[G_1(t)]$. To compute the conditional densities, we use Lemma 6:

$$f_{X(t)}(x|G_0(t)) = \mathbb{E}[Z_0(x, t)|G_0(t)], \quad f_{X(t)}(x|G_1(t)) = \mathbb{E}[Z_1(x, t)|G_1(t)].$$

16 As a consequence, $f_{X(t)}(x) = \mathbb{E}[Z(x, t)]$. □

17 *Proof of Lemma 2.* Fix $T > t_0$ and $t \in [t_0, T]$. Consider the Wronskian $W_{S_0, S_1}(t) = S_0(t)\dot{S}_1(t) - S_1(t)\dot{S}_0(t)$, where
18 the derivatives are regarded in the $L^\infty(\Omega)$ sense. Recall that $S_0(t)$ and $S_1(t)$ solve the differential equation in the
19 $L^\infty(\Omega)$ sense, see Section 1. In particular, the sample paths of $S_0(t)$ and $S_1(t)$ solve the deterministic version of the
20 differential equation, for almost every $\omega \in \Omega$. By Liouville's formula for deterministic linear differential equations
21 [38, Prop. 2.15], $W_{S_0, S_1}(t) = e^{-\int_{t_0}^t A(s) ds}$ almost surely. As $A(t)$ is a power series in $L^\infty(\Omega)$, we can lower bound

1 the Wronskian as follows:

$$W_{S_0, S_1}(t) \geq e^{-\int_{t_0}^T \|A(s)\|_\infty ds} =: \beta > 0,$$

almost surely. On the other hand, using the triangular inequality,

$$\begin{aligned} W_{S_0, S_1}(t) &\leq 2 \max\{|S_0(t)|, |S_1(t)|\} \max\{|\dot{S}_0(t)|, |\dot{S}_1(t)|\} \\ &\leq 2 \max\{|S_0(t)|, |S_1(t)|\} \max\{\|\dot{S}_0(t)\|_\infty, \|\dot{S}_1(t)\|_\infty\} \\ &\leq 2 \max\{|S_0(t)|, |S_1(t)|\} \gamma, \end{aligned}$$

2 where γ is a constant upper bound for $\|\dot{S}_0(t)\|_\infty$ and $\|\dot{S}_1(t)\|_\infty$ on $[t_0, T]$. As a consequence,

$$\max\{|S_0(t)|, |S_1(t)|\} \geq \frac{\beta}{2\gamma} =: 2\alpha,$$

3 almost surely, and (13) is proved.

4 Finally, we have that $\max\{|S_0^N(t)|, |S_1^N(t)|\}$ converges to $\max\{|S_0(t)|, |S_1(t)|\}$ as $N \rightarrow \infty$ in $L^\infty(\Omega)$, uni-
5 formly on $[t_0, T]$. This fact allows for deducing that there exists an integer $N_{T, \alpha} \geq 0$ such that (14) holds. \square

Proof of Proposition 3. Observe that, by (11) and (12),

$$|Z^N(x, t)| \leq \max\{\|f_{Y_0}\|_\infty, \|f_{Y_1}\|_\infty\} \left(\frac{1}{|S_0^N(t)|} \mathbf{1}_{G_0^N(t)} + \frac{1}{|S_1^N(t)|} \mathbf{1}_{G_1^N(t)} \right) = \frac{\max\{\|f_{Y_0}\|_\infty, \|f_{Y_1}\|_\infty\}}{\max\{|S_0^N(t)|, |S_1^N(t)|\}},$$

6 for $N \leq \infty$. Then Lemma 2 applies. \square

7 Proofs for Section 4.

Proof of Proposition 4. First, by Proposition 1, $\mathbb{E}[Z^{l_0}(x, t)] = f_{X^{l_0}(t)}(x)$. On the other hand,

$$\begin{aligned} \mathbb{E}[Z_\Delta^l(x, t)] &= \mathbb{E}[\Delta Z_0^l(x, t) \mathbf{1}_{H_0^l(t)}] + \mathbb{E}[\Delta Z_1^l(x, t) \mathbf{1}_{H_1^l(t)}] \\ &= \mathbb{E}[\Delta Z_0^l(x, t) | H_0^l(t)] \mathbb{P}[H_0^l(t)] + \mathbb{E}[\Delta Z_1^l(x, t) | H_1^l(t)] \mathbb{P}[H_1^l(t)] \\ &= \mathbb{E}[Z_0^{l+1}(x, t) | H_0^l(t)] \mathbb{P}[H_0^l(t)] - \mathbb{E}[Z_0^l(x, t) | H_0^l(t)] \mathbb{P}[H_0^l(t)] \\ &\quad + \mathbb{E}[Z_1^{l+1}(x, t) | H_1^l(t)] \mathbb{P}[H_1^l(t)] - \mathbb{E}[Z_1^l(x, t) | H_1^l(t)] \mathbb{P}[H_1^l(t)]. \end{aligned}$$

By Lemma 6 and the law of total probability,

$$\begin{aligned}\mathbb{E}[Z_{\Delta}^l(x, t)] &= f_{X^{l+1}(t)}(x | H_0^l(t)) \mathbb{P}[H_0^l(t)] - f_{X^l(t)}(x | H_0^l(t)) \mathbb{P}[H_0^l(t)] \\ &\quad + f_{X^{l+1}(t)}(x | H_1^l(t)) \mathbb{P}[H_1^l(t)] - f_{X^l(t)}(x | H_1^l(t)) \mathbb{P}[H_1^l(t)] \\ &= f_{X^{l+1}(t)}(x) - f_{X^l(t)}(x).\end{aligned}$$

1 The proposition follows from telescopic sum identities. □

2 REFERENCES

- 3 1. Soong, T.T., *Random Differential Equations in Science and Engineering*, New York: Academic Press, 1973.
- 4 2. Neckel, T. and Rupp, F., *Random Differential Equations in Scientific Computing*, London: Walter de Gruyter, 2013.
- 5 3. Strand, J.L., Random ordinary differential equations, *J. Differ. Equ.*, **7**, pp. 538–553, 1970.
- 6 4. Smith, R.C., *Uncertainty Quantification. Theory, Implementation, and Application*, New York: SIAM Computational Science
7 & Engineering, 2014.
- 8 5. Cortés, J.-C., Jódar, L., Camacho, J. and Villafuerte, L., Random Airy type differential equations: Mean square exact and
9 numerical solutions, *Comput. Math. Appl.*, **60**, pp. 1237–1244, 2010.
- 10 6. Calbo, G., Cortés, J.-C. and Jódar, L., Random Hermite differential equations: mean square power series solutions and statis-
11 tical properties, *Appl. Math. Comput.*, **218**(7), pp. 3654–3666, 2011.
- 12 7. Calatayud, J., Cortés, J.-C. and Jornet, M., Improving the approximation of the first- and second-order statistics of the response
13 stochastic process to the random Legendre differential equation, *Mediterr. J. Math.*, **16**(68), pp. 1–14, 2019.
- 14 8. Calatayud, J., Cortés, J.-C. and Jornet, M., Computational uncertainty quantification for random non-autonomous second
15 order linear differential equations via adapted gPC: a comparative case study with random Fröbenius method and Monte Carlo
16 simulation, *Open Math.*, **16**, pp. 1651–1666, 2018.
- 17 9. Xiu, D., *Numerical Methods for Stochastic Computations. A Spectral Method Approach*, New York: Cambridge Texts in
18 Applied Mathematics, Princeton University Press, 2010.
- 19 10. Golmankhaneh, A.A., Porghoveh, N.A. and Baleanu, D., Mean square solutions of second-order random differential equations
20 by using homotopy analysis method, *Rom. Rep. Phys.*, **65**, pp. 350–362, 2013.
- 21 11. Khudair, A.R., Ameen, A.A. and Khalaf, S.L., Mean square solutions of second-order random differential equations by using
22 variational iteration method, *Appl. Math. Sci.*, **5**, pp. 2505–2519, 2011.
- 23 12. Khudair, A.R., Ameen, A.A. and Khalaf, S.L., Mean square solutions of second-order random differential equations by using

- 1 Adomian decomposition method, *Appl. Math. Sci.*, **5**, pp. 2521–2535, 2011.
- 2 13. Khudair, A.K., Haddad, S.A.M. and Khalaf, S.L., Mean square solutions of second-order random differential equations by
3 using the differential transformation method, *Open Journal of Applied Sciences*, **6**, pp. 287–297, 2016.
- 4 14. Villafuerte, L. and Chen-Charpentier, B.M., A random differential transform method: Theory and applications, *Appl. Math.*
5 *Lett.*, **25**, pp. 1490–1494, 2012.
- 6 15. Calatayud, J., Cortés, J.-C., Jornet, M. and Villafuerte, L., Random non-autonomous second order linear differential equations:
7 mean square analytic solutions and their statistical properties, *Adv. Differ. Equ.*, **2018**(392), pp. 1–29, 2018.
- 8 16. Calatayud, J., Cortés, J.-C. and Jornet, M., Some notes to extend the study on random non-autonomous second order linear
9 differential equations appearing in mathematical modeling, *Mathematical and Computational Applications*, **23**(4), pp. 76–89,
10 2018.
- 11 17. Cortés, J.-C., Navarro-Quiles, A., Romero, J.-V. and Roselló, M.-D., Probabilistic solution of random autonomous first-order
12 linear systems of ordinary differential equations, *Rom. Rep. Phys.*, **68**(4), pp. 1397–1406, 2016.
- 13 18. Calatayud, J., Chen-Charpentier, B.M., Cortés, J.-C. and Jornet, M., Combining polynomial chaos expansions and the random
14 variable transformation technique to approximate the density function of stochastic problems, including some epidemiological
15 models, *Symmetry*, **11**(1), pp. 43–60, 2019.
- 16 19. El-Tawil, M.A., The approximate solutions of some stochastic differential equations using transformations, *Appl. Math. Com-*
17 *put.*, **164**(1), pp. 167–178, 2005.
- 18 20. Dorini, F.A., Ceconello, M.S. and Dorini, M.B., On the logistic equation subject to uncertainties in the environmental carrying
19 capacity and initial population density, *Commun. Nonlinear Sci. Numer. Simul.* **33**, pp. 160–173, 2016.
- 20 21. Santos, L.T., Dorini, F.A. and Cunha, M.C.C., The probability density function to the random linear transport equation, *Appl.*
21 *Math. Comput.*, **216**(5), pp. 1524–1530, 2010.
- 22 22. Calatayud, J., Cortés, J.-C. and Jornet, M., The damped pendulum random differential equation: A comprehensive stochastic
23 analysis via the computation of the probability density function, *Physica A*, **512**, 261–279, 2018.
- 24 23. Hussein, A. and Selim, M.M., Solution of the stochastic radiative transfer equation with Rayleigh scattering using RVT
25 technique, *Appl. Math. Comput.*, **218**(13), pp. 7193–7203, 2012.
- 26 24. Casabán, M.C., Cortés, J.-C., Romero, J.V. and Roselló, M.D., Solving random homogeneous linear second-order differential
27 equations: a full probabilistic description, *Mediterr. J. Math.*, **13**(6), pp. 3817–3836, 2016.
- 28 25. Jornet, M., Calatayud, J., Le Maître, O.P. and Cortés, J.-C., Second order linear differential equations with analytic uncer-
29 tainties: Stochastic analysis via the computation of the probability density function, *J. Comput. Appl. Math.*, **374**(112770), pp.
30 1–20, 2020.
- 31 26. Wolfram Research, Inc., *Mathematica, Version 11.2*, Champaign, IL, USA: Wolfram Research, Inc., 2017.

- 1 27. Giles, M.B., Multilevel Monte Carlo path simulation, *Oper. Res.*, **56**(3), pp. 607–617, 2008.
- 2 28. Giles, M.B., Multilevel Monte Carlo methods, *Acta Numer.*, **24**, pp. 259–328, 2015.
- 3 29. Cliffe, A., Giles, M.B., Scheichl, R. and Teckentrup, A., Multilevel Monte Carlo Methods and applications to elliptic PDEs
4 with random coefficients, *Computing and Visualization in Science*, **14**(1), pp. 3–15, 2011.
- 5 30. Pisoni, M., Nobile, F.B. and Leyland, P., A Continuation Multi Level Monte Carlo method for uncertainty quantification in
6 compressible inviscid aerodynamics, *Comput. Method. Appl. M.*, **326**(C): pp. 20–50, 2017.
- 7 31. Beskos, A., Jasra, A., Law, K., Tempone, R. and Zhou, Y., Multilevel sequential Monte Carlo samplers, *Stoch. Proc. Appl.*,
8 **127**(5), pp. 1417–1440, 2017.
- 9 32. Sobol, I.M., Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates, *Math. Comput.*
10 *Simulat.*, **55**(1–3), pp. 271–280, 2001.
- 11 33. Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M. and Tarantola, S., *Global Sensitivity*
12 *Analysis, The Primer*, England: John Wiley & Sons, 2008.
- 13 34. Saltelli, A., Annoni, P., Azzini, I., Campolongo, F., Ratto, M. and Tarantola, S., Variance based sensitivity analysis of model
14 output. Design and estimator for the total sensitivity index, *Comput. Phys. Commun.*, **181**(2), pp. 259–270, 2010.
- 15 35. Sobol', I.M., Tarantola, S., Gatelli, D., Kucherenko, S. and Mauntz, W., Estimating the approximation error when fixing
16 unessential factors in global sensitivity analysis, *Reliab. Eng. Syst. Safe.*, **92**(7), pp. 957–960, 2007.
- 17 36. Efron, B. and Tibshirani, R., *An Introduction to the Bootstrap*, New York: CRC Press, 1994.
- 18 37. Davison, A.C. and Hinkley D.V., *Bootstrap Methods and Their Application*, London: Cambridge University Press, 1997.
- 19 38. Chicone, C., *Ordinary Differential Equations With Applications*, New York: Springer Science+Business Media, 2006.