

ÍNDICE GENERAL

INTRODUCCIÓN GENERAL.....	1
1. El manejo integrado de plagas.....	3
2. Semioquímicos en el control de plagas.....	6
2.1 Los semioquímicos.....	6
2.2 Técnicas de identificación e investigación de semioquímicos.....	8
2.2.1 Bioensayos electrofisiológicos.....	9
2.2.2 Bioensayos de comportamiento.....	10
2.3 Técnicas de control de plagas basadas en semioquímicos.....	11
2.3.1 Control indirecto.....	12
2.3.2 Control directo.....	12
3. La confusión sexual.....	15
3.1 Descripción de la técnica.....	15
3.2 Dispositivos emisores de feromona.....	17
3.2.1 Características del emisor ideal.....	17
3.2.2 Dispositivos y formulaciones comerciales utilizados.....	19
3.3 Mecanismos que producen la confusión sexual.....	21
3.3.1 Mecanismos competitivos.....	21
3.3.2 Mecanismos no competitivos.....	22
4. Medida de semioquímicos en aire.....	24
4.1 Importancia de la medida de semioquímicos en aire.....	24
4.2 Medida de compuestos orgánicos volátiles en aire.....	25
4.3 Dificultad de la medida de feromona en aire.....	26
4.4 Antecedentes de cuantificación de feromonas en aire.....	27
4.5 Cuantificación de volátiles mediante la captación de muestras de aire.....	28
4.5.1 Captación de la muestra de aire.....	28

4.5.1.1	Técnicas de captación.....	28
4.5.1.2	Materiales o dispositivos para la toma de muestra.....	30
4.5.2	Extracción de la muestra.....	32
4.5.2.1	Extracción mediante disolventes.....	32
4.5.2.2	Extracción sin uso de disolventes.....	33
4.5.3	Concentración y limpieza de la muestra.....	33
4.5.4	Identificación y cuantificación del analito contenido en la muestra.....	34
5.	La polilla del racimo de la vid (<i>Lobesia botrana</i>).....	35
5.1	Importancia y daños.....	35
5.2	Distribución geográfica.....	35
5.3	Sistemática.....	37
5.4	Morfología.....	37
5.5	Ciclo biológico.....	39
5.6	Control de la plaga.....	40
5.6.1	Los primeros métodos de lucha.....	40
5.6.2	Control químico.....	40
5.6.3	Control biológico.....	41
5.6.4	Control biotécnico.....	42
6.	El barrenador del arroz (<i>Chilo suppressalis</i>).....	44
6.1	Importancia.....	44
6.2	Distribución geográfica.....	44
6.3	Sistemática.....	45
6.4	Morfología.....	46
6.5	Ciclo biológico y daños.....	47

6.6 Control de la plaga.....	48
6.6.1 Control cultural.....	49
6.6.2 Control químico.....	49
6.6.3 Control biológico.....	49
6.6.4 Control biotécnico.....	50
OBJETIVOS Y JUSTIFICACIÓN.....	53
CAPÍTULO 1 “Airborne pheromone quantification in treated vineyards with different mating disruption dispensers against <i>Lobesia botrana</i>”.....	59
C1.1 Introduction.....	62
C1.2 Material and methods.....	66
C1.2.1 Field Locations and Mating Disruption Systems.....	66
C1.2.2 Airborne Pheromone Quantification.....	67
C1.2.2.1 Chemicals and Reagents.....	67
C1.2.2.2 Heptyl 4,4,5,5,6,6,7,7,8,8,9,9,9-Tridecafluorononanoate (TFN) synthesis.....	68
C1.2.2.3 Field Air Sampling.....	68
C1.2.2.4 Sample Extraction and Filtration.....	69
C1.2.2.5 Pheromone Quantification.....	69
C1.2.3 Adsorbed Pheromone on Leaves.....	71
C1.2.3.1. Adsorption Capacity and Emission Assays.	71
C1.2.3.2 Leaf Sample Extractions.....	72
C1.2.4 Release Profiles Studies of the Emission Systems.....	73
C1.2.5 Efficacy of the Treatments.....	73
C1.2.6 Data Analysis.....	74
C1.3. Results.....	75
C1.3.1. Airborne Pheromone Quantification.....	75
C1.3.2. Pheromone Leaves Adsorption Capacity.....	76

C1.3.2.1. <i>Experiment 1</i>	76
C1.3.2.2. <i>Experiment 2</i>	77
C1.3.2.3 <i>Experiment 3</i>	78
C1.3.3 <i>Dispensers Pheromone Release Profile</i>	79
C1.3.3.1. <i>Passive Dispensers' Release Profile</i>	79
C1.3.3.2 <i>Aerosol Devices Profile</i>	81
C1.3.4 <i>Population Monitoring</i>	82
C1.3.5 <i>Crop Damage Prospections</i>	83
C1.4 Discussion	84
CAPÍTULO 2 “ <i>Studies on Lobesia botrana non-competitive disruption mechanisms based on male’s response after pheromone exposure</i> ”.....	91
C2.1 Introduction	94
C2.2 Material and methods	97
C2.2.1 <i>Insects</i>	97
C2.2.2 <i>Moths pheromone exposure</i>	97
C2.2.3 <i>Airborne pheromone quantification for exposure treatments</i>	99
C2.2.3.1 <i>Chemicals and reagents</i>	99
C2.2.3.2 <i>Quantification of the airborne pheromone in the conditioning chamber</i>	100
C2.2.4 <i>Wind tunnel tests</i>	101
C2.2.5 <i>Electroantennography assays</i>	102
C2.2.6 <i>Data análisis</i>	103
C2.3 Results	104
C2.3.1 <i>Airborne pheromone concentrations in the exposure treatments</i>	104
C2.3.2. <i>Wind tunnel assays</i>	104
C2.3.2.1 <i>Immediately after the exposure</i>	104
C2.3.2.2 <i>Twenty-four hours after the exposure</i>	106

C2.3.3 EAG assays.....	107
C2.3.3.1 <i>Immediately after the exposure</i>	107
C2.3.3.2 <i>Twenty-four hours after the exposure</i>	109
C2.4 Discussion	111
CAPÍTULO 3 “Influence of climate conditions in <i>Lobesia botrana</i> mating disruption dispensers’ performance”	117
C3.1 Introduction	120
C3.2 Material and methods	122
C3.2.1 Mating disruption field trials and pheromone dispensers.....	122
C3.2.2 Climate data.....	123
C3.2.3 Dispenser release kinetic studies.....	123
C3.2.4 Population monitoring.....	124
C3.2.5 Damage assessments.....	124
C3.2.6 Statistical analysis.....	125
C3.3 Results	126
C3.3.1 Population monitoring.....	126
C3.3.2 Damage assessments.....	127
C3.3.3 Dispensers release kinetics.....	129
C3.3.4 Dispensers climate dependency.....	133
C3.4 Discussion	135
CAPÍTULO 4 “Importance of the early deployment of mating disruption dispensers against the striped rice stem borer assessed with airborne pheromone variations along the crop cycle”.....	139
C4.1 Introduction	142
C4.2 Material and Methods	144
C4.2.1 Pheromone dispensers.....	144
C4.2.2 Experimental design.....	144

C4.2.3 Efficacy of the treatments.....	146
C4.2.4 Pheromone release profiles.....	146
C4.2.5 Airborne Pheromone Quantification.....	147
C4.2.5.1 <i>Field Air Sampling</i>	147
C4.2.5.2 <i>Smple extraction and filtration</i>	148
C4.2.5.3 <i>Pheromone quantification</i>	148
C4.2.6 Data analysis.....	150
C4.3 Results	151
C4.3.1 Efficacy of the tratments.....	151
C4.3.2 Dispenser release profiles.....	152
C4.3.3 Airborne pheromone quantification.....	154
C4.4 Discussion	155
DISCUSIÓN GENERAL	161
CONCLUSIONES	173
BIBLIOGRAFÍA	179
ANEJO METODOLÓGICO: Desarrollo de una metodología para la cuantificación de feromona en aire	206

ÍNDICE DE FIGURAS

INTRODUCCIÓN

Figura 1 Tendencias en la investigación en el campo de la entomología aplicada en la revista “Journal of Economic Entomology”, 1927-1957.....	4
Figura 2. Macho y hembra de <i>Lasiocampa quercus</i>	6
Figura 3 Equipo de electroantenografía.....	10
Figura 4 Túnel de viento.....	11
Figura 5 Trampa utilizada para el seguimiento del vuelo de machos de <i>Lobesia botrana</i>	12
Figura 6 Trampa funnel de trapeo masivo y trampa de atracción y muerte.....	13
Figura 7 Emisor automático de pulverización utilizado contra <i>Lobesia botrana</i> , Emisor pasivo utilizado contra <i>Lobesia botrana</i> , emisor pasivo a bajas densidades utilizado contra <i>Chilo suppressalis</i>	19
Figura 8 Presencia y daños generados por <i>L. botrana</i> en el racimo.....	35
Figura 9 Distribución mundial de <i>Lobesia botrana</i>	36
Figura 10 Huevo, Larva y adulto de <i>Lobesia botrana</i>	38
Figura 11. Componente mayoritario de la feromona de <i>Lobesia botrana</i> , acetato de (E,Z)-7,9-dodecadienilo.....	42
Figura 12 Diferentes tipos de emisores utilizados para la confusión sexual de la polilla del racimo (De izquierda a derecha: Quant Lb Pro, Isonet L, el dispositivo automático de pulverización Checkmate Puffer Lb y Lobetec).....	43
Figura 13 Distribución mundial de <i>Chilo suppressalis</i>	45
Figura 14 Puesta, Larva y adulto de <i>Chilo suppressalis</i>	47
Figura 15 Síntomas de daño de barrenador en la parte superior de la planta. Larva de barrenador en el interior del tallo.....	48

Figura 16 Componentes de la feromona de <i>Chilo suppressalis</i>	50
Figura 17 Emisor biodegradable para el control del barrenador. Trampa fúnel utilizada para el seguimiento del barrenador.....	51

CAPÍTULO 1

Figure C1.1 Mean absolute airborne pheromone concentrations in vineyards treated with passive dispensers and aerosol devices.....	76
Figure C1.2 Mean quantity of pheromone adsorbed on leaf samples, after one aerosol emission.....	77
Figure C1.3 Mean quantity of pheromone adsorbed on leaf samples, after overnight aerosol shooting.....	78
Figure C1.4 Mean quantity of pheromone adsorbed on leave samples collected in actual field conditions.....	79
Figure C1.5 Pheromone release profile of Isonet L passive dispensers.	80
Figure C1.6 Pheromone release profile of Checkmate aerosol devices.	81
Figure C1.7 Mean of <i>L. botrana</i> male captures per trap per day throughout the season in the plots with different treatments.....	83

CAPÍTULO 2

Figure C2.1 Assembly scheme of the conditioning setup to expose male moths at different airborne pheromone amounts.....	98
Figure C2.2 Percentage of male moths that showed activation, wing fanning, take off zig-zag flight and reached the source immediately after being exposed to the different treatments.....	105
Figure C2.3 Percentage of male moths that showed activation, wing fanning, take-off, zig-zag flight and reached the source, 24 h after their exposure to the different treatments.....	107
Figure C2.4 EAG signal triggered by the main component of the pheromone in the antenna of male moths immediately after their exposure to the different treatments.....	108
Figure C2.5 EAG signal triggered by 1-hexanol in the antenna of male moths immediately after their exposure to the different treatments.....	109
Figure C2.6 EAG signal triggered by the main component of the pheromone in the antenna of male moths 24 h after their exposure to the different exposure treatments.....	110

Figure C2.7 EAG signal triggered by 1-hexanol in the antenna of male moths 24 h after their exposure to the different exposure treatments.....	110
---	-----

CAPÍTULO 3

Figure C3.1 Mean <i>L. botrana</i> male captures per trap per day throughout the study period of the three trial replicates in the plots with different treatments.....	127
Figure C3.2 Mean percentage of damaged bunches obtained in each treatment in the three damage assessments.....	128
Figure C3.3 Pheromone release profiles of Ampoule dispensers in the three trials.....	130
Figure C3.4. Pheromone release profiles of Rope dispensers in the three trials.....	131
Figure C3.5 Average temperatures (minimum, mean and maximum) and wind speed in the three trial locations of the assay.....	133
Figure C3.6 Mean residual pheromone content of each type of dispenser throughout the crop cycle and monthly average mean temperatures from the three trial replicates.....	135

CAPÍTULO 4

Figure C4.1 Arrangement of the different deployment strategies at each location.....	145
Figure C4.2 Mean <i>C. suppressalis</i> male captures per trap per day of the three trials throughout the crop season with the different mating disruption treatments.....	151
Figure C4.3 Residual pheromone content of Bioselibate CS dispensers installed in May and June.....	153
Figure C4.4 Mean absolute airborne pheromone concentrations in rice field treated with passive dispensers Bioselibate CS.....	155

ÍNDICE DE TABLAS

INTRODUCCIÓN

Tabla 1. Categorías de semioquímicos y ejemplos.....	7
---	---

CAPÍTULO 1

Table C1.1 Optimum values of the MS/MS parameters for each target compound used to quantify <i>L. botrana</i> airborne pheromone.....	70
--	----

Table C1.2 Isonet L passive dispensers aging. Residual means of pheromone content throughout the time in field.....	80
--	----

Table C1.3 Checkmate aerosol devices pheromone content. Residual means of pheromone content throughout the time in field.....	82
--	----

Table C1.4 Mean percentage of damaged bunches obtained in each treatment in the three damage assessments.....	84
--	----

CAPÍTULO 2

Table C2.1 Optimum values of the MS/MS parameters for each target compound used to quantify airborne pheromone in the conditioning chamber.....	101
--	-----

CAPÍTULO 3

Table C3.1 Initial pheromone content, total pheromone released per dispenser and per ha/day and percentage of residual pheromone at the end of the season.....	132
---	-----

CAPÍTULO 4

Table C4.1. Optimum values of the MS/MS parameters for each target compound to quantify <i>C. suppressalis</i> airborne pheromone.....	149
---	-----

Table C4.2 Mean (\pm SE) of infected plants in one square meter (NIP/m ²) and percentage of attacked plants (%) obtained in each treatment in the three assessments carried out at each trial.....	152
--	-----

Table C4.3 Bioselibate CS dispensers aging. Residual means of pheromone content (mg) throughout the time in field (days), Point values of Figure 2 and the total mean of pheromone used (mg/day/ha).	154
---	-----