

INDEX

Abstract	i
Resumen	iii
Resum	v
Chapter 1: Introduction	1
1.1 Geographical distribution and invasion history	3
1.2 Distinctive morphological and biological features	4
1.3 Biology of <i>Eutetranychus banksi</i>	6
1.4 Feeding damage, sampling, and thresholds	8
1.5 Biological control	10
1.6 Objectives	13
Chapter 2: The newcomer takes it all: the invader Texas citrus mite, <i>Eutetranychus banksi</i> (Acari: Tetranychidae), displaces the resident relatives on citrus agrosystems	15
2.1 Introduction.....	18
2.2 Materials and Methods.....	20
2.3 Results.....	27
2.4 Discussion.....	38
2.5 Acknowledgments.....	46
2.6 Supplementary Materials	47
Chapter 3: Within-tree distribution and seasonal dynamics of <i>Eutetranychus banksi</i> (Acari: Tetranychidae) and <i>Euseius stipulatus</i> (Acari: Phytoseiidae) on citrus: Implications for the biological control of the pest	61
3.1 Introduction.....	64
3.2 Materials and Methods.....	65
3.3 Results.....	69
3.4 Discussion.....	82

3.5 Acknowledgments.....	93
Chapter 4: Population structure, spatial distribution, and sampling plan for the Texas citrus mite, <i>Eutetranychus banksi</i> (Acari: Tetranychidae) on citrus	95
4.1 Introduction.....	98
4.2 Materials and Methods.....	99
4.3 Results.....	104
4.4 Discussion.....	116
4.5 Acknowledgments.....	124
Chapter 5: Life-history, prey-stage preference, functional and numerical response of the predatory mite <i>Neoseiulus madeirensis</i> (Acari: Phytoseiidae) to <i>Eutetranychus banksi</i> (Acari: Tetranychidae) on citrus.....	125
5.1 Introduction.....	128
5.2 Materials and Methods.....	129
5.3 Results.....	138
5.4 Discussion.....	146
5.5 Acknowledgments.....	154
Chapter 6: General discussion	155
Chapter 7: Conclusions	167
Chapter 8: References	173

Figure Index

Chapter 1: Introduction1

Figure 1.1 *Eutetranychus banksi*, a. Deutonymph, male and exuvia next to the midrib. b. Protonymph, larva, female and eggs close to midrib. In the images it is possible to distinguish the feeding damage caused by *E. banksi* on the leaf's surface5

Figure 1.2 Morphological features to separate *Eutetranychus* species in Mediterranean citrus crops. Dorsal (a) and ventral (b) surface of *Eutetranychus* showing the naming of dorsal setae6

Figure 1.3 Morphological features to separate *Eutetranychus* species in Mediterranean citrus crops. *Eutetranychus banksi* show (a) a trapezoid pattern between the third (e1) and fourth (f1) dorsocentral hysterosomal setae, and (b) two ventral setae in the coxa II. *Eutetranychus orientalis* show (c) a square pattern between the same dorsocentral hysterosomal setae, and (d) one ventral setae in coxa I.....7

Figure 1.4 Dense colonies of *E. banksi* on clementine citrus. a, b. fruit colony and c. leaf colony where all stages of development are feeding. In addition, stippling damage can be observed, as well as the characteristic residues of accumulated moult visible at high populations.....9

Figure 1.5 Damaged fruit and leaves by *E. banksi*. a. Unripe sweet orange tree fruit, the left fruit shows its normal colouring while the right one is damaged. b. Ripe orange fruit, the upper fruit row shows damage on the sunny side, while the lower row shows normal colouring. c. Clementine citrus leaves discoloured and with the “silvery” appearance observed after severe *E. banksi* infestation.....10

Chapter 2: The newcomer takes it all: the invader Texas citrus mite, *Eutetranychus banksi* (Acari: Tetranychidae), displaces the resident relatives on citrus agrosystems15

Figure 2.1 Location of sampling sites in the Valencian region (Spain). Legend: Pink corresponds to sites sampled during 2017, Blue is sampled sites in 2018, Yellow 2019, and Green 2020. Map was obtained using *Q-sig*21

Figure 2.2 Patterns of species occurrence across the sampled orchards. Species occur more frequently singly than together with other species. a. Proportion of occupied

orchards by occurrence (i.e., number of species detected). b. Proportion of occupied orchards by species: *E. banksi*, *E. orientalis*, and *P. citri*. c. Area-proportional Euler diagram illustrating the number of orchards occupied by each species, and by multiple species (indicated in areas of overlap)28

Figure 2.3 Proportion of orchards with occurrence of each species (framed), when another one occurs (grey legend) or is absent (white legend). a. *E. orientalis* vs *E. banksi*, b. *P. citri* vs *E. banksi*, c. *P. citri* vs *E. orientalis*. For each pair, significant differences are denoted with asterisks. Chi square contingency test: *P < 0.05; **P < 0.0130

Figure 2.4 Kernel density distribution of the three mite species in the Valencian region (Spain). Kernel density estimation (KDE) was computed with the optimal bandwidth using likelihood cross-validation, and a Gaussian function to assess the probability density of the three species per square kilometre in raster maps with 3284 x 1890 pixel array and 0.1 Km² area per pixel. The yellow areas represent high mite densities while dark violet colour indicates low mite densities. The white dots indicate the citrus orchards sampled with the occurrence of each species. Maps were performed using the R-package *spatstat*.....32

Figure 2.5 Bar plots (a–c) showing the mean kernel density (KDE) with the standard errors (capped bars), grouped by presence or absence between species. Bars with different letters are significantly different according to Mann-Whitney *U*-test. Representation of the binomial (logit-link) generalized linear models (GLMs) (d–g) on the probability of mite presence depending on KDE of the potential competitor. In the graphs, the prediction line is indicated in blue, and the 95% confidence band in grey. GLMs with statistically non-significant relationships have not been represented33

Figure 2.6 a. Principal component analysis (PCA) showing all the sampled orchards grouped according to the mite occurrence and kernel density estimation (KDE) values. b. Ordination plot of the two axes of redundancy analysis (RDA) constrained by the selected environmental variables (variable codes are in table 2.1). Graphs were performed using the R-package *vegan*.....35

Chapter 3: Within-tree distribution and seasonal dynamics of *Eutetranychus banksi* (Acari: Tetranychidae) and *Euseius stipulatus* (Acari: Phytoseiidae) on citrus: Implications for the biological control of the pest61

Figure 3.1 Mean number of *E. banksi* (a–d) and phytoseiid mites (e–h) per leaf or per cm² of leaves and fruits. Bars with different letters are significantly different according to Wilcoxon *rank sum* test.70

Figure 3.2 Seasonal trends of *E. banksi* and phytoseiid mites on leaves (solid line, first y-axis) and fruits (broken line, second y-axis) in four citrus orchards in 2018. Mean number of mites collected per sampling unit (all the stages were pooled together). Note that first and second y-axis scales are different. Mean (solid line), maximum and minimum daily temperatures in °C (broken lines) and mean daily relative humidity (RH) were represented. Capped bars represent ± standard error (SE).72

Figure 3.3 Seasonal trends of *E. banksi* and phytoseiid mites on leaves (solid line, first y-axis) and fruits (broken line, second y-axis) in six citrus orchards in 2019. Mean number of mites collected per sampling unit (all the stages were pooled together). The first and second y-axis scales are different. Mean (solid line), maximum and minimum daily temperatures in °C (broken lines) and mean daily relative humidity (RH) were represented. Capped bars represent ± standard error (SE).73

Figure 3.4 Seasonal relative abundance of motile forms of phytoseiid species in four (2018) and six (2019) citrus orchards. Percentage of each species per sampling is represented. The summer decline of *E. stipulatus* is represented by arrows in the graph.75

Figure 3.5 Variation in the spatial distribution and body coloration of phytoseiids in relation to the abundance of *E. banksi* in four (2018) and six (2019) citrus orchards. Percentage of phytoseiids in each position in the tree canopy (outside/inside), leaf (adaxial/abaxial), fruits (occupied/unoccupied by phytoseiids), and different body coloration (white/red) (primary y-axis), compared with the mean number of *E. banksi* per fruit or leaf (secondary, y-axis). Capped bars represent ± standard error (SE).77

Figure 3.6 (a–d) Representation of the binomial (logit-link) generalized linear models (GLMs) showing the relationship between the proportion of leaves occupied by phytoseiids outside the canopy (canopy), on the adaxial side of the leaves (leaf), the occupation on leaves by red-coloured phytoseiids (colour), and the proportion of fruits

occupied by phytoseiids (fruit) (y-axis), compared with the mean number of *E. banksi* per fruit or leaf (x-axis). In the graphs the prediction lines were plotted in black and the 95% confidence bands in grey. (e–h) Representation of the linear models (LM) showing the relationship between the mean number of phytoseiids per sample unit on the outside canopy, on the adaxial leaf side, red-coloured, and collected on fruits (y-axis), compared with the mean number *E. banksi* per leaf or fruit (x-axis). Prediction lines and confidence bands were plotted in the same way as on the GLMs78

Figure 3.7 Proportion of leaves occupied (grey legend) or unoccupied (white legend) by phytoseiids outside the canopy (a), on adaxial side of leaves (b), red coloured phytoseiids on leaves (c), and phytoseiid on fruits (d), when *E. banksi* occurred or was absent. Significant differences are denoted with asterisks. Chi square contingency test: ***P < 0.001.80

Figure 3.8 Phytoseiid proportion per leaf outside or inside the canopy (a), on the adaxial or abaxial side of the leaves (b), white or red coloured (c), and collected on fruits (d), when *E. banksi* occurred or was absent. Capped bars represent ± standard error (SE). Significant differences are denoted with asterisks. Chi-square contingency test: ***P < 0.001.81

Chapter 4: Population structure, spatial distribution, and sampling plan for the Texas citrus mite, *Eutetranychus banksi* (Acari: Tetranychidae) on citrus95

Figure 4.1 a. Seasonal relative abundance of developmental stages and population trends of *E. banksi* on citrus leaves and fruits. Percentage of each developmental stage per sample unit (fruit or leaf) and date are plotted in colored areas, and the number of mites per sample unit by a dotted line. b. Seasonal sex ratio and population trends of *E. banksi* on leaves and fruits. Sex ratio per sampling unit (fruit or leaf) and date are represented by solid lines, while the number of mites per sample unit by a dotted line. The peaks of eggs (a) and females (b) have been highlighted with ordinal numbers.107

Figure 4.2 Representation of the linear models (LMs) showing the relationship between the mean motile developmental stages (a) or mean females (b) (x-axis) and the mean total mites (y-axis) of *E. banksi* per leaf in the periphery of the canopy. In the

graphs, the prediction line is represented in black and the 95% confidence bands in grey. The goodness of fit of the model was assessed by computing the r^2 value.....111

Figure 4.3 Representation of the linear models (LM) showing the relationship between mean density (x-axis) and variance (y-axis) of motile stages of *E. banksi* pooled together (gray) and adult females (black) on citrus leaves in the periphery of the tree canopy. The goodness of fit of the models was assessed by computing the r^2 value.112

Figure 4.4 Enumerative and binomial sampling for *E. banksi* motile mites (a) and females (b) on citrus. Number of leaves from the periphery of the canopy required (y-axis) based on the mean motile mites or mean females per leaf (x-axis). Green's method was used for enumerative (solid line) and Kuno's method for binomial sampling (dashed line), to achieve a precision level of $D = 0.25$113

Figure 4.5 Relationship between the proportion of leaves on the periphery of the canopy infested with *E. banksi* motile forms (a) and females (b) (y-axis) and the mean number of motile mites or females per sample unit (x-axis). Lines represent predicted values from the NBD Wilson & Room approach, while points represent the observed values in the field.115

Figure 4.6 Autumn leaf with females coming from the spring and summer leaves. In the picture more than 20 females can be seen at the first stages of leaf colonization, while the number of females per leaf ranges from 1 to 10 in already established populations.....118

Figure 4.7 A dense *E. banksi* colony where all stages of development can be observed. There is a large accumulation of molts and hatched eggs, as well as discoloration on the leaf surface because of feeding.121

Chapter 5: Life-history, prey-stage preference, functional and numerical response of the predatory mite *Neoseiulus madeirensis* (Acari: Phytoseiidae) to *Eutetranychus banksi* (Acari: Tetranychidae) on citrus.....125

Figure 5.1 a. *E. banksi* colony maintained on turgid castor been leaves for about a week in a plastic container with water. b. *E. banksi* rearing units on detached citrus leaves kept fresh using wet cotton wool.130

Figure 5.2 <i>Neoseiulus madeirensis</i> rearing units on detached bean leaves previously infested with <i>T. urticae</i> and kept fresh using wet cotton wool. a. Dense colony of <i>N. madeirensis</i> where all stages of development are present. b. Gravid female of <i>N. madeirensis</i> walking on the web of <i>T. urticae</i>	131
Figure 5.3 a. Experimental arena consist of 3 x 3 cm square cells on orange leaves, placed inside a Petri dish with the underside facing upwards, and on moistened cotton wool and strips of cotton wool used to prevent individuals from escaping. b. Petri dishes inside the climate-controlled incubator and closed with a perforated lid covered with a fine mesh.....	132
Figure 5.4 Age-specific survival rate (l_x) (y-first axis) and age-specific fecundity rate (m_x) (y-secondary axis) curves of <i>N. madeirensis</i> fed on <i>E. banksi</i> . The age-specific sex-ratio is represented in bars (y-first axis).....	139
Figure 5.5 Prey-stage preferences according to mean $\beta \pm SE$ value when provided varying prey ratios (1:1, 1:2 or 2:1) for the three different immature prey combinations (Larva:Nymph, Egg:Nymph or Egg:Larva).	142
Figure 5.6 Functional and numerical responses of adult female <i>N. madeirensis</i> to different developmental stages of <i>E. banksi</i> . In functional response graphs, points represent the observed mean prey consumed at each initial prey density, and solid lines were predicted by the Holling's disc equation. In numerical response graphs, points represent the observed mean eggs laid at each initial prey density, and solid lines were predicted by the hyperbolic model. Dashed lines represent the 95% confidence intervals	143

Table Index

Chapter 1: Introduction	1
Table 1.1 Intervention thresholds and sample sizes for <i>E. banksi</i> infestations in Florida citrus.....	11
Chapter 2: The newcomer takes it all: the invader Texas citrus mite, <i>Eutetranychus banksi</i> (Acari: Tetranychidae), displaces the resident relatives on citrus agrosystems	15
Table 2.1 Meaning of the seventeen variables related to temperature, relative humidity, wind speed, precipitation, and crop density	24
Table 2.2 Results for the null model co-occurrence analysis. The analysis was based on the C-score and V-ratio index for sim 2 and sim 10 algorithms.....	29
Table 2.3 Results for the probabilistic model of co-occurrence analysis	31
Table 2.4 Results of the binomial (logit-link) generalized linear models (GLMs) on the probability of presence depending on the density kernel (KDE) for the three mite species	34
Table 2.5 (a) Permutational multivariate analysis of variance (PERMANOVA) and (b) multivariate dispersion analysis (BETADISPER) of kernel density (KDE) data, in relation to the different groups of orchards grouped by the mite occurrence and KDE values. (c) PERMANOVA pairwise comparison between groups of orchards.....	35
Table 2.6 Environmental variables between sampling sites for the three mite species. Variables are given as mean (mean plus standard deviation -SD-, minimum, and maximum values recorded. Variables are grouped by temperature, relative humidity (HUM), wind speed (WIN) and precipitation (PRE). The meaning for the codes of the environmental variables is available in Table 2.1.....	37
Table 2.7 Results from redundancy (RDAs) and variation partitioning analysis (VPA) corrected by Moran spectral randomisation.....	37

Chapter 3: Within-tree distribution and seasonal dynamics of *Eutetranychus banksi* (Acari: Tetranychidae) and *Euseius stipulatus* (Acari: Phytoseiidae) on citrus: Implications for the biological control of the pest61

Table 3.1 Sampled sites, years, varieties and pest management practices.....66

Table 3.2 Abundance, relative abundance, and frequency of phytoseiid mites. Abundance represents the number of individuals of each species collected; Relative abundance is the percentage of individuals of each species with respect to the total number of phytoseiids; Frequency is the percentage of samples in which a certain phytoseiid species occurred with respect to the total number of samples collected....74

Table 3.3 Results of the binomial (logit-link) generalized linear models (GLMs) on the occupancy by phytoseiids outside the canopy, on the adaxial side of the leaves, as well as the occupation on leaves by red-colour phytoseiids, and the occupancy by phytoseiids on fruits, depending on the mean number of *E. banksi* per sample unit. .79

Table 3.4 Results of the linear models (LM) on the mean number of phytoseiids per sample unit outside the canopy, on the adaxial leaf side, red-coloured, and collected on fruits, compared with the mean number of *E. banksi* per sample unit.....79

Chapter 4: Population structure, spatial distribution, and sampling plan for the Texas citrus mite, *Eutetranychus banksi* (Acari: Tetranychidae) on citrus95

Table 4.1 Total individuals collected of different developmental stages in each sampling unit (fruit, upper and lower leaf side) and leaf type. In brackets, the proportion of each stage of development in each sample unit and leaf type. The last row summarises the sex-ratio (females/ [males + females]). Significant differences are denoted with asterisks. Chi-square test: ***P < 0.001.105

Table 4.2 Taylor's power law parameters for the different developmental stages of *E. banksi* on different plant strata. Parameter values *a* and *b* used for the sampling plans are highlighted in bold.109

Table 4.3 Resampling simulation results used to validate Green's fixed precision sequential sampling plan for *E. banksi* motile developmental stages (a) and females (b) on citrus with desired precision levels adjusted to 0.25 with replacement (Motile stages' Taylor *a* = 8.35 and *b* = 1.53; Female's Taylor *a* = 3.04 and *b* = 1.27)116

Chapter 5: Life-history, prey-stage preference, functional and numerical response of the predatory mite *Neoseiulus madeirensis* (Acari: Phytoseiidae) to *Eutetranychus banksi* (Acari: Tetranychidae) on citrus.....125

Table 5.1 Development time in hours (mean \pm SE), and survival of immature stages of *N. madeirensis* when fed on *E. banksi*.....138

Table 5.2 Mean \pm SE duration of female adult phases, longevity, fecundity, and demography parameters of *N. madeirensis* feeding on *E. banksi*.....140

Table 5.3 Mean \pm SE number of prey consumed and relative consumption of *N. madeirensis* when offered 30 prey items of different developmental stages for 24 h.141

Table 5.4 Maximum likelihood estimates from binomial (logit-link) GLMs of the proportion of prey consumed as a function of initial *E. banksi* prey density by adult female *N. madeirensis*.....144

Table 5.5 Estimates of functional response parameters (a) by generalized non-linear least squares regression (GNLS) of Holling 's Type II model, and numerical response parameters (b) by non-linear least squares regression (NLS) of the Hyperbolic model.145

Table 5.6 *Neoseiulus madeirensis* female summary of prey consumption and egg-laying on the different prey stages of *E. banksi*.....145