

ÍNDICE

Pág.

❖ Resúmenes	1-6
❖ <u>CAPÍTULO 1: INTRODUCCIÓN, JUSTIFICACIÓN y</u> OBJETIVOS	
1.1. El fuego bacteriano	9-18
1.1.1. Sintomatología y rango de hospedadores	
1.1.2. Distribución mundial	
1.1.3. Distribución en España	
1.2. <i>Erwinia amylovora</i> , el agente causal	18-28
1.2.1. Taxonomía y características generales	
1.2.2. Factores de virulencia: los exopolisacáridos	
1.3. Epidemiología: ciclo biológico de <i>E. amylovora</i>	29-33
1.4. Diagnóstico de fuego bacteriano	33-46
1.4.1. Aislamiento de <i>E. amylovora</i>	
1.4.2. Detección serológica	
1.4.3. Detección molecular	
1.4.4. Identificación	
1.4.5. Protocolo de diagnóstico (PM7/20, EPPO)	

1.5. Una estrategia de supervivencia bacteriana: 47-63
el estado viable no cultivable (VNC)

- 1.5.1. Supervivencia, viabilidad y muerte celular
- 1.5.2. El estado VNC: determinación y características
- 1.5.3. Reversión del estado VNC: recuperación o resucitación
- 1.5.4. Mantenimiento del poder patógeno en las células VNC
- 1.5.5. El estado VNC en la naturaleza: implicaciones epidemiológicas
- 1.5.6. El estado VNC en bacterias asociadas a plantas

1.5. Control del fuego bacteriano 64-76

- 1.6.1. Medidas preventivas
- 1.6.2. Control químico: cobre, antibióticos y otros
- 1.6.3. Control biológico
- 1.6.4. Control genético

1.7. Justificación y Objetivos 77-78

❖ **CAPÍTULO 2: “Survival strategy of *Erwinia amylovora* against copper: induction of the viable-but-nonculturable state” (2006, Appl. Environ. Microbiol. 72: 3482-3488)**

2.1. Abstract	81
2.2. Introduction	82-84
2.3. Materials and Methods	84-88
2.3.1. Inoculation of <i>E. amylovora</i> cells in mineral medium with copper	
2.3.2. Bacterial cell counts	
2.3.3. Culturability restoration or resuscitation of copper-induced VBNC cells	
2.3.4. SEM of <i>E. amylovora</i> cells in the presence of copper	
2.3.5. Pathogenicity assays of VBNC and resuscitated <i>E. amylovora</i> cells	
2.3.6. Statistical analysis	
2.4. Results	89-95
2.4.1. Copper induces the VBNC state in <i>E. amylovora</i>	
2.4.2. Resuscitation of copper-induced VBNC <i>E. amylovora</i> cells	
2.4.3. SEM of copper-induced VBNC <i>E. amylovora</i> cells	
2.4.4. Pathogenicity of copper-induced VBNC and resuscitated <i>E. amylovora</i> cells	
2.5. Discussion	96-101
2.6. Acknowledgements	102

❖ **CAPÍTULO 3: “Survival of *Erwinia amylovora* in water”**

3.A. Long-term starvation-survival of *Erwinia amylovora* in sterile irrigation water. (2006, Acta Hort. 704: 107-112)

3.A.1. Abstract	105-106
3.A.2. Introduction	106-107
3.A.3. Materials and Methods	108-109
3.A.3.1. Bacterial strains and water microcosms	
3.A.3.2. <i>E. amylovora</i> cell counts	
3.A.3.3. Infectivity assays in green pears	
3.A.4. Results	109-112
3.A.4.1. Survival of <i>E. amylovora</i> in water	
3.A.4.2. Pathogenicity assays	
3.A.5. Discussion	113-115

3.B. *Erwinia amylovora* survives in natural water. (Acta Hort. *in press*)

3.B.1. Abstract	116-117
3.B.2. Introduction	117-118
3.B.3. Materials and Methods	118-119
3.B.3.1. Bacterial strains and preparation of water microcosms	
3.B.3.2. <i>E. amylovora</i> cell counts	
3.B.3.3. Pathogenicity of <i>E. amylovora</i> in natural water	

3.B.4. Results	119-122
3.B.4.1. Fate of <i>E. amylovora</i> in natural water microcosms	
3.A.4.2. Pathogenicity of <i>E. amylovora</i> in natural water microcosms	
3.B.5. Discussion	122-124
3.B.6. Acknowledgements	124

❖ **CAPÍTULO 4: “Improvement of the recovery of *Erwinia amylovora* stressed cells on RESC, a simple, rapid, and differential culture medium” (Int. Microbiol., to be submitted)**

4.1. Abstract	127
4.2. Introduction	128-130
4.3. Materials and Methods	131-138
4.3.1. Bacterial strains	
4.3.2. Selection of the copper sulphate concentration for adding to KB medium	
4.3.3. Copper complexing measurements of KB broth	
4.3.4. Growth media and incubation conditions	

- 4.3.5. Evaluation of RESC medium *in vitro*: enumeration of *E. amylovora* cells from mixed cultures with epiphytic plant bacteria
- 4.3.6. Evaluation of RESC medium *in vivo*: isolation of *E. amylovora* from field samples
- 4.3.7. Evaluation of RESC medium for the enumeration of *E. amylovora* stressed cells
- 4.3.8. Statistical analysis

4.4. Results 138-148

- 4.4.1. Copper complexing ability of KB broth
- 4.4.2. Selection of copper sulphate concentration: the RESC medium
- 4.4.3. Colonial morphology of *E. amylovora* on RESC medium
- 4.4.4. Recovery efficiency of *E. amylovora* on RESC medium *in vitro*
- 4.4.5. Recovery efficiency of *E. amylovora* on RESC medium *in vivo*
- 4.4.6. Recovery efficiency of stressed *E. amylovora* cells on RESC medium

4.5. Discussion 148-153

4.6. Acknowledgements 154

❖ **CAPÍTULO 5: “Role of amylovoran and levan on the survival of *Erwinia amylovora* under copper stress and starvation” (Appl. Environ. Microbiol., to be submitted)**

5.1. Abstract	157
5.2. Introduction	158-160
5.3. Materials and Methods	160-165
5.3.1. Bacterial strains and inoculation conditions	
5.3.2. Bacterial cell counts	
5.3.3. Direct and indirect EPS measurements	
5.3.4. Staining of EPS and bacterial morphology	
5.3.5. Quantification of copper-complexing ability of EPSs	
5.3.6. Effect of EPS extracts under copper and starvation stress conditions	
5.3.7. Statistical analysis	
5.4. Results	165-176
5.4.1. Influence of EPSs in the survival of <i>E. amylovora</i>	
5.4.2. Morphological changes in <i>E. amylovora</i> cells under stress conditions	
5.4.3. Copper complexing ability of <i>E. amylovora</i> cells and EPS extracts	
5.4.4. EPS extracts prolong the culturability of <i>E. amylovora</i> under stress conditions	
5.5. Discussion	176-182
5.6. Acknowledgements	183

❖ <u>CAPÍTULO 6: DISCUSIÓN GENERAL Y CONCLUSIONES</u>	
6.1. Discusión general	187-195
6.2. Conclusiones generales	195-197
❖ <u>CAPÍTULO 7: BIBLIOGRAFÍA GENERAL</u>	199-226