

RESEARCH

Open Access

# Fixed and periodic points of generalized contractions in metric spaces

Mujahid Abbas<sup>1</sup>, Basit Ali<sup>2</sup> and Salvador Romaguera<sup>3\*</sup>

\*Correspondence:

sromague@mat.upv.es

<sup>3</sup>Instituto Universitario de Matemática Pura y Aplicada, Universitat Politècnica de València, Valencia, 46022, Spain  
Full list of author information is available at the end of the article

## Abstract

Wardowski (*Fixed Point Theory Appl.* 2012:94, 2012, doi:10.1186/1687-1812-2012-94) introduced a new type of contraction called  $F$ -contraction and proved a fixed point result in complete metric spaces, which in turn generalizes the Banach contraction principle. The aim of this paper is to introduce  $F$ -contractions with respect to a self-mapping on a metric space and to obtain common fixed point results. Examples are provided to support results and concepts presented herein. As an application of our results, periodic point results for the  $F$ -contractions in metric spaces are proved.

**MSC:** 47H10; 47H07; 54H25

**Keywords:**  $F$ -contraction; property  $P$ ; property  $Q$ ; common fixed point

## 1 Introduction and preliminaries

The Banach contraction principle [1] is a popular tool in solving existence problems in many branches of mathematics (see, e.g., [2–4]). Extensions of this principle were obtained either by generalizing the domain of the mapping or by extending the contractive condition on the mappings [5–9]. Initially, existence of fixed points in ordered metric spaces was investigated and applied by Ran and Reurings [10]. Since then, a number of results have been proved in the framework of ordered metric spaces (see [11–18]). Contractive conditions involving a pair of mappings are further additions to the metric fixed point theory and its applications (for details, see [19–23]).

Recently, Wardowski [24] introduced a new contraction called  $F$ -contraction and proved a fixed point result as a generalization of the Banach contraction principle [1]. In this paper, we introduce an  $F$ -contraction with respect to a self-mapping on a metric space and obtain common fixed point results in an ordered metric space. In the last section, we give some results on periodic point properties of a mapping and a pair of mappings in a metric space. We begin with some basic known definitions and results which will be used in the sequel. Throughout this article,  $\mathbb{N}$ ,  $\mathbb{R}_+$ ,  $\mathbb{R}$  denote the set of natural numbers, the set of positive real numbers and the set of real numbers, respectively.

**Definition 1** Let  $f$  and  $g$  be self-mappings on a set  $X$ . If  $fx = gx = w$  for some  $x$  in  $X$ , then  $x$  is called a coincidence point of  $f$  and  $g$  and  $w$  is called a coincidence point of  $f$  and  $g$ . Furthermore, if  $fgx = gfx$  whenever  $x$  is a coincidence point of  $f$  and  $g$ , then  $f$  and  $g$  are called weakly compatible mappings [22].

Let  $C(f, g) = \{x \in X : fx = gx\}$  ( $F(f, g) = \{x \in X : x = fx = gx\}$ ) denote the set of all coincidence points (the set of all common fixed points) of self-mappings  $f$  and  $g$ .

**Definition 2** ([25]) Let  $(X, d)$  be a metric space and  $f, g : X \rightarrow X$ . The mapping  $f$  is called a  $g$ -contraction if there exists  $\alpha \in (0, 1)$  such that

$$d(fx, fy) \leq \alpha d(gx, gy)$$

holds for all  $x, y \in X$ .

In 1976, Jungck [25] obtained the following useful generalization of the Banach contraction principle.

**Theorem 1** Let  $g$  be a continuous self-mapping on a complete metric space  $(X, d)$ . Then  $g$  has a fixed point in  $X$  if and only if there exists a  $g$ -contraction mapping  $f : X \rightarrow X$  such that  $f$  commutes with  $g$  and  $g(X) \subseteq f(X)$ .

Let  $F$  be the collection of all mappings  $F : \mathbb{R}_+ \rightarrow \mathbb{R}$  that satisfy the following conditions:

(C1)  $F$  is strictly increasing, that is, for all  $\alpha, \beta \in \mathbb{R}_+$  such that  $\alpha < \beta$  implies that

$$F(\alpha) < F(\beta).$$

(C2) For every sequence  $\{\alpha_n\}_{n \in \mathbb{N}}$  of positive real numbers,  $\lim_{n \rightarrow \infty} \alpha_n = 0$  and

$$\lim_{n \rightarrow \infty} F(\alpha_n) = -\infty$$
 are equivalent.

(C3) There exists  $k \in (0, 1)$  such that

$$\lim_{\alpha \rightarrow 0^+} \alpha^k F(\alpha) = 0.$$

**Definition 3** ([24]) Let  $(X, d)$  be a metric space and  $F \in F$ . A mapping  $f : X \rightarrow X$  is said to be an  $F$ -contraction on  $X$  if there exists  $\tau > 0$  such that

$$d(fx, fy) > 0 \quad \text{implies that} \quad \tau + F(d(fx, fy)) \leq F(d(x, y)) \tag{1}$$

for all  $x, y \in X$ .

Note that every  $F$ -contraction is continuous (see [24]). We extend the above definition to two mappings.

**Definition 4** Let  $(X, d)$  be a metric space,  $F \in F$  and  $f, g : X \rightarrow X$ . The mapping  $f$  is said to be an  $F$ -contraction with respect to  $g$  on  $X$  if there exists  $\tau > 0$  such that

$$\tau + F(d(fx, fy)) \leq F(d(gx, gy)) \tag{2}$$

for all  $x, y \in X$  satisfying  $\min\{d(fx, fy), d(gx, gy)\} > 0$ .

By different choices of mappings  $F$  in (1) and (2), one obtains a variety of contractions [24].

**Example 1** Let  $F_1 : \mathbb{R}_+ \rightarrow \mathbb{R}$  be given by  $F_1(\alpha) = \ln(\alpha)$ . It is clear that  $F \in \mathcal{F}$ . Suppose that  $f : X \rightarrow X$  is an  $F$ -contraction with respect to a self-mapping  $g$  on  $X$ . From (2) we have

$$\tau + \ln(d(fx, fy)) \leq \ln(d(gx, gy)),$$

which implies that

$$d(fx, fy) \leq e^{-\tau} d(gx, gy).$$

Therefore an  $F_1$ -contraction map  $f$  with respect to  $g$  reduces to a  $g$ -contraction mapping.

Now we give an example of an  $F$ -contraction with respect to a self-mapping  $g$  on  $X$  which is not a  $g$ -contraction on  $X$ .

**Example 2** Consider the following sequence of partial sums  $\{S_n\}_{n \in \mathbb{N}}$  [24, Example 2.5]:

$$\begin{aligned} S_1 &= 1, \\ S_2 &= 1 + 2, \\ S_3 &= 1 + 2 + 3, \\ &\dots \\ S_n &= 1 + 2 + \dots + n = \frac{n(n+1)}{2}, \quad n \in \mathbb{N}. \end{aligned}$$

Let  $X = \{S_n : n \in \mathbb{N}\}$  and  $d$  be the usual metric on  $X$ . Let  $f : X \rightarrow X$  and  $g : X \rightarrow X$  be defined as

$$fS_n = \begin{cases} S_{n-1}, & \text{if } n > 1, \\ S_1, & \text{if } n = 1, \end{cases} \quad gS_n = \begin{cases} S_{n+1}, & \text{if } n > 1, \\ S_1, & \text{if } n = 1. \end{cases}$$

Let  $F_1 : \mathbb{R}_+ \rightarrow \mathbb{R}$  be given by  $F_1(\alpha) = \ln(\alpha)$ . As

$$\lim_{n \rightarrow \infty} \frac{d(fS_n, fS_1)}{d(gS_n, gS_1)} = \lim_{n \rightarrow \infty} \frac{S_{n-1} - S_1}{S_{n+1} - S_1} = 1,$$

so  $f$  is not a  $g$ -contraction. If we take  $F_2(\alpha) = \ln(\alpha) + \alpha$ , then  $F_2 \in \mathcal{F}$  and  $f$  is an  $F_2$ -contraction with respect to a mapping  $g$  (taking  $\tau = 2$ ). Indeed, the following holds:

$$\frac{d(fS_n, fS_1)}{d(gS_n, gS_1)} e^{d(fS_n, fS_1) - d(gS_n, gS_1)} = \frac{S_{n-1} - S_1}{S_{n+1} - S_1} e^{S_{n-1} - S_1 - S_{n+1} + S_1} = \frac{n^2 - n - 2}{n^2 + 3n} e^{-4n-2} \leq e^{-2}$$

for all  $n > 1$ . For all  $m, n \in \mathbb{N}$  with  $m > n > 1$ , we have

$$\begin{aligned} &\frac{d(fS_m, fS_n)}{d(gS_m, gS_n)} e^{d(fS_m, fS_n) - d(gS_m, gS_n)} \\ &= \frac{S_{m-1} - S_{n-1}}{S_{m+1} - S_{n+1}} e^{S_{m-1} - S_{n-1} - S_{m+1} + S_{n+1}} \\ &= \frac{m^2 + m - n^2 - n}{m^2 + 3m - n^2 - 3n} e^{-2(m-n)} \leq e^{-2}. \end{aligned}$$

**Definition 5** ([26], Dominance condition) Let  $(X, \preceq)$  be a partially ordered set. A self-mapping  $f$  on  $X$  is said to be (i) a dominated map if  $fx \preceq x$  for each  $x$  in  $X$ , (ii) a dominating map if  $x \preceq fx$  for each  $x$  in  $X$ .

**Example 3** Let  $X = [0, 1]$  be endowed with the usual ordering and  $f, g : X \rightarrow X$  defined by  $gx = x^n$  for some  $n \in \mathbb{N}$  and  $fx = kx$  for some real number  $k \geq 1$ . Note that

$$gx = x^n \leq x \quad \text{and} \quad x \leq kx = fx$$

for all  $x$  in  $X$ . Thus  $g$  is dominated and  $f$  is a dominating map.

**Definition 6** Let  $(X, \preceq)$  be a partially ordered set. Two mappings  $f, g : X \rightarrow X$  are said to be weakly increasing if  $fx \preceq gfx$  and  $gx \preceq fgx$  for all  $x$  in  $X$  (see [27]).

**Definition 7** Let  $X$  be a nonempty set. Then  $(X, d, \preceq)$  is called an ordered metric space if  $(X, d)$  is a metric space and  $(X, \preceq)$  is a partially ordered set.

**Definition 8** Let  $(X, \preceq)$  be a partial ordered set, then  $x, y$  in  $X$  are called comparable elements if either  $x \preceq y$  or  $y \preceq x$  holds true. Moreover, we define  $\Delta \subseteq X \times X$  by

$$\Delta = \{(x, y) \in X \times X : x \preceq y \text{ or } y \preceq x\}.$$

**Definition 9** An ordered metric space  $(X, d, \preceq)$  is said to have the sequential limit comparison property if for every non-decreasing sequence (non-increasing sequence)  $\{x_n\}_{n \in \mathbb{N}}$  in  $X$  such that  $x_n \rightarrow x$  implies that  $x_n \preceq x$  ( $x \preceq x_n$ ).

## 2 Common fixed point results in ordered metric spaces

We present the following theorem as a generalization of results in [25] and [24, Theorem 2.1].

**Theorem 2** Let  $(X, \preceq)$  be a partially ordered set such that there exists a metric  $d$  on  $X$ , and let  $f : X \rightarrow X$  be an  $F$ -contraction with respect to  $g : X \rightarrow X$  on  $\Delta$  with  $f(X) \subseteq g(X)$ . Assume that  $f$  is dominating and  $g$  is dominated. Then

- $f$  and  $g$  have a coincidence point in  $X$  provided that  $g(X)$  is complete and has the sequential limit comparison property.
- $C(f, g)$  is well ordered if and only if  $C(f, g)$  is a singleton.
- $f$  and  $g$  have a unique common fixed point if  $f$  and  $g$  are weakly compatible and  $C(f, g)$  is well ordered.

*Proof* (a) Let  $x_0$  be an arbitrary point of  $X$ . Since the range of  $g$  contains the range of  $f$ , there exists a point  $x_1$  in  $X$  such that  $fx_0 = gx_1$ . As  $f$  is dominating and  $g$  is dominated, so we have

$$x_0 \preceq fx_0 = gx_1 \preceq x_1.$$

Hence  $(x_0, x_1) \in \Delta$ . Continuing this process, having chosen  $x_n$  in  $X$ , we obtain  $x_{n+1}$  in  $X$  such that

$$x_n \preceq fx_n = gx_{n+1} \preceq x_{n+1}.$$

So, we obtain  $(x_n, x_{n+1}) \in \Delta$  for every  $n \in \mathbb{N} \cup \{0\}$ . For the sake of simplicity, take

$$\gamma_n = d(gx_n, gx_{n+1}) \tag{3}$$

for all  $n \in \mathbb{N} \cup \{0\}$ . If there exists  $n_0 \in \mathbb{N} \cup \{0\}$  for which  $x_{n_0+1} = x_{n_0}$ , then  $fx_{n_0} = gx_{n_0+1}$  implies that  $fx_{n_0+1} = gx_{n_0+1}$ , that is,  $x_{n_0+1} \in C(f, g)$ . Now we assume that  $x_{n+1} \neq x_n$  for all  $n \in \mathbb{N} \cup \{0\}$ . As  $f$  is an  $F$ -contraction with respect to  $g$  on  $\Delta$ , so we obtain

$$\begin{aligned} F(\gamma_n) &= F(d(gx_n, gx_{n+1})) = F(d(fx_{n-1}, fx_n)) \\ &\leq F(d(gx_{n-1}, gx_n)) - \tau \\ &= F(d(fx_{n-2}, fx_{n-1})) - \tau \\ &\leq F(d(gx_{n-2}, gx_{n-1})) - 2\tau \leq \dots \\ &\leq F(d(gx_1, gx_2)) - (n-1)\tau = F(\gamma_1) - (n-1)\tau. \end{aligned}$$

That is,

$$F(\gamma_n) \leq F(\gamma_1) - (n-1)\tau.$$

On taking limit as  $n \rightarrow \infty$ , we obtain  $\lim_{n \rightarrow \infty} F(\gamma_n) = -\infty$ . Hence  $\lim_{n \rightarrow \infty} \gamma_n = 0$  by (C2). Now, by (C3), there exists  $k \in (0, 1)$  such that  $\lim_{n \rightarrow \infty} \gamma_n^k F(\gamma_n) = 0$ . Note that

$$\gamma_n^k F(\gamma_n) - \gamma_n^k F(\gamma_1) \leq \gamma_n^k (F(\gamma_1) - (n-1)\tau) - \gamma_n^k F(\gamma_1) = -\gamma_n^k (n-1)\tau \leq 0. \tag{4}$$

Taking limit as  $n \rightarrow \infty$  in (4), we have  $\lim_{n \rightarrow \infty} (n-1)\gamma_n^k = 0$ . Consequently,  $\lim_{n \rightarrow \infty} n\gamma_n^k = 0$ . Thus there exists  $n_1$  in  $\mathbb{N}$  such that  $n\gamma_n^k \leq 1$  for all  $n \geq n_1$ , that is,  $\gamma_n \leq 1/n^{1/k}$  for all  $n \geq n_1$ . Now, for integers  $m > n \geq 1$ , we obtain

$$\begin{aligned} d(gx_n, gx_m) &\leq d(gx_n, gx_{n+1}) + d(gx_{n+1}, gx_{n+2}) + \dots + d(gx_{m-1}, gx_m) \\ &< \sum_{i=n}^{\infty} \gamma_i \leq \sum_{i=n}^{\infty} \frac{1}{i^k} < \infty. \end{aligned}$$

This shows that  $\{gx_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $g(X)$ . As  $g(X)$  is complete, so there exists  $q$  in  $g(X)$  such that  $\lim_{n \rightarrow \infty} gx_n = q$ . Let  $p \in X$  be such that  $g(p) = q$ . The sequential limit comparison property implies that  $gx_{n+1} \leq q$ . As  $x_n \leq fx_n = gx_{n+1} \leq q = g(p) \leq p$  so  $(x_n, p) \in \Delta$ . Hence from (2) we have

$$F(d(gx_n, fp)) = F(d(fx_{n-1}, fp)) \leq F(d(gx_{n-1}, gp)) - \tau.$$

Since  $\lim_{n \rightarrow \infty} d(gx_{n-1}, gp) = 0$ , therefore by (C2) we have  $\lim_{n \rightarrow \infty} F(d(gx_{n-1}, gp)) = -\infty$ . Hence  $\lim_{n \rightarrow \infty} F(d(gx_n, fp)) = -\infty$  implies that  $\lim_{n \rightarrow \infty} d(gx_n, fp) = 0$ . That is,  $\lim_{n \rightarrow \infty} gx_n = fp$ . Uniqueness of limit implies  $fp = gp$ , that is,  $p \in C(f, g)$ .

(b) Now suppose that  $C(f, g)$  is well ordered. We prove that  $C(f, g)$  is a singleton. Assume on the contrary that there exists another point  $w$  in  $X$  such that  $fw = gw$  with  $w \neq p$ . Since  $C(f, g)$  is well ordered, so  $(w, p) \in \Delta$ . Now from (2) we have

$$\tau \leq F(d(gw, gp)) - F(d(fw, fp)) = 0,$$

a contradiction. Therefore  $w = p$ . Hence  $f$  and  $g$  have a unique coincidence point  $p$  in  $X$ . The converse follows immediately.

(c) Now if  $f$  and  $g$  are weakly compatible mappings, then we have  $fq = fgp = gfp = gq$ , that is,  $q$  is the coincidence point of  $f$  and  $g$ . But  $q$  is the only point of coincidence of  $f$  and  $g$ , so  $fq = gq = q$ . Hence  $q$  is the unique common fixed point of  $f$  and  $g$ .  $\square$

**Example 4** Let  $X = [0, 5]$  be endowed with usual metric and usual order. Define mappings  $f, g : X \rightarrow X$  by

$$gx = \begin{cases} 0 & \text{if } x \in [0, 3), \\ 3 & \text{if } x \in [3, 5), \\ 5 & \text{if } x = 5, \end{cases} \quad fx = \begin{cases} 3 & \text{if } x \in [0, 3), \\ 5 & \text{if } x \in [3, 5]. \end{cases}$$

Clearly,  $g$  is dominated and  $f$  is dominating. Define  $F : \mathbb{R}_+ \rightarrow \mathbb{R}$  as  $F(x) = \ln(x)$ . If  $x \in [0, 3)$  and  $y \in [3, 5)$ , then

$$\begin{aligned} F(d(fx, fy)) &= F(d(3, 5)) = F(2) = \ln(2) \approx 0.693 \\ &< F(d(gx, gy)) = F(d(0, 3)) \\ &= F(3) = \ln(3) \approx 1.098. \end{aligned}$$

Hence, for  $\tau \in (0, 0.40]$ , inequality (2) is satisfied. Similarly, for  $x \in [0, 3)$  and  $y = 5$ , we have

$$\begin{aligned} F(d(fx, fy)) &= F(d(3, 5)) = F(2) = \ln(2) \approx 0.693 \\ &< F(d(gx, gy)) = F(d(0, 5)) \\ &= F(5) = \ln(5) \approx 1.6094. \end{aligned}$$

Hence, for  $\tau \in (0, 0.9164]$ , inequality (2) is satisfied. We can take a  $\tau \in (0, 0.40]$  so that

$$\tau + F(d(fx, fy)) \leq F(d(gx, gy))$$

is satisfied for all  $x, y \in [0, 5]$ , whenever  $\min\{d(fx, fy), d(gx, gy)\} > 0$ . Hence  $f$  is an  $F$ -contraction with respect to  $g$  on  $[0, 5]$ . Hence all the conditions of Theorem 2 are satisfied. Moreover,  $x = 5$  is the coincidence point of  $f$  and  $g$ . Also note that  $f$  and  $g$  are weakly compatible and  $x = 5$  is the common fixed point of  $g$  and  $f$  as well.

Now we give a common fixed point result without imposing any type of commutativity condition for self-mappings  $f$  and  $g$  on  $X$ . Moreover, we relax the dominance conditions on  $f$  and  $g$  as well.

**Theorem 3** Let  $(X, \preceq)$  be a partially ordered set such that there exists a complete metric  $d$  on  $X$ . If self-mappings  $f$  and  $g$  on  $X$  are weakly increasing and for some  $\tau > 0$  satisfy

$$\tau + F(d(fx, gy)) \leq F(d(x, y)) \tag{5}$$

for all  $(x, y) \in \Delta$  such that  $\min\{d(fx, gy), d(x, y)\} > 0$ , then  $F(f, g) \neq \emptyset$ , provided that  $X$  has the sequential limit comparison property. Further,  $f$  and  $g$  have a unique common fixed point if and only if  $F(f, g)$  is well ordered.

*Proof* Let  $x_0$  be an arbitrary point of  $X$ . Define a sequence  $\{x_n\}_{n \in \mathbb{N}}$  in  $X$  as follows:  $x_{2n+1} = fx_{2n}$  and  $x_{2n+2} = gx_{2n+1}$ . Since  $f$  and  $g$  are weakly increasing, we have  $x_{2n+1} = fx_{2n} \leq gfx_{2n} = gx_{2n+1} = x_{2n+2}$  and  $x_{2n+2} = gx_{2n+1} \leq fgx_{2n+1} = fx_{2n+2} = x_{2n+3}$ . Hence  $(x_{2n+1}, x_{2n+2}) \in \Delta$  and  $(x_{2n+2}, x_{2n+3}) \in \Delta$  for every  $n \in \mathbb{N} \cup \{0\}$ . Now define

$$\gamma_{2n} = d(x_{2n+1}, x_{2n+2}) \tag{6}$$

for all  $n \in \mathbb{N} \cup \{0\}$ . Using (5) the following holds for every  $n \in \mathbb{N} \cup \{0\}$ :

$$\begin{aligned} F(\gamma_{2n}) &= F(d(x_{2n+1}, x_{2n+2})) = F(d(fx_{2n}, gx_{2n+1})) \\ &\leq F(d(x_{2n}, x_{2n+1})) - \tau = F(\gamma_{2n-1}) - \tau. \end{aligned}$$

Similarly,

$$\begin{aligned} F(\gamma_{2n+1}) &= F(d(x_{2n+3}, x_{2n+2})) = F(d(fx_{2n+2}, gx_{2n+1})) \\ &\leq F(d(x_{2n+1}, x_{2n+2})) - \tau = F(\gamma_{2n}) - \tau. \end{aligned}$$

Therefore, for all  $n \in \mathbb{N} \cup \{0\}$ , we have

$$\begin{aligned} F(\gamma_n) &\leq F(\gamma_{n-1}) - \tau \leq F(\gamma_{n-2}) - 2\tau \leq \dots \\ &\leq F(d(x_1, x_2)) - n\tau = F(\gamma_0) - n\tau. \end{aligned}$$

Thus

$$F(\gamma_n) \leq F(\gamma_0) - n\tau. \tag{7}$$

Taking limit as  $n \rightarrow \infty$  in (7), we get

$$\lim_{n \rightarrow \infty} F(\gamma_n) = -\infty.$$

By (C2) and (C3) we get  $\lim_{n \rightarrow \infty} \gamma_n = 0$  and  $k \in (0, 1)$  such that  $\lim_{n \rightarrow \infty} \gamma_n^k F(\gamma_n) = 0$ . Note that

$$\gamma_n^k F(\gamma_n) - \gamma_n^k F(\gamma_0) \leq \gamma_n^k (F(\gamma_0) - n\tau) - \gamma_n^k F(\gamma_0) = -\gamma_n^k n\tau \leq 0. \tag{8}$$

By taking limit as  $n \rightarrow \infty$  in (8), we get  $\lim_{n \rightarrow \infty} n\gamma_n^k = 0$ . This implies that there exists  $n_1$  such that  $n\gamma_n^k \leq 1$  for all  $n \geq n_1$ . Consequently, we obtain  $\gamma_n \leq 1/n^{1/k}$  for all  $n \geq n_1$ . Now, for integers  $m > n \geq 1$ , we have

$$d(x_n, x_m) \leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \dots + d(x_{m-1}, x_m) < \sum_{i=n}^{\infty} \gamma_i \leq \sum_{i=n}^{\infty} \frac{1}{i^k} < \infty.$$

This shows that  $\{x_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence in  $X$ , so there exists  $p$  in  $X$  such that  $\lim_{n \rightarrow \infty} x_n = p$ . As  $X$  has the sequential limit comparison property, so  $(x_n, p), (x_{2n}, p), (x_{2n+1}, p) \in \Delta$ . Therefore

$$\lim_{n \rightarrow \infty} F(d(x_{2n+1}, gp)) = \lim_{n \rightarrow \infty} F(d(x_{2n}, gp)) \leq F(d(x_{2n}, p)) - \tau.$$

Since  $\lim_{n \rightarrow \infty} d(x_{2n}, p) = 0$ , by (C2) we have  $\lim_{n \rightarrow \infty} F(d(x_{2n}, p)) = -\infty$ . This implies  $\lim_{n \rightarrow \infty} F(d(x_{2n+1}, gp)) = -\infty$ , which further implies that  $\lim_{n \rightarrow \infty} d(x_{2n+1}, gp) = 0$ . Hence  $d(p, gp) = 0$  and  $p = gp$ . Similarly, we obtain  $p = fp$ . This shows that  $p$  is a common fixed point of  $g$  and  $f$ . Now suppose that  $F(f, g)$  is well ordered. We prove that  $F(f, g)$  is a singleton. Assume on the contrary that there exists another point  $q$  in  $X$  such that  $q = fq = gq$  with  $q \neq p$ . Obviously,  $(q, p) \in \Delta$ . So, from (5) we have  $\tau \leq F(d(q, p)) - F(d(fq, gp)) = 0$ , a contradiction. Therefore  $q = p$ . Hence  $g$  and  $f$  have a unique common fixed point  $p$  in  $X$ . The converse follows immediately.  $\square$

### 3 Periodic point results in metric spaces

If  $x$  is a fixed point of the self-mapping  $f$ , then  $x$  is a fixed point of  $f^n$  for every  $n \in \mathbb{N}$ , but the converse is not true. In the sequel, we denote by  $F(f)$  the set of all fixed points of  $f$ .

**Example 5** Let  $f : [0, 1] \rightarrow [0, 1]$  be given by

$$f(x) = 1 - x.$$

Then  $f$  has a unique fixed point  $x = 1/2$ . Note that  $f^n x = x$  holds for every even natural number  $n$  and  $x$  in  $[0, 1]$ . On the other hand, define a mapping  $g : [0, \pi] \rightarrow [0, \pi]$  as

$$g(x) = \cos x.$$

Then  $g$  has the same fixed point as  $g^n$  for every  $n$ .

**Definition 10** The self-mapping  $f$  is said to have the property  $P$  if  $F(f^n) = F(f)$  for every  $n \in \mathbb{N}$ . A pair  $(f, g)$  of self-mappings is said to have the property  $Q$  if  $F(f) \cap F(g) = F(f^n) \cap F(g^n)$ .

For further details on these properties, we refer to [20, 28].

Let  $(X, d)$  be a metric space and  $f : X \rightarrow X$  be a self-mapping. The set  $O(x) = \{x, fx, \dots, f^n x, \dots\}$  is called the orbit of  $x$  [29]. A mapping  $f$  is called orbitally continuous at  $p$  if  $\lim_{n \rightarrow \infty} f^n x = p$  implies that  $\lim_{n \rightarrow \infty} f^{n+1} x = fp$ . A mapping  $f$  is orbitally continuous on  $X$  if  $f$  is orbitally continuous for all  $x \in X$ .

In this section we prove some periodic point results for self-mappings on complete metric spaces.

**Theorem 4** Let  $X$  be a nonempty set such that there exists a complete metric  $d$  on  $X$ . Suppose that  $f : X \rightarrow X$  satisfies

$$\tau + F(d(fx, f^2x)) \leq F(d(x, fx)) \tag{9}$$



for some  $\tau > 0$  and for all  $x$  in  $X$  such that  $d(fx, f^2x) > 0$ . Then  $f$  has the property  $P$  provided that  $f$  is orbitally continuous on  $X$ .

*Proof* First we show that  $F(f) \neq \emptyset$ . Let  $x_0 \in X$ . Define a sequence  $\{x_n\}_{n \in \mathbb{N}}$  in  $X$ , such that  $x_{n+1} = fx_n$ , for all  $n \in \mathbb{N} \cup \{0\}$ . Denote  $\gamma_n = d(x_n, x_{n+1})$  for all  $n \in \mathbb{N} \cup \{0\}$ . If there exists  $n_0 \in \mathbb{N} \cup \{0\}$  for which  $x_{n_0+1} = x_{n_0}$ , then  $fx_{n_0} = x_{n_0}$  and the proof is finished. Suppose that  $x_{n+1} \neq x_n$  for all  $n \in \mathbb{N} \cup \{0\}$ . Using (9), we obtain

$$\begin{aligned} F(\gamma_n) &= F(d(x_n, x_{n+1})) = F(d(fx_{n-1}, f^2x_{n-1})) \\ &\leq F(d(x_{n-1}, fx_{n-1})) - \tau = F(d(fx_{n-2}, f^2x_{n-2})) - \tau \\ &\leq F(d(x_{n-2}, fx_{n-2})) - 2\tau \leq \dots \\ &\leq F(d(x_1, x_2)) - (n-1)\tau \\ &= F(d(fx_0, f^2x_0)) - (n-1)\tau \leq F(d(x_0, x_1)) - n\tau \\ &= F(\gamma_0) - n\tau \end{aligned}$$

for every  $n \in \mathbb{N} \cup \{0\}$ . By taking limit as  $n \rightarrow \infty$  in the above inequality, we obtain that  $\lim_{n \rightarrow \infty} F(\gamma_n) = -\infty$ , which together with (C2) gives  $\lim_{n \rightarrow \infty} \gamma_n = 0$ . From (C3), there exists  $k \in (0, 1)$  such that  $\lim_{n \rightarrow \infty} \gamma_n^k F(\gamma_n) = 0$ . Note that

$$\begin{aligned} \gamma_n^k F(\gamma_n) - \gamma_n^k F(\gamma_0) &\leq \gamma_n^k (F(\gamma_0) - n\tau) - \gamma_n^k F(\gamma_0) \\ &= -\gamma_n^k n\tau \leq 0. \end{aligned}$$

On taking limit as  $n \rightarrow \infty$ , we get  $\lim_{n \rightarrow \infty} n\gamma_n^k = 0$ . Hence there exists  $n_1$  such that  $n\gamma_n^k \leq 1$  for all  $n \geq n_1$ . Consequently  $\gamma_n \leq 1/n^{1/k}$  for all  $n \geq n_1$ . Now, for integers  $m > n \geq 1$  such that

$$\begin{aligned} d(f^n x_0, f^m x_0) &= d(x_n, x_m) \leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \dots + d(x_{m-1}, x_m) \\ &< \sum_{i=n}^{\infty} \gamma_i \leq \sum_{i=n}^{\infty} \frac{1}{i^k} < \infty. \end{aligned}$$

This shows that  $\{f^n x_0\}_{n \in \mathbb{N}}$  is a Cauchy sequence. Since  $\{f^n x_0 : n \in \mathbb{N}\} \subseteq O(x_0) \subseteq X$  and  $X$  is complete, which implies that there exists  $x$  in  $X$  such that  $\lim_{n \rightarrow \infty} f^n x_0 = x$ . Since  $f$  is orbitally continuous at  $x$ , so  $x = \lim_{n \rightarrow \infty} f^n x_0 = f(\lim_{n \rightarrow \infty} f^{n-1} x_0) = fx$ . Hence  $f$  has a fixed point and  $F(f^n) = F(f)$  is true for  $n = 1$ . Now assume  $n > 1$ . Suppose on the contrary that  $u \in F(f^n)$  but  $u \notin F(f)$ , then  $d(u, fu) = \alpha > 0$ . Now consider

$$\begin{aligned} F(\alpha) &= F(d(u, fu)) = F(d(f(f^{n-1}u), f^2(f^{n-1}u))) \\ &\leq F(d(f^{n-1}u, f^n u)) - \tau \\ &\leq F(d(f^{n-2}u, f^{n-1}u)) - 2\tau \leq \dots \\ &\leq F(d(u, fu)) - n\tau. \end{aligned}$$

Thus  $F(\alpha) \leq \lim_{n \rightarrow \infty} F(d(u, fu)) - n\tau = -\infty$ . Hence  $F(\alpha) = -\infty$ . By (C2)  $\alpha = 0$ , a contradiction. So  $u \in F(f)$ . □

**Theorem 5** Let  $(X, \preceq)$  be a partially ordered set such that there exists a complete metric  $d$  on  $X$  and  $f, g$  self-mappings on  $X$ . Further assume that  $f, g$  are weakly increasing and satisfy

$$\tau + F(d(fx, gy)) \leq F(d(x, y))$$

for some  $\tau > 0$ , for all  $x, y$  in  $X$  such that  $\min\{d(fx, gy), d(x, y)\} > 0$ . Then  $f$  and  $g$  have the property  $Q$  provided that  $X$  has the sequential limit comparison property.

*Proof* By Theorem 3,  $f$  and  $g$  have a common fixed point. Suppose on the contrary that

$$u \in F(f^n) \cap F(g^n)$$

but  $u \notin F(f) \cap F(g)$ , then there are three possibilities (a)  $u \in F(f) \setminus F(g)$ , (b)  $u \in F(g) \setminus F(f)$ , (c)  $u \notin F(f)$  and  $u \notin F(g)$ . Without loss of generality, let  $u \notin F(g)$ , that is,  $d(u, gu) = \alpha > 0$ , so we get

$$\begin{aligned} F(\alpha) &= F(d(u, gu)) = F(d(f^{n-1}u, g(g^n u))) \\ &\leq F(d(f^{n-1}u, g^n u)) - \tau \\ &\leq F(d(f^{n-2}u, g^{n-1}u)) - 2\tau \leq \dots \\ &\leq F(d(u, gu)) - n\tau. \end{aligned}$$

As  $\lim_{n \rightarrow \infty} F(d(u, gu)) - n\tau = -\infty$ , so we have  $F(\alpha) = -\infty$ . By (C2)  $\alpha = 0$ , a contradiction. Hence  $u \in F(g) \cap F(f)$ .  $\square$

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

The three authors contributed equally in writing this article. They read and approved the final manuscript.

#### Author details

<sup>1</sup>Department of Mathematics and Applied Mathematics, University of Pretoria, Hatfield, Pretoria, South Africa. <sup>2</sup>Department of Mathematics, Syed Babar Ali School of Science and Engineering, Lahore University of Management Sciences, Lahore, 54792, Pakistan. <sup>3</sup>Instituto Universitario de Matemática Pura y Aplicada, Universitat Politècnica de València, Valencia, 46022, Spain.

#### Acknowledgements

The third author thanks for the support of the Ministry of Economy and Competitiveness of Spain, Grant MTM2012-37894-C02-01.

Received: 1 July 2013 Accepted: 8 October 2013 Published: 07 Nov 2013

#### References

1. Banach, S: Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales. *Fundam. Math.* **3**, 133-181 (1922)
2. Dix, JG, Karakostas, GL: A fixed point theorem for  $S$ -type operators on Banach spaces and its applications to boundary-value problems. *Nonlinear Anal.* **71**, 3872-3880 (2009)
3. Latrach, K, Taoudib, MA, Zeghal, A: Some fixed point theorems of the Schauder and the Krasnosel'skii type and application to nonlinear transport equations. *J. Differ. Equ.* **221**, 256-271 (2006)
4. Rousseau, C: Point fixe de Banach (in French). *Accromath* **5**, hiver-printemps (2010) ([www.accromath.ca](http://www.accromath.ca))
5. Arandjelović, I, Kadelburg, Z, Radenović, S: Boyd-Wong-type common fixed point results in cone metric spaces. *Appl. Math. Comput.* **217**, 7167-7171 (2011)
6. Boyd, DW, Wong, JSW: On nonlinear contractions. *Proc. Am. Math. Soc.* **20**, 458-464 (1969)

7. Huang, LG, Zhang, X: Cone metric spaces and fixed point theorems of contractive mappings. *J. Math. Anal. Appl.* **332**, 1468-1476 (2007)
8. Rakotch, E: A note on contractive mappings. *Proc. Am. Math. Soc.* **13**, 459-465 (1962)
9. Tarafdar, E: An approach to fixed-point theorems on uniform spaces. *Trans. Am. Math. Soc.* **191**, 209-225 (1974)
10. Ran, ACM, Reurings, MCB: A fixed point theorem in partially ordered sets and some application to matrix equations. *Proc. Am. Math. Soc.* **132**, 1435-1443 (2004)
11. Abbas, M, Khamsi, MA, Khan, AR: Common fixed point and invariant approximation in hyperbolic ordered metric spaces. *Fixed Point Theory Appl.* **2011**, Article ID 25 (2011). doi:10.1186/1687-1812-2011-25
12. Altun, I, Simsek, H: Some fixed point theorems on ordered metric spaces and application. *Fixed Point Theory Appl.* **2010**, Article ID 621492 (2010)
13. Amini-Harandi, A, Emami, H: A fixed point theorem for contraction type maps in partially ordered metric spaces and application to ordinary differential equations. *Nonlinear Anal.* **72**, 2238-2242 (2010)
14. Ćirić, L, Abbas, M, Saadati, R, Hussain, N: Common fixed points of almost generalized contractive mappings in ordered metric spaces. *Appl. Math. Comput.* **217**, 5784-5789 (2011)
15. Harjani, J, Sadarangani, K: Fixed point theorems for weakly contractive mappings in partially ordered sets. *Nonlinear Anal.* **71**, 3403-3410 (2009)
16. Kadelburg, Z, Pavlović, M, Radenović, S: Common fixed point theorems for ordered contractions and quasicontractions in ordered cone metric spaces. *Comput. Math. Appl.* **59**, 3148-3159 (2010)
17. Nieto, JJ, Rodríguez-López, R: Contractive mapping theorems in partially ordered sets and applications to ordinary differential equations. *Order* **22**, 223-239 (2005)
18. Samet, B: Coupled fixed point theorems for a generalized Meir-Keeler contraction in partially ordered metric spaces. *Nonlinear Anal.* **72**, 4508-4517 (2010)
19. Al-Thagafi, MA: Common fixed points and best approximation. *J. Approx. Theory* **85**, 318-320 (1996)
20. Beg, I, Abbas, M: Coincidence point and invariant approximation for mappings satisfying generalized weak contractive condition. *Fixed Point Theory Appl.* **2006**, Article ID 74503 (2006)
21. Jungck, G: Compatible mappings and common fixed points. *Int. J. Math. Math. Sci.* **9**, 771-779 (1986)
22. Jungck, G: Common fixed points for noncontinuous nonself maps on nonmetric spaces. *Far East J. Math. Sci.* **4**, 199-215 (1996)
23. Sessa, S: On a weak commutativity condition of mappings in fixed point consideration. *Publ. Inst. Math. (Belgr.)* **32**, 149-153 (1982)
24. Wardowski, D: Fixed points of new type of contractive mappings in complete metric spaces. *Fixed Point Theory Appl.* (2012). doi:10.1186/1687-1812-2012-94
25. Jungck, G: Commuting mappings and fixed points. *Am. Math. Mon.* **83**, 261-263 (1976)
26. Abbas, M, Nazir, T, Radenović, S: Common fixed points of four maps in partially ordered metric spaces. *Appl. Math. Lett.* **24**, 1520-1526 (2011)
27. Altun, I, Damjanović, B, Djorić, D: Fixed point and common fixed point theorems on ordered cone metric spaces. *Appl. Math. Lett.* **23**, 310-316 (2010)
28. Jeong, GS, Rhoades, BE: Maps for which  $F(T) = F(T^n)$ . In: *Fixed Point Theory and Applications*, vol. 6, pp. 71-105. Nova Science Publishers, New York (2007)
29. Hicks, TL, Rhoades, BE: A Banach type fixed point theorem. *Math. Jpn.* **24**(3), 327-330 (1979)

10.1186/1687-1812-2013-243

**Cite this article as:** Abbas et al.: Fixed and periodic points of generalized contractions in metric spaces. *Fixed Point Theory and Applications* 2013, **2013**:243

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](http://springeropen.com)

---