

## **INDEX**

<b>1.</b>	<b>INTRODUCTION</b>	<b>3</b>
<b>1.1.</b>	<b>CITRUS</b>	<b>3</b>
<b>1.2.</b>	<b>INTEGRATED PEST MANAGEMENT IN SPANISH CITRUS</b>	<b>4</b>
<b>1.2.1.</b>	<b>Primary citrus pests in Spain and their management</b>	<b>4</b>
<b>1.2.2.</b>	<b>Aphids as citrus pests</b>	<b>6</b>
<b>1.3.</b>	<b>APHIS SPIRAECOLA AS A CITRUS PEST</b>	<b>8</b>
<b>1.3.1.</b>	<b>Origin and geographical distribution</b>	<b>8</b>
<b>1.3.2.</b>	<b>Morphological description</b>	<b>9</b>
<b>1.3.3.</b>	<b>Life cycle</b>	<b>10</b>
<b>1.3.4.</b>	<b>Seasonal abundance</b>	<b>11</b>
<b>1.3.5.</b>	<b>Ecology</b>	<b>12</b>
<b>1.3.6.</b>	<b>Damages and economic thresholds</b>	<b>13</b>
<b>1.4.</b>	<b>INTEGRATED MANAGEMENT OF APHIS SPIRAECOLA IN CLEMENTINE CITRUS</b>	<b>15</b>
<b>1.4.1.</b>	<b>Chemical control</b>	<b>15</b>
<b>1.4.2.</b>	<b>Cultural control</b>	<b>16</b>
<b>1.4.3.</b>	<b>Biological control</b>	<b>17</b>
<b>1.4.3.1.</b>	<b>Parasitoids</b>	<b>17</b>
<b>1.4.3.2.</b>	<b>Predators</b>	<b>21</b>
<b>1.5.</b>	<b>RATIONALE AND OBJECTIVES</b>	<b>30</b>
<b>1.5.1.</b>	<b>Poor performance of parasitism</b>	<b>30</b>
<b>1.5.2.</b>	<b>Asynchrony between aphids and predators</b>	<b>31</b>
<b>1.5.3.</b>	<b>Cover management as conservation biological control of aphids in citrus</b>	<b>32</b>

**REFERENCES 34**

**2. UNTANGLING THE APHID-PARASITOID FOOD WEB IN CITRUS: CAN HYPERPARASITIDS DISRUPT BIOLOGICAL CONTROL? 53**

**2.1. INTRODUCTION 55**

**2.2. MATERIAL AND METHODS 59**

**2.2.1. Experimental site and sampling dates 59**

**2.2.2. Field sampling 59**

**2.2.2.1. Parasitism rates 59**

**2.2.2.2. Parasitoid complex 60**

**2.2.3. Molecular identification 60**

**2.2.3.1. DNA sequencing of adult specimens and design of multiplex qPCR primers 60**

**2.2.3.2. Multiplex qPCR assay of aphid mummies 65**

**2.2.4. Secondary sex ratio 66**

**2.3. RESULTS 66**

**2.3.1. Parasitism rate 66**

**2.3.2. Primary parasitoid and hyperparasitoid complex 68**

**2.3.2.1. Emergence and morphological identification assay 70**

**2.3.2.2. Quantitative PCR assay 70**

**2.3.3. Seasonal trend 74**

**2.3.4. Hyperparasitism effect on the primary parasitoid sex ratio 75**

**2.4. DISCUSSION 78**

**REFERENCES 85**

**3. EARLY ARRIVAL OF PREDATORS CONTROLS APHIS SPIRAECOLA COLONIES IN CITRUS CLEMENTINES. 97**

**3.1. INTRODUCTION 99**

**3.2. MATERIALS AND METHODS 102**

**3.2.1. Study orchards 102**

**3.2.2. Aphid colonies and predators 103**

**3.2.3. Damage and intervention thresholds 104**

**3.2.4. Data analysis 105**

**3.3. RESULTS 106**

**3.3.1. Colony parameters 106**

**3.3.1.1. Maximum number of aphids per colony 106**

**3.3.1.2. Colony survival 107**

**3.3.1.3. Colony phenology 109**

**3.3.2. Aphid predators 111**

**3.3.2.1. Abundance of aphid predators 111**

**3.3.2.2. Attack ratio and time of the first attack by aphid predators 113**

**3.3.3. Effect of the first attack by predators on colony parameters and damages 114**

**3.4. DISCUSSION 117**

**REFERENCES 123**

**4. A SOWN COVER WITH WILD PLANTS IMPROVES THE BIOLOGICAL CONTROL OF APHIDS IN CITRUS. 133**

**4.1. INTRODUCTION 135**

**4.2. MATERIAL AND METHODS 138**

4.2.1. Orchards	138
4.2.2. Ground cover sampling	139
4.2.3. Citrus canopy sampling	141
4.2.4. Statistical analysis	143
4.3. RESULTS	144
4.3.1. Ground cover characterization	144
4.3.2. Aphid community in the ground cover	145
4.3.2.1. Quantitative analysis	145
4.3.2.2. Qualitative analysis	146
4.3.3. The effect of ground cover on aphid natural enemies in the canopy	153
4.3.3.1. Ratio of attacked colonies	153
4.3.3.2. Cumulative predators and parasitoids per day	154
4.3.4. Aphid damage	156
4.4. DISCUSSION	157
REFERENCES	163
5. GENERAL DISCUSSION AND CONCLUSIONS	173
REFERENCES	179